

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 235/08, A61K 31/415, 31/40, C07D 235/14, 235/30, 209/08, 209/42, 235/12, 235/24, 235/06, 235/10, 235/26, 209/12, 401/04, 401/06</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/24771</p>
		<p>(43) International Publication Date: 11 June 1998 (11.06.98)</p>
<p>(21) International Application Number: PCT/JP97/04192</p>	<p>(74) Agent: SEKI, Hideo; Fujisawa Pharmaceutical Co., Ltd., Osaka Factory, 1-6, Kashima 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532 (JP).</p>	
<p>(22) International Filing Date: 18 November 1997 (18.11.97)</p>		
<p>(30) Priority Data: PO 3953 2 December 1996 (02.12.96) AU</p>	<p>(81) Designated States: AU, CA, CN, HU, IL, JP, KR, MX, US, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p>	
<p>(71) Applicant (for all designated States except US): FUJISAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).</p>	<p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
<p>(72) Inventors; and</p>		
<p>(75) Inventors/Applicants (for US only): SETOI, Hiroyuki [JP/JP]; 4-13-1, Namiki, Tsukuba-shi, Ibaraki 305 (JP). OHKAWA, Takehiko [JP/JP]; 944-2, Mukoishige, Ishigemachi, Yuki-gun, Ibaraki 300-27 (JP). ZENKOH, Tatsuya [JP/JP]; 6-3-3-A101, Keyakidai, Moriyamachi, Kitasouma-gun, Ibaraki 302-01 (JP). SAWADA, Hitoshi [JP/JP]; 2-25-10, Matsushiro, Tsukuba-shi, Ibaraki 305 (JP). SAWADA, Yuki [JP/JP]; 2-28-40, Kamikashiwada, Ushiku-shi, Ibaraki 300-12 (JP). OKU, Teruo [JP/JP]; 8-2, Midorigaoka, Tsukuba-shi, Ibaraki 305 (JP).</p>		

(54) Title: BENZAMIDE DERIVATIVES HAVING A VASOPRESSIN ANTAGONISTIC ACTIVITY

(57) Abstract

This invention relates to new benzamide derivatives having a vasopressin antagonistic activity, etc. and represented by general formula (I) wherein R¹ is aryl optionally substituted with lower alkoxy, etc., R² is lower alkyl, etc., R³ is hydrogen etc., A is NH, etc., E is (a), etc., X is -CH=CH-, -CH=N-, or S, and Y is a condensed heterocyclic group, etc., and pharmaceutically acceptable salts thereof, to processes for preparation thereof and to a pharmaceutical composition comprising the same.



(I)



(a)

and pharmaceutically acceptable salts thereof, to processes for preparation thereof and to a pharmaceutical composition comprising the same.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DESCRIPTION

BENZAMIDE DERIVATIVES HAVING A VASOPRESSIN ANTAGONISTIC ACTIVITY

5 TECHNICAL FIELD

This invention relates to new benzamide derivatives and pharmaceutically acceptable salts thereof which are useful as a medicament.

10 BACKGROUND ART

Some benzamide derivatives have been known as vasopressin antagonist, for example, in PCT International Publication Nos. WO 91/05549, WO 95/29152 and WO 96/41795, and EP Application Publication No. 0620216.

15

DISCLOSURE OF INVENTION

This invention relates to new benzamide derivatives and pharmaceutically acceptable salts thereof.

More particularly, it relates to new benzamide
20 derivatives and pharmaceutically acceptable salts thereof which possess activities as vasopressin antagonistic activity, vasodilating activity, hypotensive activity, activity for inhibiting saccharide release in liver, activity for inhibiting growth of mesangium cells, water diuretic
25 activity, platelet agglutination inhibitory activity, oxytocin antagonistic activity and the like, to a pharmaceutical composition comprising the salt and to a method for the treatment and/or prevention of hypertension, heart failure, renal insufficiency, edema, ascites,
30 vasopressin parasecretion syndrome, hepatocirrhosis, hyponatremia, hypokalemia, diabetic, circulation disorder, cerebrovascular disease (e.g. cerebral edema, cerebral infarction, etc.), Meniere's syndrome (e.g. Meniere's disease, etc.), motion sickness and the like in human beings
35 or animals.


One object of this invention is to provide new and useful benzamide derivatives which possess aforesaid activities.

Another object of this invention is to provide processes for the preparation of said benzamide derivatives and salts thereof.

A further object of this invention is to provide a pharmaceutical composition comprising, as an active ingredient, said benzamide derivatives and pharmaceutically acceptable salts thereof.

Still further object of this invention is to provide a therapeutical method for the treatment and/or prevention of aforesaid diseases in human beings or animals, using said benzamide derivatives and pharmaceutically acceptable salts thereof.

The object benzamide derivatives of this invention are new and can be represented by the following general formula (I) :



wherein

R^1 is aryl, cyclo(lower)alkyl or a heterocyclic group, each of which may be substituted with substituent(s) selected from the group consisting of halogen; hydroxy; nitro; protected amino; amino; acyl; substituted acyl; acyl(lower)alkylsulfinyl; acyl(lower)alkylsulfonyl; acyloxy; lower alkylamino(lower)alkylcarbamoyloxy; aryl; cyano; a heterocyclic group;

- lower alkenyl optionally substituted with acyl, substituted acyl, aryl or acyl-substituted aryl; lower alkynyl optionally substituted with amino, acylamino or substituted acylamino;
- 5 lower alkyl optionally substituted with halogen, amino, lower alkylamino, acylamino, substituted acylamino, hydroxy, acyloxy, acyl(lower)alkanoyloxy, acyl, substituted acyl, acyl(lower)alkoxyimino, aryl or acyl-substituted aryl;
- 10 lower alkylthio optionally substituted with acyl or substituted acyl; alkoxy optionally substituted with aryl, substituted aryl, hydroxy, acyloxy, amino, lower alkylamino, protected amino, a heterocyclic group, acyl-substituted
- 15 pyridyl, substituted acyl-substituted pyridyl, halogen, acyl(lower)alkylamino, N-protected-acyl(lower)-alkylamino, N-acyl(lower)alkyl-N-lower alkylamino, acyl, substituted acyl, acylamino, substituted acylamino, lower alkylhydrazinocarbonylamino, hydroxyimino,
- 20 acyl(lower)alkoxyimino, substituted acyl(lower)alkoxyimino, acyl(lower)alkoxy, guanidino or N-protected guanidino; and lower alkenyloxy optionally substituted with acyl or substituted acyl;
- R^2 is hydrogen; lower alkyl optionally substituted with
- 25 hydroxy, aryl or acyl; or cyclo(lower)alkyl;
- R^3 is hydrogen; halogen; hydroxy; acyloxy; substituted acyloxy; lower alkyl optionally substituted with hydroxy or lower alkoxy; lower alkoxy optionally substituted with aryl, amino, protected amino, acyl, hydroxy, cyano
- 30 or lower alkylthio; nitro; amino; acyl; substituted acyl; or cyclo(lower)alkyloxy;
- A is a single bond, O or NH;
- E is lower alkylene, lower alkenylene, $\text{-}\overset{\text{O}}{\parallel}\text{C-}$, $\text{-}\overset{\text{O}}{\parallel}\text{S-}$, or a group of the formula :

-G-J-

in which G is lower alkylene or $\text{-}\overset{\text{O}}{\parallel}\text{C-}$ and

5 J is O or $\text{-}\overset{\text{R}^4}{\underset{|}{\text{N}}}\text{-}$

(wherein R^4 is hydrogen or N-protective group);

X is -CH=CH- , -CH=N- or S; and

Y is aryl which may be substituted with acyl, protected
amino(lower)alkanoyl, protected amino and nitro, amino
10 and nitro or diamino;
or a condensed heterocyclic group which may be
substituted with substituent(s) selected from the group
consisting of halogen, acyl, lower alkoxy, hydroxy,
guanidino, mercapto, acylamino, amino, a heterocyclic
15 group, cyanoamino, amino(lower)alkyl(lower)alkylamino,
lower alkylamino, lower alkylamino(lower)alkylamino,
substituted-heterocyclic group, lower alkylhydrazino,
aryloxy, lower alkylthio, aryl, protected amino,
N-protected lower alkylamino(lower)alkylamino,
20 N-protected amino(lower)alkyl(N'-lower alkyl)amino,
amino(lower)alkyl(N-lower alkyl)amino, lower
alkylamino(lower)alkyl(N-lower alkyl)amino, lower
alkoxy(lower)alkylamino and lower alkyl optionally
substituted with aryl, ar(lower)alkoxy, cyano,
25 hydroxyimino, mercapto, lower alkylamino, acyloxy,
halogen, lower alkoxy, protected hydroxy, hydroxy, lower
alkoxyaryl, protected amino, amino, a heterocyclic group
or substituted heterocyclic group;

provided that when Y is phenyl which may be substituted with
30 lower alkyl or acyl,


then A is a single bond and

E is $\text{-}\overset{\text{OR}^4}{\parallel}\text{CN-}$ (wherein R^4 is as defined
above);


35 and pharmaceutically acceptable salt thereof.

The object compound (I) or its salt can be prepared by the processes as illustrated in the following reaction schemes.

5 Process 1




Process 2



Process 3


5



10

Process 4


15





20

25 Process 5


30








35

Process 6Process 7


25

Process 8

Process 9Process 10Process 11

Process 12Process 13


25

Process 14

Process 15


5

10

Process 16


15

20




Process 17



25

30




35

Process 18Process 19Process 20


Process 21Process 22

25


Process 23

Process 24

5





reduction



10

Process 25



15



20


Process 26




25



30

35

Process 27

Process 30Process 31Process 32

wherein

R^1 , R^2 , R^3 , A, E, X and Y are each as defined above,

Ea is $-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ or $-\overset{\text{O}}{\underset{\text{O}}{\text{S}}}-$,

5

Ya is indolyl,

R^5 is lower alkyl,

Z^1 is an acid residue,

Yb is N-(lower alkyl)indolyl,

10 Yc is phenyl substituted with amino and nitro,

Yd is phenyl substituted with diamino,

Ye is benzimidazolyl optionally 2-position substituted with
aryl, phenoxy, sulfamoylamino, cyanoamino, guanidino,
mercapto, amino, lower alkoxy carbonylamino, lower alkoxy
15 or lower alkyl optionally substituted with cyano,
mercapto, hydroxy, halogen, protected amino or a
heterocyclic group;

Yf is quinoxalinyll or benzotriazolyl,

Yg is N-acylindolyl,

20 Yh is (N-acyl)acylindolyl, N-acylindolyl,
(N-acyl)hydroxy(lower)alkylindolyl,
lower alkylamino(lower)alkylamino(N-acyl)indolyl,
(N-lower alkoxyarylmethyl)acylbenzimidazolyl,
(N-lower alkoxy carbonyl)phthalimido(lower)alkylindolyl,
25 N-protected lower alkylamino(lower)alkylamino(N-acyl)-
benzimidazolyl, (N-acyl)benzimidazolyl, (N-acyl)(lower)-
alkylbenzimidazolyl, N-protected amino(lower)alkyl(N-
lower alkyl)amino(N-acyl)benzimidazolyl, N-acylindolyl,
(N-acyloxymethyl)indolyl, (N-acyl)acylindolyl,
30 (N-arylmethyl)lower alkoxy(lower)alkylbenzimidazolyl or
(N-lower alkoxyarylmethyl)acylbenzimidazolyl;

Yi is acylindolyl, indolyl, hydroxy(lower)alkylindolyl,
lower alkylamino(lower)alkylaminoindolyl,
acylbenzimidazolyl, phthalimido(lower)alkylindolyl,
35 amino(lower)alkylindolyl, lower

- alkylamino(lower)alkylaminobenzimidazolyl,
benzimidazolyl, lower alkylbenzimidazolyl,
amino(lower)alkyl(N-lower alkyl)aminobenzimidazolyl,
indolyl, acylindolyl, lower alkoxy(lower)-
5 alkylbenzimidazolyl or acylbenzimidazolyl;
Yj is aryl which is substituted with protected amino and
nitro; or a condensed heterocyclic group which is
substituted with protected amino or lower alkyl
substituted with protected amino;
10 Yk is aryl which is substituted with amino and nitro; or
a condensed heterocyclic group which is substituted with
amino or lower alkyl substituted with amino;
Yl is aryl substituted with esterified carboxy, or
a condensed heterocyclic group substituted with
15 esterified carboxy,
Ym is aryl substituted with carboxy, or a condensed
heterocyclic group substituted with carboxy,
Yn is aryl or a condensed heterocyclic group, each of which
is substituted with substituted or unsubstituted
20 N-containing heterocycliccarbonyl, carbamoyl,
heterocycliccarbamoyl, or substituted or unsubstituted
lower alkylcarbamoyl;
Yo is a condensed (N-acyl)N-containing heterocyclic group or
a condensed heterocyclic group, each of which is
25 substituted with methoxy or lower alkyl substituted with
protected hydroxy;
Yp is a condensed (N-acyl)N-containing heterocyclic group or
a condensed heterocyclic group, each of which is
substituted with hydroxy or lower alkyl substituted with
30 hydroxy;
Yq is a condensed heterocyclic group which is substituted
with amino or amino(lower)alkyl,
Yr is a condensed heterocyclic group which is substituted
with acylamino or acylamino(lower)alkyl,
35 Ys is indolyl which is substituted with methyl substituted

with lower alkylamino,

Yt is a condensed heterocyclic group which is substituted
with lower alkyl substituted with hydroxy,

5 Yu is a condensed heterocyclic group which is substituted
with lower alkyl substituted with formyl,

R_a^1 is aryl substituted with esterified carboxy or lower
alkoxy substituted with esterified carboxy,

R_b^1 is aryl substituted with carboxy or lower alkoxy
substituted with carboxy,

10 R_a^2 is lower alkyl,

R_a^3 is hydrogen or lower alkoxy,

Yv is benzimidazolyl optionally substituted with lower alkyl
or protected amino(lower)alkyl,

15 R_c^1 is aryl substituted with methoxy which is substituted with
substituted or unsubstituted aryl,

R_d^1 is aryl substituted with hydroxy,

20 R_e^1 is aryl substituted with N-protected piperazinylcarbonyl,
oxopiperidinylcarbonyl, carbamoyl, lower alkylcarbamoyl,
lower alkylaminocarbamoyl or lower alkylamino(lower)-
alkyl(N-lower)alkylcarbamoyl, or aryl which is
substituted with lower alkoxy substituted with
N-protected piperazinylcarbonyl, oxopiperidinylcarbonyl,
carbamoyl, lower alkylcarbamoyl, lower
alkylaminocarbamoyl or lower alkylamino(lower)alkyl(N-
25 lower)alkylcarbamoyl,

R_f^1 is aryl which is substituted with lower alkoxy substituted
with oxopiperidinylcarbonyl,

R_g^1 is aryl which is substituted with lower alkoxy substituted
with hydroxypiperidinylcarbonyl,

30 R_h^1 is aryl substituted with acyloxy,

R_i^1 is aryl which is substituted with lower alkoxy substituted
with protected amino,

R^6 is lower alkyl substituted with protected amino,

Z^2 is an acid residue,

35 R_j^1 is aryl which is substituted with lower alkoxy substituted

- with amino,
- R_K^1 is aryl which is substituted with acylamino,
- R_L^1 is aryl which is substituted with lower alkylamino,
- R_m^1 is aryl substituted with nitro,
- 5 R_n^1 is aryl substituted with amino,
- R_{ba}^1 is aryl substituted with carboxy,
- R_{bb}^1 is aryl which is substituted with lowre alkoxy
substituted with carboxy,
- R_o^1 is aryl which is substituted with lower alkoxy substituted
10 with hydroxymethyl,
- R_p^1 is aryl which is substituted with lower alkoxy substituted
with hydroxy,
- R_q^1 is aryl which is substituted with lower alkoxy substituted
with acyloxy,
- 15 R_r^1 is aryl which is substituted with lower alkoxy substituted
with phthalimido,
- Yw is benzimidazolyl substituted with halogen,
- Yx is benzimidazolyl substituted with N-lower alkylpiperidyl,
morpholino, lower alkylamino, di(lower)alkylamino-
20 piperidino, di(lower)alkylhydrazino,
amino(lower)alkyl(N-lower alkyl)amino or
di(lower)alkylamino(lower)alkylamino,
- Yy is benzimidazolyl substituted with N-protected piperidyl,
- Yz is benzimidazolyl substituted with piperidyl,
- 25 Y_1 is benzimidazolyl or indolyl, each of which is substituted
with formyl or cyano(lower)alkyl, and
- Y_2 is benzimidazolyl or indolyl, each of which is substituted
with hydroxyiminomethyl or amino(hydroxyimino)(lower)-
alkyl.

30

In the above and subsequent description of the present
specification, suitable examples of the various definitions
to be included within the scope of the invention are
explained in detail in the following.

35

The term "lower" is intended to mean a group having 1 to 6 carbon atom(s), unless otherwise provided.

The "higher" is intended to mean 7 to 20 carbon atoms, unless otherwise provided.

5 The lower moiety in the terms "cyclo(lower)alkyl" and "cyclo(lower)alkyloxy" is intended to mean a group having 3 to 6 carbon atoms.

10 The lower moiety in the terms "lower alkenyl", "lower alkenyloxy" and "lower alkynyl" is intended to mean a group having 2 to 6 carbon atoms.

The term "alkoxy" may include lower alkoxy and higher alkoxy.

15 Suitable "lower alkoxy" and lower alkoxy moiety in the terms "ar(lower)alkoxy", "lower alkoxy(lower)alkylamino", "acyl(lower)alkoxy", "acyl(lower)alkoxyimino", "esterified carboxy(lower)alkoxyimino", "carboxy(lower)-alkoxyimino", "N-containing heterocycliccarbonyl(lower)-alkoxyimino", "carbamoyl(lower)alkoxyimino", "lower alkylcarbamoyl(lower)alkoxyimino" and "lower alkoxycarbonyl" may be straight or branched C₁-C₆ alkoxy such as methoxy, ethoxy, propoxy, isopropoxy, methylpropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, hexyloxy or the like.

20 Suitable "higher alkoxy" may be straight or branched C₇-C₂₀ alkoxy such as heptyloxy, octyloxy, nonyloxy, decyloxy, undecyloxy, dodecyloxy, tridecyloxy, tetradecyloxy, pentadecyloxy, hexadecyloxy, heptadecyloxy, octadecyloxy, nonadecyloxy, eicosyloxy, methylheptyloxy, methyldecyloxy, methylnonyloxy, methyldecyloxy, ethylheptyloxy, ethyloctyloxy, ethylnonyloxy, ethyldecyloxy or the like, in
30 which preferable one is heptyloxy.

35 Suitable "lower alkyl" and lower alkyl moiety in the terms "amino(lower)alkyl(lower)alkylamino", "lower alkylamino(lower)alkylamino", "mercapto(lower)alkyl", "lower alkylhydrazino", "lower alkylthio", "N-protected lower alkylamino(lower)alkylamino", "N-protected

amino(lower)alkyl(N'-lower alkyl)amino",
 "amino(lower)alkyl(N-lower alkyl)amino", "lower
 alkylamino(lower)alkyl(N-lower alkyl)amino, "lower
 alkoxy(lower)alkylamino", "acyl(lower)alkylsulfinyl",
 5 "acyl(lower)alkylsulfonyl", "lower
 alkylamino(lower)alkylcarbamoxyloxy", "acyl(lower)alkylamino",
 "N-protected-acyl(lower)alkylamino", "N-acyl(lower)alkyl-N-
 lower alkylamino", "lower alkylhydrazinocarbonylamino",
 "esterified carboxy(lower)alkylamino", "N-protected-
 10 esterified carboxy(lower)alkylamino", "N-esterified
 carboxy(lower)alkyl-N-lower alkylamino",
 "carboxy(lower)alkylamino",
 "N-protected-carboxy(lower)alkylamino",
 "N-carboxy(lower)alkyl-N-lower alkylamino", "lower
 15 alkylcarbamoxy", "lower alkylcarbamoxy(lower)alkoxyloxy",
 "lower alkylcarbamoxy(lower)alkoxyimino", "N-protected-
 (substituted or unsubstituted N-containing heterocyclic)-
 carbonyl(lower)alkylamino", "N-protected-carbamoyl(lower)-
 alkylamino", "N-protected-substituted or unsubstituted lower
 20 alkylcarbamoxy(lower)alkylamino:", "N-(substituted or
 unsubstituted N-containing heterocyclic)carbonyl(lower)alkyl-
 N-lower alkylamino", "N-carbamoyl(lower)alkyl-N-lower
 alkylamino", "N-lower alkylcarbamoxy-N-lower alkylamino",
 "lower alkylcarbamoxy(lower)alkoxyimino",
 25 "1-hydroxy(lower)alkyl", "1-(lower alkyl)amino(lower)alkyl",
 "mono(lower)alkylamino", "lower alkylamino(lower)alkyl",
 "acyloxy(lower)alkyl", "halo(lower)alkyl", "lower
 alkoxy(lower)alkyl", "protected hydroxy(lower)alkyl",
 "hydroxy(lower)alkyl", "ar(lower)alkyl", "protected
 30 amino(lower)alkyl", "amino(lower)alkyl",
 "a heterocyclic(lower)alkyl", "acyl(lower)alkyl",
 "di(lower)alkylamino", "lower alkylsulfonyl" and "lower
 alkylamino" may be straight or branched C₁-C₆ alkyl such as
 methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-
 35 butyl, pentyl, ethylpropyl, hexyl or the like.

Suitable "cyclo(lower)alkyl" and cyclo(lower)alkyl moiety in the term "cyclo(lower)alkyloxy" may be cyclo-(C₃-C₆)alkyl such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

5 Suitable "aryl", aryl moiety in the terms "aryloxy", "haloaryl", "arylsulfonyl", "acyl-substituted aryl", "(N-arylmethyl)lower alkoxy(lower)alkylbenzimidazolyl" and "(N-lower alkoxyarylmethyl)acylbenzimidazolyl" and ar moiety
10 in the terms "ar(lower)alkyl" and "ar(lower)alkoxy" may be phenyl, naphthyl, phenyl substituted with lower alkyl [e.g. tolyl, xylyl, mesityl, cumenyl, di(tert-butyl)phenyl, etc.] and the like, in which preferable one is phenyl, tolyl or xylyl.

15 Suitable "substituted aryl" may be aryl substituted with suitable substituent(s) such as acyl, substituted acyl, N-protected piperazinylsulfonyl, piperazinylsulfonyl, N-lower alkylpiperazinylsulfonyl, hydroxy(lower)alkyl, a heterocyclic(lower)alkyl, halogen, nitro, amino, lower
20 alkylamino, a heterocyclic group [e.g. thiazolyl, oxazolyl, tetrazolyl, oxazolinyl, pyridyl, pyrimidinyl, pyrrolyl optionally substituted with lower alkyl and cyano, etc.], cyano, lower alkoxy or the like, in which preferable one for the substituent of alkoxy for R¹ is aryl substituted with N-lower alkylpiperazinylcarbonyl.

25 Suitable "halogen" and halo moiety in the terms "halo(lower)alkyl" and "haloaryl" may be fluorine, chlorine, bromine and iodine, in which preferable one is chlorine or bromine.

30 Suitable "lower alkylamino" and lower alkylamino moiety in the terms "amino(lower)alkyl(lower)alkylamino", "lower alkylamino(lower)alkylamino", "N-protected lower alkylamino(lower)alkylamino", "N-protected amino(lower) (N'-lower alkyl)amino", "amino(lower)alkyl(N-lower alkyl)amino", "lower alkylamino(lower)alkyl(N-lower alkyl)amino", "lower
35 alkoxy(lower)alkylamino", "lower

alkylamino(lower)alkylcarbamoxyloxy", "acyl(lower)alkylamino",
"esterified carboxy(lower)alkylamino",
"carboxy(lower)alkylamino",
"N-containing heterocycliccarbonyl(lower)alkylamino",
5 "carbamoxy(lower)alkylamino",
"lower alkylcarbamoxy(lower)alkylamino",
"lower alkylamino(lower)alkyl" and "lower
alkylaminopiperidylcarbonyl" may be mono or di(lower
alkyl)amino such as methylamino, ethylamino, propylamino,
10 isopropylamino, butylamino, tert-butylamino, isobutylamino,
pentylamino, hexylamino, dimethylamino, diethylamino,
dipropylamino, dibutylamino, diisopropylamino, dipentylamino,
dihexylamino, N-methylethylamino or the like, in which
preferable one is methylamino, dimethylamino or diethylamino.

15 Suitable "lower alkylhydrazino" may be 2-mono or
2,2-di(lower alkyl)hydrazino such as 2-methylhydrazino,
2,2-dimethylhydrazino, 2-ethylhydrazino, 2,2-diethylhydrazino
or the like, in which preferable one 2,2-dimethylhydrazino.

20 Suitable "1-hydroxy(lower)alkyl" may be 1-hydroxy-
(C₁-C₆)alkyl such as hydroxymethyl, 1-hydroxyethyl,
1-hydroxypropyl, 1-hydroxybutyl, 1-hydroxy-3-methylpropyl or
the like, in which preferable one is hydroxymethyl or
1-hydroxyethyl.

25 Suitable "1-(lower alkyl)amino(lower)alkyl" may be
1-mono or di(C₁-C₆ alkyl)amino(C₁-C₆)alkyl such as
methylaminomethyl, dimethylaminomethyl, 1-methylaminoethyl,
1-dimethylaminoethyl, ethylaminomethyl, 1-ethylaminoethyl or
the like, in which preferable one is methylaminomethyl,
dimethylaminomethyl, 1-methylaminoethyl or
30 1-dimethylaminoethyl.

Suitable "lower alkylamino(lower)alkyl" may be mono or
di(lower alkyl)amino(lower)alkyl such as methylaminomethyl,
dimethylaminomethyl, dimethylaminoethyl or the like.

35 Suitable "amino(lower)alkyl(lower)alkylamino" may be
aminomethylmethylamino, aminomethylethylamino,

aminoethylmethyamino, aminoethylethylamino and the like, in which preferable one is aminoethylmethyamino.

Suitable "lower alkylamino(lower)alkylamino" may be mono or di(lower alkyl)amino(lower)alkylamino such as
5 methylethylamino, dimethylethylamino and the like.

Suitable "N-protected lower alkylamino(lower)alkylamino" may be N-tert-butoxycarbonyl(lower)alkylamino(lower)-alkylamino such as N-tert-butoxycarbonylmethylaminoethylamino or the like.

10 Suitable "N-protected amino(lower)alkyl(N'-lower alkyl)amino" may be N-tert-butoxycarbonylamino(lower)alkyl-(N'-lower alkyl)amino such as N-tert-butoxycarbonylaminoethyl(N-methyl)amino or the like.

Suitable "amino(lower)alkyl(N-lower alkyl)amino" may be
15 aminoethyl(N-methyl)amino or the like.

Suitable "lower alkylamino(lower)alkyl(N-lower alkyl)-amino" may be mono or di(lower alkyl)amino(lower)alkyl(N-lower alkyl)amino such as dimethylaminoethyl(N-methyl)amino or the like.

20 Suitable "lower alkoxy(lower)alkylamino" may be methoxyethylamino and the like.

Suitable "acyloxy(lower)alkyl" may be pivaloyloxymethyl and the like.

Suitable "lower alkoxy(lower)alkyl" may be methoxymethyl
25 and the like.

"hydroxy-protective group" in protected hydroxy moiety in the term "protected hydroxy(lower)alkyl" may be common hydroxy-protective group such as substituted or unsubstituted arylmethyl (e.g. benzyl, lower alkoxybenzyl, etc.), acyl,
30 substituted silyl (e.g. tert-butyldiphenylsilyl, etc.) or the like.

Suitable "heterocyclic group" may be one containing at least one hetero atom selected from nitrogen, sulfur and oxygen atom, and may include saturated or unsaturated,
35 monocyclic or condensed heterocyclic group, and preferable

heterocyclic group may be N-containing heterocyclic group such as unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl [e.g. 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.], tetrazolyl [e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.], etc.;

saturated 3 to 7-membered heteromonocyclic group containing 1 to 4 nitrogen atoms [e.g. pyrrolidinyl, imidazolidinyl, piperidyl, piperazinyl, homopiperazinyl, etc.];

saturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, quinuclidinyl, etc.

unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, imidazopyridyl [e.g. imidazo[4,5-b]pyridyl, imidazo[1,2-a]pyridyl, imidazo[3,4-a]pyridyl, etc.], purinyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl [e.g. tetrazolo[1,5-b]pyridazinyl, etc.], indolinyl, tetrahydroquinolyl, quinoxalinyl, 1H-indazolyl, 1H-pyrazolo[1,5-b][1,2,4]-triazolyl, quinazolinyl, etc.;

unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.;

saturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, 1H-tetrahydropyranyl, tetrahydrofuranlyl, etc.;

unsaturated, 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms, for example, thienyl, etc.;

unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl [e.g. 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.], oxazolinyl [e.g. 2,5-oxazolinyl, etc.], oxazinyl [e.g. 3H,4H,5H-2,6-oxazinyl, etc.], etc.;

saturated 3 to 6-membered heteromonocyclic group containing 1

to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. morpholinyl, etc.];
unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. benzofurazanyl, benzoxazolyl, benzoxadiazolyl, benzoxazinyl, etc.];
5 unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl [e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.], etc.;
10 saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., thiazolidinyl, etc.];
unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., benzothiazolyl, benzothiadiazolyl, etc.];
15 unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms [e.g. benzofuranyl, benzodioxolyl, etc.] and the like.

Said "heterocyclic group" includes one substituted with lower alkyl as exemplified above or oxo, and spiro-typed one substituted with C₂-C₆ alkylene, in which preferable one is N-methylpiperazinyl, tetrazolyl, morpholinyl, pyrrolidinyl, N-methylpiperidyl, N-methylhomopiperazinyl, 1H-tetrahydropyranyl, thienyl, pyridyl, piperidyl, 20 oxopiperidyl, pyrrolyl, oxazolyl, 2,5-oxazolinyl, 4,4-dimethyl(2,5-oxazolinyl), 1-aza-3-oxaspiro[4.4]non-1-en-2-yl, 3H,4H,5H-2,6-oxazinyl.

Suitable "condensed heterocyclic group" may be saturated or unsaturated one above-mentioned, in which preferable one is indolyl, benzimidazolyl, benzoxazolyl, benzotriazolyl, imidazopyridyl (e.g. imidazo[4,5-b]pyridyl, imidazo[1,2-a]pyridyl, imidazo[3,4-a]pyridyl, etc.), purinyl, indolinyl, tetrahydroquinolyl, quinoxalinyl, 1H-indazolyl, 1H-pyrazolo[1,5-b][1,2,4]triazolyl, quinazolinyl, 2H-1,4-benzoxazin-3-oxo-8-yl.
35

Suitable acyl and acyl moiety in the terms
"acyl(lower)alkylsulfinyl", "acyl(lower)alkylsulfonyl",
"acyloxy", "acyloxy(lower)alkyl", "acylamino",
"acyl(lower)alkanoyloxy", "acyl(lower)alkoxyimino",
5 "acyl(lower)alkylamino", "N-protected-acyl(lower)alkylamino",
"N-acyl(lower)alkyl-N-lower alkylamino" and
"acyl(lower)alkoxy" may be carboxy, esterified carboxy,
carbamoyl, lower alkylcarbamoyl, lower alkanoyl, aroyl, a
heterocycliccarbonyl, lower alkylsulfonyl, arylsulfonyl,
10 sulfamoyl, lower alkylsulfamoyl and the like.

The esterified carboxy may be substituted or
unsubstituted lower alkoxycarbonyl [e.g. methoxycarbonyl,
ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl,
tert-butoxycarbonyl, hexyloxycarbonyl, 2-iodoethoxycarbonyl,
15 2,2,2-trichloroethoxycarbonyl, dimethylaminopropoxycarbonyl,
dimethylaminoethoxycarbonyl, etc.], substituted or
unsubstituted aryloxycarbonyl [e.g. phenoxycarbonyl,
4-nitrophenoxycarbonyl, 2-naphthyloxycarbonyl, etc.],
substituted or unsubstituted ar(lower)alkoxycarbonyl [e.g.
20 benzyloxycarbonyl, phenethyloxycarbonyl,
benzhydryloxycarbonyl, 4-nitrobenzyloxycarbonyl,
3-methoxy-4-nitrobenzyloxycarbonyl, etc.], N-containing
heterocyclicoxycarbonyl [e.g. N-methylpiperidyloxycarbonyl,
etc.] and the like.

25 The lower alkylcarbamoyl may be mono or di(lower alkyl)-
carbamoyl such as methylcarbamoyl, ethylcarbamoyl,
propylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl,
N-methyl-N-ethylcarbamoyl or the like.

The lower alkanoyl may be substituted or unsubstituted
30 C₁-C₆ alkanoyl such as formyl, acetyl, propionyl, butyryl,
isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl,
trifluoroacetyl or the like, in which preferable one is
formyl or acetyl.

The aroyl may be benzoyl, naphthoyl, toluoyl, di(tert-
35 butyl)benzoyl and the like, in which preferable one is

benzoyl.

The heterocyclic moiety in the terms "a heterocyclic-carbonyl", "heterocyclicoxycarbonylamino", "heterocycliccarbamoyl" and "heterocyclicsulfonyl" may be one
5 mentioned above as a heterocyclic group.

Preferred "a heterocycliccarbonyl" may be N-containing heterocycliccarbonyl.

The "N-containing heterocycliccarbonyl" may be one containing at least one nitrogen atom in heterocyclic group
10 mentioned above, in which preferable one is N-(lower alkyl)-piperazinylcarbonyl (e.g. N-methylpiperazinylcarbonyl, etc.), N-(lower alkyl)homopiperazinylcarbonyl (e.g. N-methylhomopiperazinylcarbonyl, etc.), piperazinylcarbonyl, pyrrolidinylcarbonyl, piperidylcarbonyl, morpholinocarbonyl,
15 lower alkylpiperidylcarbonyl (e.g. methylpiperidylcarbonyl, etc.) or oxopiperidylcarbonyl.

Suitable "substituted acyl" may be carbamoyl substituted with amino, a heterocyclic group [e.g. N-(lower alkyl)piperazinyl, pyridyl, etc.], lower alkylsulfonyl or
20 arylsulfonyl, substituted lower alkylcarbamoyl [e.g. N-lower alkylamino-N-lower alkylcarbamoyl, pyridyl(lower)alkylcarbamoyl, morpholino(lower)alkylcarbamoyl, bis[hydroxy(lower)alkyl]carbamoyl, hydroxy(lower)alkylcarbamoyl, carbamoyl(lower)alkylcarbamoyl, lower alkylamino(lower)alkylcarbamoyl, N-lower alkyl-N-lower alkylcarbamoyl, etc.], substituted N-containing heterocycliccarbonyl [e.g. trifluoroacetyl-piperazinylcarbonyl,
25 pyridylpiperazinylcarbonyl, hydroxypiperidylcarbonyl, dimethylaminopiperidylcarbonyl, diethylaminopiperidylcarbonyl, carbamoylpyrrolidinylcarbonyl, dimethylaminopiperazinylcarbonyl, hydroxyethoxyethylpiperazinylcarbonyl,
30 pyrrolidinylcarbonylmethylpiperazinylcarbonyl, etc.],

N-protected-N-containing heterocycliccarbonyl [e.g. N-t-butoxycarbonylpiperidylcarbonyl, N-t-butoxycarbonylpiperazinylcarbonyl, etc.], N-protected amino(lower)alkanoyl, amino(lower)alkanoyl, benzyloxybenzoyl, and the like.

"N-Protective group" in "protected amino" may be common N-protective group such as substituted or unsubstituted lower alkanoyl [e.g. formyl, acetyl, propionyl, trifluoroacetyl, etc.], phthaloyl, lower alkoxycarbonyl [e.g. tert-butoxycarbonyl, tert-amylloxycarbonyl, etc.], substituted or unsubstituted aralkyloxycarbonyl [e.g. benzyloxycarbonyl, p-nitrobenzyloxycarbonyl, etc.], 9-fluorenylmethoxycarbonyl, substituted or unsubstituted arenesulfonyl [e.g. benzenesulfonyl, tosyl, etc.], nitrophenylsulfonyl, aralkyl [e.g. trityl, benzyl, etc.] or the like, in which preferable one is phthaloyl, tert-butoxycarbonyl or 9-fluorenylmethoxycarbonyl.

"N-protective group" in "N-protected guanidino" may be common N-protective group such as lower alkoxycarbonyl [e.g. tert-butoxycarbonyl, etc.] or the like.

Suitable "acid residue" may be halogen [e.g. fluoro, chloro, bromo, iodo], arenesulfonyloxy [e.g. benzenesulfonyloxy, tosyloxy, etc.], alkanesulfonyloxy [e.g. mesyloxy, ethanesulfonyloxy, etc.], and the like, in which preferable one is halogen.

Suitable "lower alkylsulfonyl" may be (C₁-C₆)-alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, propylsulfonyl or the like, in which preferable one is methylsulfonyl.

Suitable "arylsulfonyl" may be phenylsulfonyl, tolylsulfonyl and the like.

The substituent(s) on aryl for R¹ or a condensed heterocyclic group for Y, and the substituent(s) on lower alkyl as substituent of a condensed heterocyclic group for Y may be plural and in such case the substituents may be the

same or different.

Preferred "aryl" for R^1 may be phenyl or phenyl substituted with lower alkyl.

Preferred "a heterocyclic group" as substituent of aryl
5 for R^1 may be piperidino, N-lower alkylpiperazinyl [e.g. N-methylpiperazinyl, etc.], morpholino, 4,4-dimethyl(2,5-oxazoliny), pyrrolyl, 2,5-oxazolyl, 2,5-oxazoliny, 3H,4H,5H-2,6-oxaziny or 1-aza-3-oxaspiro[4.4]non-1-en-2-yl.

Preferred "a heterocyclic group" in a heterocyclic-
10 (lower)alkyl as substituent of a condensed heterocyclic group for Y may be pyridyl, N-lower alkylpiperazinyl [e.g. N-methylpiperazinyl, etc.], morpholino, imidazolyl, pyrrolidinyl.

Preferred "substituted-heterocyclic group" in
15 substituted heterocyclic(lower)alkyl as substituent of a condensed heterocyclic group for Y may be substituted-piperidyl such as lower alkylaminopiperidyl including mono or di(lower alkyl)aminopiperidyl [e.g. dimethylaminopiperidyl, etc.] or the like.

20 Preferred compound (I) is one having tolyl which is substituted with lower alkoxy substituted with N-(lower alkyl)piperazinylcarbonyl for R^1 , lower alkyl for R^2 , lower alkoxy for R^3 , NH for A and $\overset{\text{O}}{\underset{\text{||}}{\text{C}}}$ for E, or a single bond for A

25 $\overset{\text{O}}{\underset{\text{||}}{\text{C}}}\text{NH}-$ for E, $-\text{CH}=\text{CH}-$ for X and benzimidazolyl which is substituted with lower alkyl optionally substituted with hydroxy, amino or N-lower alkyl piperazinyl for Y.

Suitable pharmaceutically acceptable salts of the object
30 compound (I) are conventional non-toxic salts and include an acid addition salt such as an inorganic acid addition salt [e.g. hydrochloride, hydrobromide, sulfate, phosphate, etc.], an organic acid addition salt [e.g. formate, acetate, trifluoroacetate, maleate, tartrate, methanesulfonate,
35 benzenesulfonate, toluenesulfonate, etc.], a metal salt such

as an alkali metal salt [e.g. sodium salt, potassium salt, etc.] and an alkaline earth metal salt [e.g. calcium salt, magnesium salt, etc.] and the like.

- 5 The processes for preparing the object compound (I) are explained in detail in the following.

Process 1

10 The object compound (Ia) or its salt can be prepared by reacting a compound (II) or its salt with a compound (III) or its reactive derivative at the carboxy group or the sulfo group, or a salt thereof.

Suitable salts of the compounds (Ia) and (II) may be the same as those exemplified for the compound (I).

- 15 Suitable salts of the compound (III) and its reactive derivative at the carboxy group or the sulfo group may be base salts as exemplified for the compound (I).

Suitable reactive derivative at the carboxy group or the sulfo group of the compound (III) may include an acid halide, an acid anhydride containing intramolecular, intermolecular and a mixed ones, an activated amide, an activated ester, and the like. Suitable examples of the reactive derivatives may be an acid chloride; an acid azide; a mixed acid anhydride with an acid such as substituted phosphoric acid [e.g. dialkylphosphoric acid, phenylphosphoric acid, diphenylphosphoric acid, dibenzylphosphoric acid, halogenated phosphoric acid, etc.], dialkylphosphorous acid, sulfurous acid, thiosulfuric acid, sulfuric acid, sulfonic acid [e.g. methanesulfonic acid, etc.], aliphatic carboxylic acid [e.g. acetic acid, propionic acid, butyric acid, isobutyric acid, pivalic acid, pentanoic acid, isopentanoic acid, 2-ethylbutyric acid, trichloroacetic acid, etc.] or aromatic carboxylic acid [e.g. benzoic acid, etc.]; a symmetrical acid anhydride; an activated amide with imidazole, 4-substituted imidazole, dimethylpyrazole, triazole or tetrazole; or

20
25
30
35

an activated ester [e.g. ethyl ester, cyanomethyl ester, methoxymethyl ester, dimethyliminomethyl $[(\text{CH}_3)_2\text{N}^+=\text{CH}-]$ ester, intramolecular trifluoromethyl-substituted iminomethyl ester, vinyl ester, propargyl ester, p-nitrophenyl ester, 2,4-dinitrophenyl ester, trichlorophenyl ester, pentachlorophenyl ester, mesylphenyl ester, phenylazophenyl ester, phenyl thioester, p-nitrophenyl thioester, p-cresyl thioester, carboxymethyl thioester, pyranyl ester, pyridyl ester, piperidyl ester, 8-quinolyl thioester, etc.] or
an ester with an N-hydroxy compound [e.g. N,N-dimethylhydroxylamine, 1-hydroxy-2-(1H)-pyridone, N-hydroxysuccinimide, N-hydroxyphthalimide, 1-hydroxy-1H-benzotriazole, etc.], and the like. These reactive derivatives can optionally be selected from them according to the kind of the compound (III) to be used.

The reaction is usually carried out in a conventional solvent such as water, alcohol [e.g. methanol, ethanol, etc.], acetone, dioxane, acetonitrile, chloroform, methylene chloride, ethylene chloride, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide, pyridine or any other organic solvent which does not adversely influence the reaction. These conventional solvents may also be used in a mixture with water.

In this reaction, when the compound (III) is used in a free acid form or its salt form, the reaction is preferably carried out in the presence of a conventional condensing agent such as N,N'-dicyclohexylcarbodiimide; N-cyclohexyl-N'-morpholinoethylcarbodiimide; N-cyclohexyl-N'-(4-diethylaminocyclohexyl)carbodiimide; N,N'-diethylcarbodiimide, N,N'-diisopropylcarbodiimide; N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide; N,N'-carbonylbis-(2-methylimidazole); pentamethyleneketene-N-cyclohexylimine; diphenylketene-N-cyclohexylimine; ethoxyacetylene; 1-alkoxy-1-chloroethylene; trialkyl phosphite; ethyl

polyphosphate; isopropyl polyphosphate; phosphorus
oxychloride (phosphoryl chloride); phosphorus trichloride;
diphenylphosphoryl azide; diphenyl chlorophosphate;
diphenylphosphinic chloride; thionyl chloride; oxalyl
5 chloride; lower alkyl haloformate [e.g. ethyl chloroformate,
isopropyl chloroformate, etc.]; triphenylphosphine;
2-ethyl-7-hydroxybenzisoxazolium salt;
2-ethyl-5-(m-sulfophenyl)isoxazolium hydroxide intramolecular
salt; 1-(p-chlorobenzenesulfonyloxy)-6-chloro-1H-
10 benzotriazole; so-called Vilsmeier reagent prepared by the
reaction of N,N-dimethylformamide with thionyl chloride,
phosgene, trichloromethyl chloroformate, phosphorus
oxychloride, etc.; or the like.

The reaction may also be carried out in the presence of
15 an inorganic or organic base such as an alkali metal
bicarbonate, tri(lower)alkylamine, pyridine,
4-dimethylaminopyridine, N-(lower)alkylmorpholine,
N,N-di(lower)alkylbenzylamine, or the like.

The reaction temperature is not critical, and the
20 reaction is usually carried out under cooling to heating.

In this reaction, in case that the compound (II) having
aryl substituted with phthalimido for R¹, the compound (Ia)
having aryl substituted with amino may be obtained according
to reaction condition.

25 This case is included within the scope of this reaction.

Process 2

The object compound (Ic) or its salt can be prepared by
reacting a compound (Ib) or its salt with a compound (IV) in
30 the presence of a base.

Suitable salts of the compounds (Ib) and (Ic) may be the
same as those exemplified for the compound (I).

Suitable base may be an alkali metal (e.g. sodium,
potassium, etc.), an alkali metal hydride (e.g. sodium
35 hydride), an alkali metal alkoxide (e.g. potassium tert-

butoxide, etc.) and the like.

The reaction is carried out in a solvent such as N,N-dimethylformamide, tetrahydrofuran, dioxane, a mixture thereof or any other solvent which does not adversely
5 influence the reaction.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process 3

10 The object compound (Ie) or its salt can be prepared by subjecting a compound (Id) or its salt to reduction.

Suitable salts of the compounds (Id) and (Ie) may be the same as those exemplified for the compound (I).

The reduction may include chemical reduction and
15 catalytic reduction, which are carried out in a conventional manner.

Suitable reducing agents to be used in chemical reduction are a metal [e.g. tin, zinc, iron, nickel, etc.], a combination of such metal and/or metallic compound [e.g.
20 nickel chloride, chromium chloride, chromium acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.], a combination of such metal and/or metallic compound and base
25 [e.g. ammonia, ammonium chloride, sodium hydroxide, etc.], a metal hydride compound such as aluminum hydride compound [e.g. lithium aluminum hydride, sodium aluminum hydride, aluminum hydride, lithium trimethoxyaluminum hydride, lithium tri-t-butoxyaluminum hydride, etc.], borohydride compound
30 [e.g. sodium borohydride, lithium borohydride, sodium cyanoborohydride, tetramethylammonium borohydride, borane, diborane, etc.], a phosphorus compound [e.g. phosphorus trichloride, phosphorus tribromide, triphenylphosphine, triethylphosphine, etc.] and the like.

35 Suitable catalysts to be used in catalytic reduction are

conventional ones such as platinum catalysts [e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalyst [e.g. spongy palladium, palladium black, palladium oxide, palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalyst [e.g. reduced nickel, nickel oxide, Raney nickel, etc.], cobalt catalyst [e.g. reduced cobalt, Raney cobalt, etc.], iron catalyst [e.g. reduced iron, Raney iron, etc.], copper catalyst [e.g. reduced copper, Raney copper, Ullman copper, etc.], or the like.

The reduction is usually carried out in a solvent. A suitable solvent to be used may be water, an alcohol [e.g. methanol, ethanol, propanol, etc.], acetonitrile or any other conventional organic solvent such as diethyl ether, dioxane, tetrahydrofuran, etc. or a mixture thereof.

The reaction temperature is not critical, and the reaction is preferably carried out under cooling to heating.

20 Process 4

The object compound (If) or its salt can be prepared by reacting a compound (Ie) or its salt with aroyl halide, cyano(lower)alkylcarboxylic acid, mercapto(lower)alkylcarboxylic acid, lower alkyl lactone, 1,1-dihalo-1,1-diphenoxymethane, diphenyl N-sulfamoylcarbonimidate, diphenyl N-cyanocarbonimidate, dicyandiamide, 1,1'-thiocarbonyl-diimidazole, cyanogen bromide, lower alkoxycarbonyl isothiocyanate, tri(lower)alkyl orthoformate, tetra(lower)alkyl orthocarbonate, lower alkylcarboxylic acid, halo(lower)alkylcarboxylic acid, protected amino(lower)-alkylcarbonyl halide or a heterocyclic(lower)alkylcarbonyl halide.

Suitable salts of the compounds (Ie) and (If) may be the same as those exemplified for the compound (I).

35 The reaction is carried out in no solvent or a solvent

such as water, hydrochloric acid, tetrahydrofuran, ethyl acetate, acetonitrile, benzene, acetic acid, dichloromethane, pyridine, an alcohol (e.g. methanol, ethanol, isopropanol, etc.), a mixture thereof or any other solvent which does not adversely influence the reaction.

The reaction is also preferably carried out in the presence of base (e.g. sodium carbonate, etc.) or a conventional condensing agent such as N,N'-dicyclohexylcarbodiimide, p-toluenesulfonic acid, or the like.

The reaction temperature of this reaction is not critical and the reaction is usually carried out under cooling to heating.

Process 5

The object compound (Ig) or its salt can be prepared by reacting a compound (Ie) or its salt with glyoxal and sodium hydrogen sulfite, or sodium nitrite.

Suitable salts of the compounds (Ie) and (Ig) may be the same as those exemplified for the compound (I).

The reaction is usually carried out in a solvent such as water, acetic acid, an alcohol (e.g. methanol, ethanol, etc.), a mixture thereof or any other solvent which does not adversely influence the reaction.

The reaction temperature of this reaction is not critical and the reaction is usually carried out under cooling to heating.

Process 6

The object compound (Ih) or its salt can be prepared by reacting a compound (Ib) or its salt with an acylating agent.

Suitable salts of the compounds (Ib) and (Ih) may be the same as those exemplified for the compound (I).

The acylating agent may include an organic acid represented by the formula : R^7-OH , in which R^7 is acyl as illustrated above, or its reactive derivative.

The suitable reactive derivative of organic acid may be a conventional one such as an acid halide [e.g. acid chloride, acid bromide, etc.], an acid azide, an acid anhydride containing intramolecular and intermolecular ones,
5 an activated amide, an activated ester or the like.

When free acid is used as an acylating agent, the acylation reaction may preferably be conducted in the presence of a conventional condensing agent such as N,N'-dicyclohexylcarbodiimide or the like.

10 The reaction is usually carried out in a conventional solvent such as water, pyridine, acetone, dioxane, chloroform, methylene chloride, acetonitrile, ethylene chloride, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide, pyridine or any other organic solvent which does
15 not adversely influence the reaction, or a mixture thereof.

The reaction is also preferably carried out in the presence of a conventional base such as triethylamine, pyridine, N,N-dimethylaminopyridine, sodium hydroxide or the like.

20 The reaction temperature is not critical, and the reaction can be carried out under cooling to heating.

Process 7

The object compound (Ij) or its salt can be prepared by subjecting a compound (Ii) or its salt to elimination
25 reaction of the N-substituent group.

Suitable salts of the compounds (Ii) and (Ij) may be the same as those exemplified for the compound (I).

The reaction is carried out in accordance with a conventional method such as hydrolysis, reduction or the like.

30 In case that the N-substituent group is acyl, acyloxymethyl or lower alkoxyarylmethyl, the reaction is preferably carried out in accordance with hydrolysis, and in case that the N-substituent is arylmethyl, the reaction is preferably carried out in accordance with reduction.

35 The hydrolysis is preferably carried out in the presence

of a base or an acid including Lewis acid.

Suitable base may include an inorganic base and an organic base such as an alkali metal [e.g. sodium, potassium, etc.], an alkaline earth metal [e.g. magnesium, calcium, etc.], the hydroxide or carbonate or bicarbonate or lower alkoxide thereof, hydrazine, alkylamine [e.g. methylamine, trimethylamine, triethylamine, etc.], picoline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, 1,8-diazabicyclo[5.4.0]undec-7-ene, or the like.

Suitable acid may include an organic acid [e.g. formic acid, acetic acid, propionic acid, trichloroacetic acid, trifluoroacetic acid, etc.], an inorganic acid [e.g. hydrochloric acid, hydrobromic acid, sulfuric acid, hydrogen chloride, hydrogen bromide, hydrogen fluoride, etc.] and an acid addition salt compound [e.g. pyridine hydrochloride, etc.].

The elimination using trihaloacetic acid [e.g. trichloroacetic acid, trifluoroacetic acid, etc.] or the like is preferably carried out in the presence of cation trapping agents [e.g. anisole, phenol, etc.].

The reaction is usually carried out in a solvent such as water, an alcohol [e.g. methanol, ethanol, etc.], methylene chloride, chloroform, tetrachloromethane, tetrahydrofuran, a mixture thereof or any other solvent which does not adversely influence the reaction. A liquid base or acid can be also used as the solvent. The reaction temperature is not critical and the reaction is usually carried out under cooling to heating.

The reduction method applicable for the elimination reaction may include chemical reduction and catalytic reduction.

Suitable reducing agents to be used in chemical reduction are a combination of metal [e.g. tin, zinc, iron, etc.] or metallic compound [e.g. chromium chloride, chromium

acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.).

5 Suitable catalysts to be used in catalytic reduction are conventional ones such as platinum catalysts [e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalysts [e.g. spongy palladium, palladium black, palladium oxide, 10 palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalysts [e.g. reduced nickel, nickel oxide, Raney nickel, etc.], cobalt catalysts [e.g. reduced cobalt, Raney cobalt, etc.], iron catalysts [e.g. reduced iron, Raney iron, etc.], 15 copper catalysts [e.g. reduced copper, Raney copper, Ullman copper, etc.] and the like.

 In case that the N-substituent group is benzyl, the reduction is preferably carried out in the presence of a combination of palladium catalysts [e.g. palladium black, 20 palladium on carbon, etc.] and formic acid or its salt [e.g. ammonium formate, etc.].

 The reduction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, methanol, ethanol, propanol, N,N-dimethylformamide, 25 or a mixture thereof. Additionally, in case that the above-mentioned acids to be used in chemical reduction are in liquid, they can also be used as a solvent. Further, a suitable solvent to be used in catalytic reduction may be the above-mentioned solvent, and other conventional solvent such as diethyl ether, dioxane, tetrahydrofuran, etc. or a mixture 30 thereof.

 The reaction temperature of this reduction is not critical and the reaction is usually carried out under cooling to heating.

In this reaction, in case that the compound (Ii) having (N-lower alkoxy carbonyl)phthalimido(lower)alkylindolyl, (N-lower alkoxy carbonyl)lower alkylamino(lower)alkylamino-(N-lower alkoxy carbonyl)benzimidazolyl or N-lower alkoxy carbonylamino(lower)alkyl(N-lower alkyl)amino(N-lower alkoxy carbonyl)benzimidazolyl for Yh, the compound (Ij) having amino(lower)alkylindolyl, lower alkylamino(lower)-alkylaminobenzimidazolyl or amino(lower)alkyl(N-lower)-alkylaminobenzimidazolyl for Yi may be obtained according to reaction condition. This case is included within the scope of this reaction.

Process 8

The object compound (I_l) or its salt can be prepared by subjecting a compound (I_k) or its salt to elimination reaction of the N-protective group.

Suitable salts of the compounds (I_k) and (I_l) may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as Process 7, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 7.

Process 9

The object compound (I_n) or its salt can be prepared by subjecting a compound (I_m) or its salt to deesterification reaction.

Suitable salts of the compounds (I_m) and (I_n) may be the same as those exemplified for the compound (I).

The reaction is carried out in accordance with a conventional method such as hydrolysis, reduction or the like.

The hydrolysis is preferably carried out in the presence of a base or an acid including Lewis acid. Suitable base may

include an inorganic base and an organic base such as an alkali metal [e.g. lithium, sodium, potassium, etc.], an alkaline earth metal [e.g. magnesium, calcium, etc.], the hydroxide or carbonate or bicarbonate thereof, trialkylamine
5 [e.g. trimethylamine, triethylamine, etc.], picoline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]-octane, 1,8-diazabicyclo[5.4.0]undec-7-ene, or the like. Suitable acid may include an organic acid [e.g. formic acid, acetic acid, propionic acid, trichloroacetic acid,
10 trifluoroacetic acid, etc.], an inorganic acid [e.g. hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, etc.] and Lewis acid [e.g. boron tribromide, etc.].

The reaction is usually carried out in a solvent such as
15 water, an alcohol [e.g. methanol, ethanol, etc.], xylene, diethylene glycol monomethyl ether, methylene chloride, tetrahydrofuran, a mixture thereof or any other solvent which does not adversely influence the reaction. A liquid base or acid can be also used as the solvent.

20 The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

The reduction can be applied preferably for elimination of the ester moiety such as 4-nitrobenzyl, 2-iodoethyl, 2,2,2-trichloroethyl, or the like. The reduction method
25 applicable for the elimination reaction may include chemical reduction and catalytic reduction.

Suitable reducing agents to be used in chemical reduction are a combination of metal [e.g. tin, zinc, iron, palladium back, etc.] or metallic compound [e.g. chromium
30 chloride, chromium acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, carbamic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.].

Suitable catalysts to be used in catalytic reduction are
35 conventional ones such as platinum catalysts [e.g. platinum

plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalyst [e.g. spongy palladium, palladium black, palladium oxide, palladium hydroxide, palladium on carbon, colloidal
5 palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalyst [e.g. reduced nickel, nickel oxide, Raney nickel, etc.], cobalt catalyst [e.g. reduced cobalt, Raney cobalt, etc.], iron catalyst [e.g. reduced iron, Raney iron, etc.], copper catalyst [e.g.
10 reduced copper, Raney copper, Ullman copper, etc.] or the like.

The reduction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, an alcohol [e.g. methanol, ethanol, propanol,
15 etc.], N,N-dimethylformamide, or a mixture thereof. Additionally, in case that the above-mentioned acids to be used in chemical reduction are in liquid, they can also be used as a solvent. Further, a suitable solvent to be used in catalytic reduction may be the above-mentioned solvent, and
20 other conventional solvent such as diethyl ether, dioxane, tetrahydrofuran, etc., or a mixture thereof.

The reaction temperature of this reduction is not critical and the reaction is usually carried out under cooling to warming.
25

Process 10

The object compound (Io) or its salt can be prepared by reacting a compound (In) or its reactive derivative at the carboxy group or a salt thereof with an amine or its salt.

30 Suitable salt of amine may be an acid addition salt as exemplified for the compound (I).

Suitable salts of the compounds (In) and (Io) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

35 Suitable "amine" may be ammonia, substituted or

unsubstituted lower alkylamine, substituted or unsubstituted N-containing heterocyclic compound and the like.

The substituted or unsubstituted lower alkylamine may be mono or di(lower)alkylamine (e.g. methylamine, ethylamine, propylamine, isopropylamine, butylamine, isobutylamine, pentylamine, hexylamine, dimethylamine, diethylamine, dipropylamine, dibutylamine, di-isopropylamine, dipentylamine, dihexylamine, etc.), pyridyl(lower)alkylamine, (e.g. pyridylmethylamine, etc.), lower alkylamino(lower)alkylamine (e.g. N-dimethylaminoethylamine, N-dimethylaminopropylamine, N-diethylaminoethyl-N-methylamine, etc.), morpholino(lower)alkylamine (e.g. morpholinoethylamine, etc.) or the like.

The substituted or unsubstituted N-containing heterocyclic compound may be a heterocyclic group substituted with amino (e.g. aminopyridine, N-methyl-N'-aminopiperazine, etc.), saturated 5 or 6-membered N-, or N- and S-, or N- and O-containing heterocyclic compound such as pyrrolidine, imidazolidine, piperidine, piperidone, piperazine, lower alkylaminopiperidine (e.g. dimethylaminopiperidine, etc.), N-(lower)alkylhomopiperazine (e.g. N-methylhomopiperazine, etc.), N-(lower)alkylpiperazine (e.g. N-methylpiperazine, N-ethylpiperazine, etc.), morpholine, thiomorpholine, N-pyridylpiperazine, N-hydroxy(lower)alkoxy(lower)-alkylpiperazine (e.g. N-hydroxyethoxyethylpiperazine, etc.), N-pyrrolidinylcarbonyl(lower)alkylpiperazine (e.g. N-pyrrolidinylcarbonylmethylpiperazine, etc.), or the like, in which preferable one is N-methylpiperazine.

This reaction can be carried out in substantially the same manner as Process 1, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 1.

35 Process 11

The object compound (Iq) or its salt can be prepared by subjecting a compound (Ip) or its salt to elimination reaction of methyl or the hydroxy-protective group.

5 Suitable salts of the compounds (Ip) and (Iq) may be the same as those exemplified for the compound (I).

 This reaction can be carried out in substantially the same manner as Process 7, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in
10 Process 7.

 In case that the hydroxy-protective group is tert-butyldiphenylsilyl, the reaction is preferably carried out in the presence of tetrabutylammonium fluoride.

15 Process 12

 The object compound (Is) or its salt can be prepared by reacting a compound (Ir) or its salt with an acylating agent.

 Suitable salts of the compounds (Ir) and (Is) may be the same as those exemplified for the compound (I).

20 This reaction can be carried out in substantially the same manner as Process 6, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in
 Process 6.

25

Process 13

 The object compound (It) or its salt can be prepared by reacting a compound (Ib) or its salt with N-lower alkylmethylenammonium halide.

30 Suitable salts of the compounds (Ib) and (It) may be the same as those exemplified for the compound (I).

 Suitable N-lower alkylmethylenammonium halide may be N-mono or di(lower alkyl)methylenammonium halide such as N-methylmethylenammonium chloride, N,N-dimethylmethylenammonium chloride or the like, in which preferable one is
35

N,N-dimethylmethyleneammonium chloride.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as chloroform, methylene chloride or the like.

5 The reaction temperature of this reaction is not critical and the reaction is usually carried out under cooling to heating.

Process 14

10 The object compound (Iv) or its salt can be prepared by subjecting a compound (Iu) or its salt to oxidation reaction.

Suitable salts of the compounds (Iu) and (Iv) may be the same as those exemplified for the compound (I).

15 Suitable oxidizing agent used in this reaction may be manganese dioxide, dimethyl sulfoxide, a mixture of dimethyl sulfoxide and oxalyl chloride or dimethyl sulfoxide and sulfur trioxide pyridine complex, and the like.

20 The reaction is usually carried out in a conventional solvent such as pentane, hexane, benzene, diethyl ether, dimethoxyethane, acetone, chloroform, dichloromethane or any other solvent which does not adversely influence the reaction.

Additionally in case that the above-mentioned oxidizing agent is liquid, it can be used as a solvent.

25 In this reaction, in case that dimethyl sulfoxide or a mixture of dimethyl sulfoxide and oxalyl chloride or dimethyl sulfoxide and sulfur trioxide pyridine complex is used as an oxidizing agent, the reaction is preferably carried out in the presence of alkali metal iodide (e.g. sodium iodide, etc.) and alkali metal carbonate (e.g. sodium carbonate) or
30 tri(lower)alkylamine (e.g. triethylamine, etc.).

The reaction temperature is not critical, and the reaction is usually carried out under cooling to heating.

35 Process 15

The object compound (Ix) or its salt can be prepared by subjecting a compound (Iw) or its salt to deesterification reaction.

5 Suitable salts of the compounds (Iw) and (Ix) may be the same as those exemplified for the compound (I).

 This reaction can be carried out in substantially the same manner as Process 9, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in
10 Process 9.

Process 16

 The object compound (Iz) or its salt can be prepared by subjecting a compound (Iy) or its salt to elimination
15 reaction of methyl substituted with aryl or substituted aryl.

 Suitable salts of the compounds (Iy) and (Iz) may be the same as those exemplified for the compound (I).

 This reaction can be carried out in substantially the same manner as Process 11, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as
20 explained in Process 11.

Process 17

25 The object compound (I-1) or its salt can be prepared by reacting a compound (Ix) or its reactive derivative at the carboxy group or a salt thereof with an amine or its salt.

 Suitable salts of the compounds (Ix) and (I-1) may be the same as those exemplified for the compound (I).

30 Suitable salts of amine may be an acid addition salt as exemplified for the compound (I).

 Suitable "amine" may be N-protected piperazine, oxopiperidine, lower alkylamine (e.g. dimethylamine, etc.), ammonia, lower alkylaminoamine (e.g. N,N-dimethylhydrazine, etc.), lower alkylamino(lower)alkyl(N-lower alkyl)amine (e.g.
35

dimethylaminoethyl(N-methyl)amine and the like.

This reaction can be carried out in substantially the same manner as Process 1, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those explained in Process 1.

Process 18

The object compound (I-3) or its salt can be prepared by reacting a compound (I-2) or its salt with a reducing agent.

Suitable salts of the compounds (I-2) and (I-3) may be the same as those exemplified for the compound (I).

Suitable reducing agent may be alkali metal borohydride (e.g. sodium borohydride, etc.), and the like.

The reaction is carried out in a solvent such as an alcohol (e.g. methanol, ethanol, etc.), tetrahydrofuran, or the like.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process 19

The object compound (I-4) or its salt can be prepared by reacting a compound (Iz) or its salt with an acylating agent.

Suitable salts of the compounds (Iz) and (I-4) may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as Process 6, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 6.

Process 20

The object compound (I-5) or its salt can be prepared by reacting a compound (Iz) or its salt with a compound (V).

Suitable salts of the compounds (Iz) and (I-5) may be

the same as those exemplified for the compound (I).

When the compound (V) having halogen for Z^2 is used in this reaction, the reaction is preferably carried out in the presence of a base such as an alkali metal (e.g. sodium, potassium, etc.), an alkaline earth metal (e.g. magnesium, calcium, etc.), the hydride or hydroxide or carbonate or bicarbonate thereof.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, dioxane, alcohol (e.g. methanol, ethanol, etc.), acetonitrile, tetrahydrofuran, acetic acid, N,N-dimethylformamide, or a mixture thereof.

The reaction temperature is not critical and the reaction can be carried out under cooling to heating.

Process 21

The object compound (I-6) or its salt can be prepared by subjecting a compound (I-5) or its salt to elimination reaction of the N-protective group.

Suitable salts of the compounds (I-5) and (I-6) may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as Process 8, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 8.

Process 22

The object compound (I-7) or its salt can be prepared by reacting a compound (I-6) or its salt with an acylating agent.

Suitable salts of the compounds (I-6) and (I-7) may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as Process 6, and therefore the reaction mode and

reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 6.

5 Process 23

The object compound (I-8) or its salt can be prepared by reacting a compound (I-6) or its salt with lower alkanal in the presence of a reducing agent.

10 Suitable salts of the compounds (I-6) and (I-8) may be the same as those exemplified for the compound (I).

Suitable lower alkanal may be C₁-C₆ alkanal such as formaldehyde, ethanal, propanal or the like, in which preferable one is formaldehyde.

15 Suitable reducing agent may be diborane, borane-organic amine complex [e.g. borane-pyridine complex, etc.], alkali metal cyanoborohydride [e.g. sodium cyanoborohydride, lithium cyanoborohydride, etc.], sodium borohydride and the like.

The reaction is preferably carried out in the presence of molecular sieves.

20 The reaction is usually carried out in a conventional solvent such as water, an alcohol [e.g. methanol, ethanol, etc.], dioxane, tetrahydrofuran, a mixture thereof or any other organic solvent which does not adversely influence the reaction.

25 The reaction may also be carried out in an acidic condition [e.g. presence of acetic acid, sulfuric acid, etc.] and the reaction temperature is not critical, and the reaction is usually carried out under cooling to warming.

30 Process 24.

The object compound (I-10) or its salt can be prepared by subjecting a compound (I-9) or its salt to reduction.

Suitable salts of the compounds (I-9) and (I-10) may be the same as those exemplified for the compound (I).

35 The reduction may include chemical reduction and

catalytic reduction, which are carried out in a conventional manner.

Suitable reducing agents to be used in chemical reduction are a metal [e.g. tin, zinc, iron, nickel, etc.],
5 a combination of such metal and/or metallic compound [e.g. nickel chloride, chromium chloride, chromium acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.],
10 a combination of such metal and/or metallic compound and base [e.g. ammonia, ammonium chloride, sodium hydroxide, etc.], a metal hydride compound such as aluminum hydride compound [e.g. lithium aluminum hydride, sodium aluminum hydride, aluminum hydride, lithium trimethoxyaluminum hydride, lithium
15 tri-t-butoxyaluminum hydride, etc.], borohydride compound [e.g. sodium borohydride, lithium borohydride, sodium cyanoborohydride, tetramethylammonium borohydride, borane, diborane, etc.], a phosphorus compound [e.g. phosphorus trichloride, phosphorus tribromide, triphenylphosphine, triethylphosphine, etc.] and the like.
20

Suitable catalysts to be used in catalytic reduction are conventional ones such as platinum catalysts [e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalyst
25 [e.g. spongy palladium, palladium black, palladium oxide, palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalyst [e.g. reduced nickel, nickel oxide, Raney nickel, etc.], cobalt catalyst [e.g. reduced cobalt, Raney cobalt, etc.],
30 iron catalyst [e.g. reduced iron, Raney iron, etc.], copper catalyst [e.g. reduced copper, Raney copper, Ullman copper, etc.], or the like.

The reduction is usually carried out in a solvent. A suitable solvent to be used may be water, an alcohol [e.g.
35 methanol, ethanol, propanol, etc.], acetonitrile or any other

conventional organic solvent such as dimethyl ether, dioxane, tetrahydrofuran, etc. or a mixture thereof.

The reaction temperature is not critical, and the reaction is preferably carried out under cooling to heating.

5

Process 25

The object compound (I-10) or its salt can be prepared by reacting a compound (I-11) or its salt with an azide compound.

10 Suitable salts of the compounds (I-10) and (I-11) may be the same as those exemplified for the compound (I).

 Suitable azide compound may be sodium azide, diphenylphosphorylazide and the like.

15 The reaction is usually carried out in a conventional solvent such as water, tetrahydrofuran, dioxane or any other organic solvent which does not adversely influence the reaction, or a mixture thereof.

 The reaction is also preferably carried out in the presence of a conventional base such as triethylamine, pyridine or the like.

20

 The reaction temperature is not critical, and the reaction is preferably carried out under warming to heating.

Process 26

25 The object compound (I-13) or its salt can be prepared by reacting a compound (I-12) or its reactive derivative at the carboxy group or a salt thereof with a reducing agent.

 Suitable salts of the compounds (I-13), (I-12) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

30

 Suitable reactive derivative at the carboxy group of the compound (I-12) may include an activated imide (e.g. phthalimido, etc.), an activated amide, an activated ester and the like.

35

 Suitable reducing agent may be aluminum hydride compound

[e.g. lithium aluminum hydride, lithium tri-t-butoxyaluminum hydride, etc.], borohydride compound [e.g. lithium borohydride, etc.], aluminum alkoxide [e.g. aluminum isopropoxide, etc.] and the like.

5 The reaction is usually carried out in a conventional solvent, such as diethyl ether, dioxane, or any other organic solvent which does not adversely influence the reaction, or a mixture thereof.

10 The reaction temperature is not critical, and the reaction is usually carried out under cooling to heating.

Process 27

15 The object compound (I-15) or its salt can be prepared by reacting a compound (I-14) or its salt with an acylating agent.

Suitable salts of the compounds (I-14) and (I-15) may be the same as those exemplified for the compound (I).

20 This reaction can be carried out in substantially the same manner as Process 6, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 6.

Process 28

25 The object compound (I-16) or its salt can be prepared by reacting a compound (I-15) or its salt with an alkali metal salt of phthalimide.

Suitable salts of the compounds (I-15) and (I-16) may be the same as those exemplified for the compound (I).

30 Suitable alkali metal salt of phthalimide may be potassium phthalimide and the like.

35 The reaction is usually carried out in a conventional solvent such as dimethyl sulfoxide, tetrahydrofuran or any other solvent which does not adversely influence the reaction.

The reaction temperature is not critical, and the reaction can be carried out under cooling to heating.

Process 29

5 The object compound (I-18) or its salt can be prepared by reacting a compound (I-17) or its salt with an amine.

Suitable salts of the compounds (I-17) and (I-18) may be the same as those exemplified for the compound (I).

10 Suitable amine may be N-lower alkylpiperazine, morpholine, dimethylamine, di(lower)alkylaminopiperidine, di(lower)alkylhydrazine, amino(lower)alkyl(N-lower alkyl)-amine, di(lower)alkylamino(lower)alkylamine and the like.

15 The reaction is carried out in no solvent or a solvent such as tetrahydrofuran, dioxane or any other solvent which does not adversely influence the reaction.

The reaction temperature is not critical, and the reaction is preferably carried out under warming to heating.

Process 30

20 The object compound (I-20) or its salt can be prepared by subjecting a compound (I-19) or its salt to elimination reaction of N-protective group.

Suitable salts of the compounds (I-19) and (I-20) may be the same as those exemplified for the compound (I).

25 This reaction can be carried out in substantially the same manner as Process 8, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 8.

30

Process 31

The object compound (I-22) or its salt can be prepared by reacting a compound (I-21) with hydroxylamine or its salt.

35 Suitable salts of the compounds (I-21) and (I-22) may be the same as those exemplified for the compound (I).

Suitable salt of hydroxylamine may be an acid addition salt as exemplified for the compound (I).

The reaction is preferably carried out in the presence of a conventional base such as sodium acetate, sodium
5 hydrogen carbonate or the like.

The reaction is usually carried out in a solvent which does not influence the reaction such as water, an alcohol (e.g. methanol, ethanol, etc.), pyridine or a mixture thereof.

10 The reaction temperature is not critical and the reaction is preferably carried out warming to heating.

Process 32

The object compound (I-23) or its salt can be prepared
15 by reacting a compound (VI) or its reactive derivative at the carboxy group or a salt thereof with a compound (VII) or its salt.

Suitable salts of the compounds (I-23), (VII), (VI) and its reactive derivative at the carboxy group may be the same
20 as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as Process 1, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in
25 Process 1.

The compounds obtained by the above processes can be isolated and purified by a conventional method such as pulverization, recrystallization, column chromatography,
30 reprecipitation, or the like.

It is to be noted that the compound (I) and the other compounds may include one or more stereoisomer(s) such as optical isomer(s) or geometrical isomer(s) due to asymmetric carbon atom(s) and double bond(s), and all of such isomers
35 and mixture thereof are included within the scope of this

invention.

Additionally, it is to be noted that any hydrate of the compound (I) is also included within the scope of this invention.

5 The object compound (I) and pharmaceutically acceptable salts thereof possess activities as vasopressin antagonistic activity, vasodilating activity, hypotensive activity, activity for inhibiting saccharide release in liver, activity for inhibiting growth of mesangium cells, water diuretic
10 activity, platelet agglutination inhibitory activity, oxytocin antagonistic activity and the like, and are useful for the treatment and/or prevention of hypertension, heart failure, renal insufficiency, edema, ascites, vasopressin parasecretion syndrome, hepatocirrhosis, hyponatremia,
15 hypokalemia, diabetic, circulation disorder, cerebrovascular disease (e.g. cerebral edema, cerebral infarction, etc.), Meniere's syndrome (e.g. Meniere's disease, etc.), motion sickness and the like in human being and animals.

20 In order to illustrate the usefulness of the object compound (I), the pharmacological data of the compound (I) are shown in the following.

Test 1

25

Vasopressin 1 (V1) receptor binding

(i) Test Method :

Blood was obtained by venipuncture from normal subjects.
30 Platelet-rich plasma (PRP) was prepared by centrifugation of whole blood at 200 xg for 10 minutes. PRP was centrifuged at 45,000 xg for 30 minutes. The remaining pellet was resuspended in 10 volume of ice cold 100 mM Tris-HCl (pH 7.4) buffer (containing 5 mM MgCl₂, 0.1% bovine serum albumin and
35 1 mM EDTA), and centrifuged at 45,000 xg for 30 minutes

again. The final pellet was resuspended in 100 mM Tris-HCl buffer. The resulting membrane preparation was used immediately for the binding assay.

Competition assays were conducted at equilibrium (15 minutes at 30°C) by using 1.5 nM ³H-vasopressin (40-87 Ci/mmol; New England Nuclear) in 100 mM Tris-HCl (pH 7.4) buffer. Nonspecific binding was determined by using 1 μM vasopressin. After incubation, reaction was terminated by adding 5 ml of ice-cold 100 mM Tris-HCl (pH 7.4) buffer, and then filtered rapidly through Whatman glass filter (GF/C). The filter was washed twice with the same buffer. The glass filter was mixed with liquid scintillation cocktail, and radioactivity was counted in a liquid scintillation counter. Competition activity of the test compound was represented by IC₅₀ values.

(ii) Test Result :

Test Compound (Example No.)	IC ₅₀ (nM)
39-27)	1.5
39-35)	<1.0
115-71)	0.7
115-81)	4.5

Test 2

Vasopressin 2 (V2) receptor binding

(i) Test Method :

For binding assays, the receptor cDNA was permanently expressed in Chinese hamster ovary (CHO) cells. CHO cells were transfected with a vector directing expression of the cDNA for the human V2 receptor and the clonal cell lines

expressing human V2 receptor was established essentially as described previously (Nakajina, Y., et. al. J. Biol. Chem., 1992, 267, 2437).

5 DNA-transfected cells were harvested and homogenized in ice cold 250 mM sucrose buffer containing 25 mM Tris-HCl (pH 7.4), 10 mM MgCl₂, 1 mM EDTA and 5 µg/ml p-amidinophenylmethylsulfonyl fluoride (A-PMSF). The homogenate was centrifuged at 500 xg for 10 minutes. The supernatant was centrifuged at 100,000 xg for 1 hour. The
10 final pellet was suspended in 25 mM Tris-HCl (pH 7.4) buffer (containing 10 mM MgCl₂, 1 mM EDTA and 5 µg/ml A-PMSF), and stored in small aliquots at -80°C.

Competition assays were conducted at equilibrium (2 hours at 22°C) by using 0.5 nM ³H-vasopressin (40-87 Ci/mmol, New England Nuclear) in 100 mM Tris-HCl (pH 7.4) buffer
15 (containing 5 mM MgCl₂, 5 µg/ml A-PMSF, 4 µg/ml leupeptin, 40 µg/ml bacitracin, 20 µg/ml chymostatin and 0.1% bovine serum albumin). Nonspecific binding was determined by using 1 µM vasopressin. After incubation, reaction mixture was rapidly
20 filtered through Whatman glass filter (GF/C). The filter was washed twice with the same buffer. The radioactivity was counted in a liquid scintillation counter. Competition activity of the test compound was represented by IC₅₀ values.

25 (ii) Test Result :

Test Compound (Example No.)	IC ₅₀ (nM)
39-27)	460
39-35)	380

30 For therapeutic purpose, the compound (I) of the present invention can be used in a form of pharmaceutical preparation containing one of said compounds, as an active ingredient, in

admixture with a pharmaceutically acceptable carrier such as an organic or inorganic solid, semi-solid or liquid excipient suitable for oral, parenteral or external (topical) administration. The pharmaceutical preparations may be capsules, tablets, dragees, granules, suppositories, solution, lotion, suspension, emulsion, ointment, gel, or the like. If desired, there may be included in these preparations, auxiliary substances, stabilizing agents, wetting or emulsifying agents, buffers and other commonly used additives.

While the dosage of the compound (I) will vary depending upon the age and condition of the patient, an average single dose of about 0.1 mg, 1 mg, 10 mg, 50 mg, 100 mg, 250 mg, 500 mg and 1000 mg of the compound (I) may be effective for treating the above-mentioned diseases. In general, amounts between 0.1 mg/body and about 1,000 mg/body may be administered per day.

The following Preparations and Examples are given for the purpose of illustrating this invention.

- to be continued on the next page -

Preparation 1

To a suspension of sodium hydride (133 mg) in tetrahydrofuran (5.0 ml) was added dropwise a solution of benzyl indole-4-carboxylate (580 mg) in tetrahydrofuran (5.0 ml) at 0°C and the mixture was stirred at 0°C for 1 hour. 4-Toluenesulfonyl chloride (440 mg) was added to the mixture and the solution was stirred at ambient temperature for 1 hour. The reaction was quenched with 1N hydrochloric acid and then the aqueous solution was extracted with ethyl acetate. Drying, filtering and removal of solvents afforded a crude product. The crude product was purified by column chromatography (eluent; n-hexane:ethyl acetate = 15:1) to give benzyl 1-(4-toluenesulfonyl)indole-4-carboxylate (560 mg) as a colorless syrup.

NMR (CDCl₃, δ) : 2.32 (3H, s), 5.39 (2H, s), 7.19-7.23 (2H, m), 7.31-7.48 (7H, m), 7.67 (1H, d, J=4Hz), 7.72 (2H, d, J=9Hz), 8.00 (1H, d, J=8Hz), 8.20 (1H, d, J=8Hz)

Preparation 2

To a suspension of sodium hydride (174 mg) in tetrahydrofuran (8.0 ml) was added dropwise a solution of benzyl indole-7-carboxylate (700 mg) in tetrahydrofuran (7.0 ml) at 0°C and the mixture was stirred at 0°C for 1 hour. Chloromethyl pivalate (461 mg) was added to the mixture and the solution was stirred at ambient temperature for 3 hours. The reaction was quenched with 1N hydrochloric acid and then the aqueous solution was extracted with ethyl acetate. Drying, filtering and removal of solvents afforded benzyl 1-pivaloyloxymethylindole-7-carboxylate (1.08 g) as a yellow oil.

NMR (CDCl₃, δ) : 1.01 (9H, s), 5.42 (2H, s), 6.40 (2H, s), 6.58 (1H, d, J=4Hz), 7.17 (1H, t, J=8Hz), 7.28 (1H, d, J=4Hz), 7.33-7.42 (3H, m), 7.47-7.51 (2H, m), 7.73-7.80 (2H, m)

Preparation 3

To a solution of 2,2,6,6-tetramethylpiperidine (322 mg) in tetrahydrofuran (5.0 ml) was added dropwise a solution of n-butyllithium (1.6M n-hexane solution 1.3 ml) at -70 ~ -60°C and the solution was stirred at 0°C for 30 minutes. A solution of benzyl 1-tert-butoxycarbonylindole-4-carboxylate (500 mg) in tetrahydrofuran (2.5 ml) was added dropwise to the above solution at -70 ~ -60°C and the mixture was stirred at -70°C for 30 minutes. To the mixture was added a solution of ethyl chloroformate (185 mg) in tetrahydrofuran (2.5 ml) at such a rate as to maintain the temperature below -60°C. The solution was stirred at -70°C for 2 hours and the reaction was quenched with aqueous saturated ammonium chloride solution at -20°C. The aqueous solution was extracted with ethyl acetate. Drying, filtering and removal of solvents afforded a crude product. The crude product was chromatographed on silica gel (eluent; n-hexane:ethyl acetate = 15:1) to give benzyl 1-tert-butoxycarbonyl-2-ethoxycarbonylindole-4-carboxylate (100 mg) as a colorless oil.

NMR (CDCl₃, δ) : 1.39 (3H, t, J=7Hz), 1.62 (9H, s), 4.38 (2H, q, J=7Hz), 5.43 (2H, s), 7.31-7.50 (6H, m), 7.78 (1H, s), 8.04 (1H, d, J=8Hz), 8.32 (1H, d, J=9Hz)

Preparation 4

To a solution of 2-amino-3-nitrobenzoic acid (4.47 g) in 1,2-dichloroethane (50 ml) was added trifluoroacetic anhydride (10.3 g) at 5°C and the mixture was stirred at ambient temperature for 5 hours. To the mixture was added trifluoroacetic anhydride (5.15 g) and it was stirred at ambient temperature for additional 1 hour. The solution was concentrated in vacuo to give 8-nitro-2-trifluoromethyl-3,1-benzoxazin-4-one as a slight yellow powder (6.35 g).

NMR (CDCl₃, δ) : 7.86 (1H, t, J=7Hz), 8.29 (1H, d,

J=7Hz), 8.51 (1H, d, J=7Hz)

Preparation 5

To a solution of ethyl 2-(N-benzylamino)-3-nitrobenzoate
5 (400 mg) in N,N-dimethylaniline (3 ml) was added
methoxyacetyl chloride (318 mg) at ambient temperature and
the mixture was stirred at 90°C for 4 hours. The reaction
mixture was poured into water and the aqueous solution was
10 extracted with ethyl acetate. The organic layer was washed
successively with 1N hydrochloric acid, water and brine and
the solution was dried over magnesium sulfate. The solvent
was evaporated in vacuo and the residue was chromatographed
on silica gel eluting with a mixture of n-hexane and ethyl
acetate (3:1) to give ethyl 2-(N-benzyl-N-methoxyacetyl)-
15 amino-3-nitrobenzoate (480 mg) as an oil.

NMR (CDCl₃, δ) : 1.29 (3H, t, J=7Hz), 3.37 (3H, s),
3.92 (2H, s), 4.09 (2H, q, J=7Hz), 4.74 (1H, d,
J=13Hz), 4.83 (1H, d, J=13Hz), 7.00-7.11 (2H, m),
7.11-7.31 (3H, m), 7.59 (1H, t, J=8Hz), 7.96 (1H,
20 d, J=8Hz), 8.08 (1H, d, J=8Hz)

Preparation 6

A mixture of 2,3-diaminotoluene (2.0 g) and ethyl
N-methyloxamate (2.36 g) in N,N-dimethylformamide (10 ml) was
25 stirred at 175°C for 8 hours. After being cooled to ambient
temperature, the mixture was poured into a mixture of
saturated aqueous sodium bicarbonate solution and ethyl
acetate and the organic layer was separated. The organic
layer was washed with brine and the solution was dried over
30 magnesium sulfate. The solvent was evaporated in vacuo and
the residue was chromatographed on silica gel eluting with a
mixture of chloroform and methanol (50:1) to give 4-methyl-2-
(N-methylcarbamoyl)-1H-benzimidazole (1.17 g) as a powder.

NMR (CDCl₃, δ) : 2.60 (3H x 1/2, s), 2.66 (3H x 1/2,
35 s), 3.10 (3H x 1/2, s), 3.12 (3H x 1/2, s), 7.06-

7.19 (1H, m), 7.19-7.29 (1H, m), 7.36 (1H x 1/2, d, J=8Hz), 7.61 (1H x 1/2, d, J=8Hz), 7.71 (1H, br peak)

5 Preparation 7

To a suspension of 4-methyl-2-(N-methylcarbamoyl)-1H-benzimidazole (1.0 g) in 1N-aqueous sodium hydroxide solution (15 ml) was added portionwise potassium permanganate (3.34 g) at 100°C and the reaction mixture was stirred at the same
10 temperature for 15 minutes. The reaction mixture was filtered through a bed of celite and the filtrate was washed with chloroform. The aqueous layer was adjusted to pH 3 with 4N hydrochloric acid. The precipitate was collected by vacuum filtration to give 2-(N-methylcarbamoyl)-1H-
15 benzimidazole-4-carboxylic acid (647 mg) as a solids.

NMR (DMSO-d₆, δ) : 2.85 (3H, d, J=5Hz), 7.41 (1H, t, J=8Hz), 7.90 (1H, d, J=8Hz), 7.95 (1H, d, J=8Hz), 9.06 (1H, q-like)

20 Preparation 8

To a solution of methyl 2-hydroxymethyl-1H-benzimidazole-4-carboxylate (1.0 g) in N,N-dimethylformamide (10 ml) were added tert-butylchlorodiphenylsilane (1.87 g) and imidazole (495 mg) at ambient temperature and the mixture
25 was stirred at the same temperature for 28 hours. The reaction mixture was poured into water and the aqueous solution was extracted with ethyl acetate. The organic layer was washed with water and brine and dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue
30 was chromatographed on silica gel eluting with a mixture of n-hexane and ethyl acetate (10:1) to give methyl 2-tert-butyl-diphenylsiloxymethyl-1H-benzimidazole-4-carboxylate (1.45 g) as an oil.

NMR (CDCl₃, δ) : 1.18 (9H, s), 4.02 (3H, s), 5.06 (2H, s), 7.30 (1H, t, J=8Hz), 7.35-7.50 (6H, m), 7.67-

35

7.73 (4H, m), 7.89 (1H, d, J=8Hz), 7.92 (1H, d, J=8Hz)

Preparation 9

5 To a solution of methyl 2-tert-butyldiphenylsiloxymethyl-1H-benzimidazole-4-carboxylate (500 mg) in pyridine (3 ml) was added lithium iodide (602 mg) under nitrogen at ambient temperature and the mixture was heated to reflux for 3 hours. The reaction mixture was concentrated in vacuo and
10 the residue was dissolved in chloroform. The solution was washed with water and brine and the organic layer was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was chromatographed on silica gel eluting with a mixture of chloroform and methanol (chloroform only-
15 50:1-25:1-10:1) to give 2-tert-butyldiphenylsiloxymethyl-1H-benzimidazole-4-carboxylic acid (425 mg) as a powder.

NMR (DMSO-d₆, δ) : 1.03 (9H, s), 5.00 (2H, s), 7.28 (1H, t, J=8Hz), 7.38-7.52 (6H, m), 7.70-7.75 (4H, m), 7.80 (1H, d, J=8Hz), 7.91 (1H, d, J=8Hz)

20

Preparation 10

The mixture of benzyl 1-(4-toluenesulfonyl)indole-4-carboxylate (550 mg) and 10% palladium on charcoal (200 mg) in methanol (20 ml) and water (2 ml) was hydrogenated at
25 ambient temperature (an initial hydrogen pressure was set to 3.5 atm.). The theoretical amount of hydrogen was absorbed in 6 hours. The resulting mixture was filtered through a bed of celite and the filtrate was evaporated in vacuo. The residue was diluted with chloroform and the solution was
30 dried over magnesium sulfate. Filtering and removal of solvents afforded a crude product. The crude product was triturated with diethyl ether-n-hexane (1:3) to give 1-(4-toluenesulfonyl)indole-4-carboxylic acid (330 mg) as a brown powder.

35 NMR (DMSO-d₆, δ) : 2.32 (3H, s), 7.30 (1H, d, J=4Hz),

7.37-7.43 (2H, m), 7.47 (1H, d, J=8Hz), 7.89 (3H, d, J=8Hz), 7.97 (1H, d, J=4Hz), 8.20 (1H, d, J=8Hz)

Preparation 11

5 The following compounds were obtained according to a similar manner to that of Preparation 10.

1) 1-Pivaloyloxymethylindole-7-carboxylic acid

10 NMR (DMSO-d₆, δ) : 1.00 (9H, s), 6.40 (2H, s), 6.61 (1H, d, J=3Hz), 7.16 (1H, t, J=8Hz), 7.53 (1H, d, J=3Hz), 7.60 (1H, d, J=8Hz), 7.78 (1H, d, J=8Hz)

2) 1-Methylindole-7-carboxylic acid

15 NMR (CDCl₃, δ) : 3.88 (3H, s), 6.52 (1H, d, J=4Hz), 7.00-7.12 (2H, m), 7.72-7.79 (2H, m)

Preparation 12

20 To a mixture of 10% palladium on charcoal (130 mg) in 5.0% formic acid-methanol (5.0 ml) was added a solution of benzyl 1-tert-butoxycarbonyl-2-ethoxycarbonylindole-4-carboxylate (130 mg) in 5.0% formic acid-methanol (5.0 ml). The mixture was stirred under nitrogen atmosphere at ambient temperature for 30 minutes. The resulting mixture was filtered through a bed of celite and the filtrate was
25 evaporated in vacuo to give 1-tert-butoxycarbonyl-2-ethoxycarbonylindole-4-carboxylic acid (87 mg) as a white crystal.

30 NMR (CDCl₃, δ) : 1.44 (3H, t, J=7Hz), 1.66 (9H, s), 4.42 (2H, q, J=7Hz), 7.50 (1H, t, J=9Hz), 7.83 (1H, s), 8.12 (1H, d, J=9Hz), 8.39 (1H, d, J=9Hz)

Preparation 13

35 To an ice water bath cooled 4N hydrogen chloride solution in 1,4-dioxane (5 ml) was added 2-(N-tert-butoxycarbonyl-N-methyl)amino-3-nitrobenzoic acid (900 mg)

and the solution was stirred at ambient temperature for 2 hours. The reaction mixture was concentrated in vacuo and the residue was washed with diethyl ether and collected by vacuum filtration to give 2-(N-methylamino)-3-nitrobenzoic acid hydrochloride (687 mg) as a powder.

NMR (DMSO-d₆, δ) : 2.70 (3H, s), 6.73 (1H, t, J=8Hz), 7.98 (1H, d, J=8Hz), 8.04 (1H, d, J=8Hz)

Preparation 14

The following compound was obtained according to a similar manner to that of Preparation 13.

Ethyl 2-(N-benzylamino)-3-nitrobenzoate

NMR (CDCl₃, δ) : 1.38 (3H, t, J=7Hz), 4.16 (2H, d, J=6Hz), 4.35 (2H, q, J=7Hz), 6.72 (1H, t, J=8Hz), 7.22-7.46 (5H, m), 8.00 (1H, d, J=8Hz), 8.11 (1H, d, J=8Hz), 8.80 (1H, br s)

Preparation 15

To a suspension of sodium hydride (60% dispersion in mineral oil, 142 mg) in N,N-dimethylformamide (1 ml) was added dropwise a solution of ethyl 2-(N-tert-butoxycarbonyl)amino-3-nitrobenzoate (1.0 g) in N,N-dimethylformamide (5 ml) under nitrogen in ice water bath and stirred at the same temperature for 1 hour. To the mixture was added methyl iodide (526 mg) at 0°C under nitrogen and the solution was stirred at the same temperature for 2 hours. The reaction mixture was poured into water and the aqueous solution was extracted with ethyl acetate. The organic layer was washed with water and brine, dried over magnesium sulfate and the solvent was evaporated to give ethyl 2-(N-tert-butoxycarbonyl-N-methyl)amino-3-nitrobenzoate (1.05 g) as an oil.

NMR (CDCl₃, δ) : 1.28 (9H, s), 1.41 (3H, t, J=7.5Hz), 3.10-3.20 (3H, m), 4.30-4.48 (2H, m), 7.52 (1H, t,

J=8Hz), 7.95 (1H, d, J=8Hz), 8.10 (1H, d, J=8Hz)

Preparation 16

5 The following compounds were obtained according to a similar manner to that of Preparation 15.

- 1) Ethyl 2-(N-benzyl-N-tert-butoxycarbonyl)amino-3-nitrobenzoate

10 NMR (CDCl₃, δ) : 1.17-1.38 (12H, m), 4.08-4.23 (2H, m), 4.53 (1H, d, J=13Hz), 4.81 (1H, d, J=13Hz), 7.03-7.16 (2H, m), 7.16-7.29 (3H, m), 7.45 (1H, t, J=8Hz), 7.88 (1H, d, J=8Hz), 8.04 (1H, d, J=8Hz)

- 2) 3-(N-Acetyl-N-methyl)amino-2-nitrobenzoic acid

15 NMR (CDCl₃, δ) : 1.87 (3H, s), 3.20 (3H, s), 7.55 (1H, d, J=8Hz), 7.69 (1H, t, J=8Hz), 8.20 (1H, d, J=8Hz)

- 3) 3-(N-Acetyl-N-ethyl)amino-2-nitrobenzoic acid

20 NMR (DMSO-d₆, δ) : 1.00 (3H, t, J=7Hz), 1.72 (3H, s), 3.08 (1H, m), 3.86 (1H, m), 7.78-7.91 (2H, m), 8.12 (1H, d, J=8Hz)

Preparation 17

25 To a solution of ethyl 2-(N-tert-butoxycarbonyl-N-methyl)amino-3-nitrobenzoate (1.0 g) in ethanol (10 ml) was added 1N aqueous sodium hydroxide solution (3.5 ml) and the solution was stirred at ambient temperature for 1 day. The reaction mixture was concentrated and the residue was dissolved in water. The aqueous layer was washed with
30 diethyl ether and the aqueous solution was adjusted to pH 4 with 1N hydrochloric acid. The solution was extracted with chloroform and the organic layer was separated. The solution was washed with water and brine and the solution was dried over magnesium sulfate. The solvent was evaporated in vacuo
35 to give 2-(N-tert-butoxycarbonyl-N-methyl)amino-3-

nitrobenzoic acid (910 mg) as a powder.

NMR (CDCl₃, δ) : 1.26 (9H, s), 3.20 (3H, s), 7.55 (1H, t, J=8Hz), 8.00 (1H, d, J=8Hz), 8.22 (1H, d, J=8Hz)

5 Preparation 18

The following compounds were obtained according to a similar manner to that of Preparation 17.

10 1) 3-Benzyl-2-methoxymethyl-3H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 3.40 (3H, s), 4.75 (2H, s), 5.95 (2H, s), 6.83-6.91 (2H, m), 7.16-7.26 (3H, m), 7.33 (1H, t, J=8Hz), 7.72 (1H, d, J=8Hz), 7.89 (1H, d, J=8Hz)

15 2) 1,2-Dimethyl-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 2.64 (3H, s), 3.82 (3H, s), 7.37 (1H, t, J=8Hz), 7.76 (1H, d, J=8Hz), 7.86 (1H, d, J=8Hz)

20 3) 1-Ethyl-2-methyl-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 1.33 (3H, t, J=7Hz), 2.68 (3H, s), 4.33 (2H, q, J=7Hz), 7.37 (1H, t, J=8Hz), 7.78 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz)

25 4) 2-Methyl-1-propyl-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 0.94 (3H, t, J=7Hz), 1.82 (2H, m), 2.88 (3H, s), 4.41 (2H, t, J=7Hz), 7.65 (1H, t, J=8Hz), 8.05 (1H, d, J=8Hz), 8.24 (1H, d, J=8Hz)

30 5) 1-(4-Methoxybenzyl)-2-(N-methylcarbamoyl)-1H-benzimidazole-4-carboxylic acid

NMR (CD₃OD, δ) : 2.96 (3H, s), 3.72 (3H, s), 6.00 (2H, s), 6.83 (2H, d, J=8Hz), 7.20 (2H, d, J=8Hz), 7.45 (1H, t, J=8Hz), 7.84 (1H, d, J=8Hz), 8.03 (1H, d,

J=8Hz)

Preparation 19

To a solution of ethyl 2-(N-benzyl-N-methoxyacetyl)amino-3-nitrobenzoate (478 mg) in ethanol (5 ml) were added iron powder (358 mg) and acetic acid (771 mg) and the mixture was refluxed for 2 hours. The reaction mixture was filtered through a bed of celite and the filtrate was concentrated in vacuo. The residue was diluted with a mixture of ethyl acetate and saturated aqueous sodium bicarbonate solution and the mixture was filtered through a bed of celite again. The organic layer was separated and washed with water and brine. The solution was dried over magnesium sulfate and the solvent was evaporated in vacuo to give ethyl 3-benzyl-2-methoxymethyl-3H-benzimidazole-4-carboxylate (364 mg) as an oil.

NMR (CDCl₃, δ) : 1.20 (3H, t, J=7Hz), 3.40 (3H, s), 4.15 (2H, q, J=7Hz), 4.75 (2H, s), 5.91 (2H, s), 6.78-6.89 (2H, m), 7.14-7.41 (4H, m), 7.68 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz)

Preparation 20

The following compound was obtained according to a similar manner to that of Preparation 19.

Methyl 2-methyl-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 2.67 (3H, s), 4.00 (3H, s), 7.25 (1H, t, J=8Hz), 7.85 (1H, d, J=8Hz), 7.89 (1H, d, J=8Hz)

Preparation 21

The following compounds were obtained by using methyl 3-(N-acetyl-N-methyl)amino-2-nitrobenzoate as a starting compound according to a similar manner to that of Preparation 19.

A mixture of methyl 1,2-dimethyl-1H-benzimidazole-4-carboxylate and ethyl 1,2-dimethyl-1H-benzimidazole-4-carboxylate

5 Methyl 1,2-dimethyl-1H-benzimidazole-4-carboxylate
NMR (CDCl₃, δ) : 2.69 (3H, s), 3.75 (3H, s), 4.02 (3H, s), 7.28 (1H, t, J=8Hz), 7.48 (1H, d, J=8Hz), 7.95 (1H, d, J=8Hz)

10 Ethyl 1,2-dimethyl-1H-benzimidazole-4-carboxylate
NMR (CDCl₃, δ) : 1.44 (3H, t, J=7Hz), 2.67 (3H, s), 3.72 (3H, s), 4.48 (2H, q, J=7Hz), 7.26 (1H, t, J=8Hz), 7.45 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz)

15 Preparation 22

The following compound was obtained by using methyl 2-(N-acetyl-N-ethyl)amino-3-nitrobenzoate as a starting compound according to a similar manner to that of Preparation 19.

20 Ethyl 1-ethyl-2-methyl-1H-benzimidazole-4-carboxylate
NMR (CDCl₃, δ) : 1.37 (3H, t, J=7Hz), 1.43 (3H, t, J=7Hz), 2.66 (3H, s), 4.19 (2H, q, J=7Hz), 4.48 (2H, q, J=7Hz), 7.25 (1H, t, J=8Hz), 7.47 (1H, d, J=8Hz), 7.89 (1H, d, J=8Hz)

25

Preparation 23

To a solution of 3-(N-acetyl-N-methyl)amino-2-nitrobenzoic acid (700 mg) in 20% methanol in benzene solution (5 ml) was added dropwise 2N trimethylsilyldiazomethane in n-hexane solution (5 ml) in ice water bath and the mixture was allowed to stand at ambient temperature for 1 hour. The reaction mixture was concentrated in vacuo and the residue was dissolved in ethyl acetate. The solution was washed successively with saturated

30

35

aqueous sodium bicarbonate solution, water and brine and the organic layer was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was chromatographed on silica gel eluting with a mixture of n-hexane - ethyl acetate (10:1) to give methyl 3-(N-acetyl-N-methyl)amino-2-nitrobenzoate (547 mg) as an oil.

NMR (CDCl₃, δ) : 1.85 (3H, s), 3.19 (3H, s), 3.94 (3H, s), 7.54 (1H, d, J=8Hz), 7.68 (1H, t, J=8Hz), 8.13 (1H, d, J=8Hz)

Preparation 24

The following compounds were obtained according to a similar manner to that of Preparation 23.

1) Methyl 3-(N-acetyl-N-ethyl)amino-2-nitrobenzoate

NMR (CDCl₃, δ) : 1.11 (3H, t, J=7Hz), 1.83 (3H, s), 3.20 (1H, m), 3.93 (3H, s), 4.10 (1H, m), 7.49 (1H, d, J=8Hz), 7.67 (1H, t, J=8Hz), 8.13 (1H, d, J=8Hz)

2) Methyl 2-(N-methylcarbamoyl)-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 3.08 (3H, d, J=5Hz), 4.03 (3H, s), 7.40 (1H, t, J=8Hz), 7.47 (1H, br s), 7.98 (1H, d, J=8Hz), 8.04 (1H, d, J=8Hz)

Preparation 25

2-(N-Methylamino)-3-nitrobenzoic acid hydrochloride (250 mg) in methanol (5 ml) was hydrogenated under medium pressure (3 atm.) at ambient temperature for 3 hours. The reaction mixture was filtered through a bed of celite and the filtrate was concentrated in vacuo. The crude 3-amino-2-(N-methylamino)benzoic acid hydrochloride was used without further purification.

NMR (CD₃OD, δ) : 3.06 (3H, s), 7.23 (1H, d, J=8Hz), 7.32 (1H, t, J=8Hz), 7.59 (1H, d, J=8Hz)

Preparation 26

The following compound was obtained according to a similar manner to that of Example 13.

- 5 3-Methyl-3H-benzimidazole-4-carboxylic acid hydrochloride
NMR (DMSO-d₆, δ) : 4.07 (3H, s), 7.50 (1H, t, J=8Hz),
7.88 (1H, d, J=8Hz), 8.00 (1H, d, J=8Hz), 9.02 (1H,
s)

10 Preparation 27

The following compound was obtained by using 2-amino-3-hydroxybenzoic acid as a starting compound according to a similar manner to that of Example 13.

- 15 4-Benzoxazolecarboxylic acid
NMR (DMSO-d₆, δ) : 7.55 (1H, t, J=8Hz), 7.92 (1H, d,
J=8Hz), 8.03 (1H, d, J=8Hz), 8.85 (1H, s)

Preparation 28

- 20 The following compound was obtained by using 3-amino-2-hydroxybenzoic acid as a starting compound according to a similar manner to that of Example 13.

- 7-Benzoxazolecarboxylic acid
25 NMR (DMSO-d₆, δ) : 7.51 (1H, t, J=8Hz), 7.96 (1H, d,
J=8Hz), 8.07 (1H, d, J=8Hz), 8.88 (1H, s)

Preparation 29

- The following compound was obtained according to a
30 similar manner to that of Example 5.

- Benzyl 1-methylindole-7-carboxylate
NMR (CDCl₃, δ) : 3.83 (3H, s), 5.41 (2H, s), 6.53 (1H,
d, J=3Hz), 7.01-7.11 (2H, m), 7.31-7.42 (3H, m),
35 7.47-7.50 (2H, m), 7.70 (1H, d, J=8Hz), 7.77 (1H,

d, J=8Hz)

Preparation 30

To a solution of benzyl indole-4-carboxylate (1.85 g)
5 and N,N-dimethylaminopyridine (180 mg) in acetonitrile (10 ml) was added portionwise di-tert-butyl dicarbonate (1.61 g), and then the mixture was stirred at ambient temperature for 2 hours and stand overnight. The resulting mixture was concentrated in vacuo and the residue was diluted with ethyl acetate (30 ml). The organic layer was washed successively
10 with 1N hydrochloric acid, saturated sodium bicarbonate aqueous solution and brine. The solution was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by silica gel column chromatography (eluent;
15 n-hexane:ethyl acetate = 10:1) to give benzyl 1-tert-butoxycarbonylindole-4-carboxylate (2.26 g).

NMR (CDCl₃, δ) : 1.68 (9H, s), 5.42 (2H, s), 7.26 (1H, d, J=4Hz), 7.31-7.43 (4H, m), 7.47-7.51 (2H, m), 7.69 (1H, d, J=4Hz), 8.02 (1H, d, J=8Hz), 8.40 (1H, d, J=9Hz)
20

Preparation 31

To a solution of methyl 2-methyl-1H-benzimidazole-4-carboxylate (250 mg) in N,N-dimethylformamide (4 ml) were
25 added potassium carbonate (363 mg) and n-propyl bromide at ambient temperature and the mixture was stirred at the same temperature for 2 days. The reaction mixture was poured into water and the aqueous solution was extracted with ethyl acetate. The organic layer was washed with water and brine
30 and the solution was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was chromatographed on silica gel eluting with a mixture of n-hexane and ethyl acetate (1:1-1:2-1:3-ethyl acetate only) to give methyl 2-methyl-1-propyl-1H-benzimidazole-4-
35 carboxylate (173 mg) as an oil.

NMR (CDCl₃, δ) : 0.97 (3H, t, J=7Hz), 1.84 (2H, m),
2.69 (3H, s), 4.05 (3H, s), 4.11 (2H, t, J=7Hz),
7.27 (1H, t, J=8Hz), 7.49 (1H, d, J=8Hz), 7.94 (1H,
d, J=8Hz)

5

Preparation 32

The following compound was obtained according to a similar manner to that of Preparation 31.

10 Methyl 1-(4-methoxybenzyl)-2-(N-methylcarbamoyl)-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 3.04 (3H, d, J=5Hz), 3.74 (3H, s),
4.04 (3H, s), 6.01 (2H, s), 6.80 (2H, d, J=8Hz),
7.17 (2H, d, J=8Hz), 7.38 (1H, t, J=8Hz), 7.63 (1H,
15 d, J=8Hz), 8.01 (1H, d, J=8Hz), 8.05 (1H, br peak)

Example 1

To a mixture of 1H-imidazo[4,5-b]pyridine-7-carboxylic acid (203 mg) and oxalyl chloride (0.217 ml) in
20 dichloromethane (25 ml) was added 1 drop of
N,N-dimethylformamide and the mixture was stirred at ambient temperature for 2 hours. After being removed a solvent by evaporation, residual acid chloride in dichloromethane (5 ml) was added to a mixture of 4-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yloxy]-phenyl]benzamide (400 mg) and triethylamine (210 mg) in
25 dichloromethane (20 ml) and the mixture was stirred at ambient temperature for 2 hours. The mixture was washed successively with saturated aqueous sodium hydrogen carbonate and brine, and dried over sodium sulfate. The solvent was
30 removed by rotary evaporation and the residue was purified by silica gel column chromatography (SiO₂ 30 g, 3% methanol in dichloromethane) to give 4-[1H-imidazo[4,5-b]pyridin-7-yl]-carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]phenyl]-
35

benzamide (393 mg).

NMR (CDCl₃, δ) : 1.42-1.61 (2H, m), 1.63-1.92 (4H, m),
2.25 (3H, s), 2.29 (3H, s), 2.32-2.47 (6H, m), 3.34
(3H, s), 3.42-3.55 (2H, m), 3.60-3.70 (2H, m),
5 3.72-4.00 (5H, m), 6.50-6.66 (3H, m), 6.76-7.08
(3H, m), 8.03 (1H, m), 8.32 (1H, s), 8.44 (1H, m),
8.59 (1H, m)

Example 2

10 The following compounds were obtained according to a
similar manner to that of Example 1.

1) 4-[[1-(4-Toluenesulfonyl)indol-4-yl]carbonyl]amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
15 yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.48-1.58 (2H, m), 1.66-1.87 (4H, m),
2.27 (3H, s), 2.31-2.39 (8H, m), 2.42-2.53 (4H, m),
3.32 (3H, s), 3.52-3.58 (2H, m), 3.64-3.72 (2H, m),
3.77 (3H, s), 3.83-4.00 (2H, m), 6.59 (1H, d,
20 J=8Hz), 6.62 (1H, s), 6.85 (1H, d, J=8Hz), 6.92
(1H, d, J=8Hz), 7.01 (1H, s), 7.19-7.27 (3H, m),
7.38 (1H, t, J=8Hz), 7.58 (1H, d, J=8Hz), 7.69 (1H,
d, J=4Hz), 7.75 (2H, d, J=8Hz), 8.17 (1H, d,
J=8Hz), 8.28 (1H, d, J=8Hz), 8.47 (1H, s)

25

2) 4-[(1-Pivaloyloxymethylindol-7-yl)carbonyl]amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 0.84 (9H, s), 1.48-1.61 (2H, m),
30 1.68-1.90 (4H, m), 2.28 (3H, s), 2.31 (3H, s),
2.33-2.46 (6H, m), 3.33 (3H, s), 3.48-3.54 (2H, m),
3.60-3.68 (2H, m), 3.70 (3H, s), 3.88-4.00 (2H, m),
6.28 (2H, s), 6.57 (1H, d, J=3Hz), 6.59-6.66 (2H,
m), 6.83 (1H, d, J=8Hz), 6.97 (1H, d, J=8Hz), 7.02
35 (1H, s), 7.19 (1H, t, J=8Hz), 7.29 (1H, d, J=3Hz),

7.42 (1H, d, J=8Hz), 7.74 (1H, d, J=8Hz), 8.28-8.37 (2H, m)

3) 4-[(1-Methylindol-7-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.48-1.89 (6H, m), 2.29 (6H, s), 2.31-2.42 (6H, m), 3.33 (3H, s), 3.46-3.51 (2H, m), 3.59-3.67 (2H, m), 3.72 (3H, s), 3.80 (3H, s), 3.88-4.00 (2H, m), 6.55 (1H, d, J=4Hz), 6.61 (1H, d, J=8Hz), 6.67 (1H, s), 6.81-7.12 (5H, m), 7.33 (1H, d, J=8Hz), 7.72 (1H, d, J=8Hz), 8.28-8.36 (2H, m)

4) 4-(1-tert-Butoxycarbonyl-2-ethoxycarbonylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.40 (3H, t, J=8Hz), 1.50-1.89 (6H, m), 1.63 (9H, s), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.43 (6H, m), 3.34 (3H, s), 3.47-3.52 (2H, m), 3.60-3.68 (2H, m), 3.79 (3H, s), 3.87-4.00 (2H, m), 4.38 (2H, q, J=8Hz), 6.60 (1H, d, J=8Hz), 6.64 (1H, s), 6.86 (1H, d, J=8Hz), 6.95 (1H, d, J=8Hz), 7.06 (1H, s), 7.48 (1H, t, J=8Hz), 7.62 (1H, d, J=8Hz), 7.69 (1H, s), 8.27-8.33 (2H, m), 8.54 (1H, s)

5) 4-[2-Chloro-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.45-1.60 (2H, m), 1.63-1.92 (4H, m), 2.25 (3H, s), 2.29 (3H, s), 2.31-2.49 (6H, m), 3.35 (3H, s), 3.44-3.55 (2H, m), 3.59-3.70 (2H, m), 3.71-4.01 (5H, m), 6.52-6.66 (2H, m), 6.80-7.06 (3H, m), 7.24-7.37 (1H, m), 7.42-8.50 (3H, m)

6) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(purin-6-yl)carbonylaminobenzamide

5 NMR (DMSO-d₆, δ) : 1.40-1.50 (2H, m), 1.50-1.62 (2H, m), 1.70-1.79 (2H, m), 2.14 (3H, s), 2.18-2.36 (9H, m), 3.20 (3H, s), 3.35-3.43 (6H, m), 3.75 (3H, s), 3.81-3.99 (2H, m), 6.65 (1H, d, J=7Hz), 6.82 (1H, s), 6.95-7.08 (3H, m), 8.27 (1H, d, J=6Hz), 8.83 (1H, s), 9.10 (1H, s)

10

7) 4-(3-Benzyl-2-methoxymethyl-3H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

15 NMR (CDCl₃, δ) : 1.49-1.64 (2H, m), 1.64-1.81 (2H, m), 1.81-1.95 (2H, m), 2.21-2.31 (6H, m), 2.31-2.44 (6H, m), 3.34 (3H, s), 3.43 (3H, s), 3.46-3.54 (2H, m), 3.58 (3H, s), 3.60-3.70 (2H, m), 3.90-4.04 (2H, m), 4.82 (2H, s), 5.63 (2H, s), 6.60-6.75 (6H, m), 6.75-6.83 (1H, m), 6.83-6.93 (2H, m), 6.99 (1H, d, J=8Hz), 7.19-7.31 (2H, m), 7.55 (1H, s), 7.90 (1H, d, J=8Hz), 8.20 (1H, d, J=8Hz)

20

8) 4-(1,2-Dimethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

25

30 NMR (CDCl₃, δ) : 1.45-1.76 (4H, m), 1.76-1.91 (2H, m), 2.25 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 2.69 (3H, s), 3.33 (3H, s), 3.45-3.52 (2H, m), 3.59-3.68 (2H, m), 3.78 (3H, s), 3.81-3.90 (4H, m), 3.90-4.01 (1H, m), 6.54-6.64 (2H, m), 6.86 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.01 (1H, s), 7.34 (1H, t, J=8Hz), 7.44 (1H, d, J=8Hz), 8.14 (1H, d, J=8Hz), 8.53 (1H, d, J=8Hz)

30

35 9) 4-(1-Ethyl-2-methyl-1H-benzimidazol-4-yl)carbonylamino-

3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

5 NMR (CDCl₃, δ) : 1.44 (3H, t, J=7Hz), 1.48-1.61 (2H, m), 1.65-1.75 (2H, m), 1.75-1.98 (2H, m), 2.27 (3H, s), 2.29 (3H, s), 2.31-2.43 (6H, m), 2.70 (3H, s), 3.34 (3H, s), 3.44-3.53 (2H, m), 3.59-3.68 (2H, m), 3.79-3.90 (4H, m), 3.90-4.00 (1H, m), 4.22 (2H, q, J=7Hz), 6.53-6.63 (2H, m), 6.86 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.01 (1H, s), 7.34 (1H, t, J=8Hz), 7.46 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz), 8.52 (1H, d, J=8Hz)

10 10) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-methyl-1-propyl-1H-benzimidazol-4-yl)carbonylamino benzamide

15 NMR (CDCl₃, δ) : 0.97 (3H, t, J=7Hz), 1.44-1.74 (4H, m), 1.74-1.92 (4H, m), 2.24 (3H, s), 2.27 (3H, s), 2.31-2.42 (6H, m), 2.67 (3H, s), 3.32 (3H, s), 3.43-3.53 (2H, m), 3.58-3.66 (2H, m), 3.76-3.90 (4H, m), 3.90-4.00 (1H, m), 4.13 (2H, t, J=7Hz), 6.52-6.62 (2H, m), 6.86 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.00 (1H, s-like), 7.33 (1H, t, J=8Hz), 7.45 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz), 8.53 (1H, d, J=8Hz)

25 11) 3-Methoxy-4-[1-(4-methoxybenzyl)-2-(N-methylcarbamoyl)-1H-benzimidazol-4-yl]carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

30 NMR (CDCl₃, δ) : 1.46-1.63 (2H, m), 1.63-1.77 (2H, m), 1.77-1.91 (2H, m), 2.25 (3H, s), 2.28 (3H, s), 2.31-2.41 (6H, m), 3.13 (3H, d, J=5Hz), 3.34 (3H, s), 3.44-3.51 (2H, m), 3.56-3.65 (2H, m), 3.74 (3H, s), 3.82-4.01 (5H, m), 6.00 (2H, s), 6.58 (1H, d, J=8Hz), 6.64 (1H, s), 6.82 (2H, d, J=8Hz), 6.87 (1H, d, J=8Hz), 6.97 (1H, d, J=8Hz), 7.10 (1H, s),

1.21 (2H, d, J=8Hz), 7.46 (1H, t, J=8Hz), 7.60 (1H, d, J=8Hz), 7.95 (1H, br peak), 8.24 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz)

- 5 12) 4-(2-tert-Butyldiphenylsiloxymethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.04 (9H, s), 1.35-1.49 (2H, m),
1.49-1.64 (2H, m), 1.64-1.80 (2H, m), 2.11 (3H, s),
10 2.13-2.25 (7H, m), 2.30 (2H, t, J=7.5Hz), 3.17 (3H, s), 3.27-3.45 (7H, m), 3.84 (1H, br peak), 3.96 (1H, br peak), 5.03 (2H, s), 6.63 (1H, d, J=8Hz), 6.80 (2H, s-like), 6.94 (1H, d, J=8Hz), 7.02 (1H, d, J=8Hz), 7.30-7.50 (7H, m), 7.69 (4H, d, J=8Hz),
15 7.79 (1H, d, J=8Hz), 7.91 (1H, d, J=8Hz), 8.35 (1H, d, J=8Hz)

- 13) 4-(Benzoxazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
20

NMR (CDCl₃, δ) : 1.47-1.65 (2H, m), 1.65-1.78 (2H, m),
1.78-1.92 (2H, m), 2.26 (3H, s), 2.30 (3H, s),
2.32-2.43 (6H, m), 3.35 (3H, s), 3.44-3.54 (2H, m),
3.58-3.68 (2H, m), 3.79-4.02 (5H, m), 6.54-6.66
25 (2H, m), 6.87 (1H, d, J=8Hz), 6.98 (1H, d, J=8Hz), 7.04 (1H, s-like), 7.55 (1H, t, J=8Hz), 7.77 (1H, d, J=8Hz), 8.23-8.31 (2H, m), 8.44 (1H, d, J=8Hz)

- 14) 4-(Benzoxazol-7-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
30

NMR (CDCl₃, δ) : 1.45-1.76 (4H, m), 1.76-1.90 (2H, m),
3.33 (3H, s), 3.43-3.52 (2H, m), 3.56-3.68 (2H, m),
3.83 (3H, s), 3.86-4.01 (2H, m), 6.54-6.65 (2H, m),
35 6.85 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.06 (1H,

s), 7.52 (1H, t, J=8Hz), 7.97 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz), 8.23 (1H, s), 8.36 (1H, d, J=8Hz), 9.50 (1H, s)

- 5 15) 4-(3-Bromo-2-methylimidazo[1,2-a]pyridin-8-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

10 NMR (CDCl₃, δ) : 1.44-1.61 (2H, m), 1.61-1.78 (2H, m), 1.78-1.92 (2H, m), 2.28 (3H, s), 2.30 (3H, s), 2.33-2.45 (3H, m), 2.54 (3H, s), 3.33 (3H, s), 3.45-3.55 (2H, m), 3.60-3.70 (2H, m), 3.88 (3H, s), 3.91-4.01 (2H, m), 6.53-6.64 (2H, m), 6.86 (1H, d, J=8Hz), 6.93-7.03 (2H, m), 7.07 (1H, t, J=8Hz), 8.19 (1H, d, J=8Hz), 8.25 (1H, d, J=8Hz), 8.45 (1H, d, J=8Hz)

- 16) 3-Methoxy-N-methyl-4-(2-methylimidazo[1,2-a]pyridin-4-yl)carbonylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide

20 NMR (CDCl₃, δ) : 1.45-1.65 (2H, m), 1.65-1.75 (2H, m), 1.75-1.90 (2H, m), 2.25 (3H, s), 2.29 (3H, s), 2.31-2.42 (6H, m), 2.52 (3H, s), 3.33 (3H, s), 3.42-3.52 (2H, m), 3.57-3.66 (2H, m), 3.77-3.90 (4H, m), 3.90-4.02 (1H, m), 6.51-6.62 (2H, m), 6.80-7.04 (4H, m), 7.41 (1H, s), 8.16 (2H, d-like), 8.48 (1H, d, J=8Hz)

Example 3

To a suspension of 3-methyl-3H-benzimidazole-4-carboxylic acid hydrochloride (112 mg) in dichloromethane 3. (2 ml) was added oxalyl chloride (79 mg) in an ice water bath under nitrogen and then added 1 drop of N,N-dimethylformamide. After being stirred under the same condition for 2 hours, the reaction mixture was concentrated in vacuo. The residue was added to a solution of 4-amino-3-methoxy-N-

methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (150 mg) in pyridine (2 ml) under nitrogen at ambient temperature and the mixture was stirred for 2 hours and allowed to stand at same temperature overnight. The reaction mixture was concentrated in vacuo and the residue was dissolved in chloroform. The solution was washed with water and saturated aqueous sodium bicarbonate solution and the organic layer was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by preparative thin-layer chromatography (ethyl acetate-methanol = 1:1) to give 3-methoxy-N-methyl-4-(3-methyl-3H-benzimidazol-4-yl)carboxylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (52 mg) as a powder.

NMR (CDCl₃, δ) : 1.45-1.63 (2H, m), 1.63-1.79 (2H, m), 1.79-1.91 (2H, m), 2.30 (6H, s), 2.32-2.43 (6H, m), 3.33 (3H, s), 3.43-3.52 (2H, m), 3.52-3.68 (2H, m), 3.75 (3H, s), 3.83-4.03 (5H, m), 6.55-6.72 (2H, m), 6.88 (1H, d, J=8Hz), 6.97 (1H, d, J=8Hz), 7.04 (1H, s), 7.22-7.35 (1H, m), 7.50 (1H, d, J=8Hz), 7.88 (1H, s), 7.95 (1H, d, J=8Hz), 8.30 (1H, d, J=6Hz), 8.38 (1H, s)

Example 4

To a solution of 4-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (6.0 g) in 1,4-dioxane (200 ml) was added 8-nitro-2-trifluoromethyl-3,1-benzoxazin-4-one (3.24 g) and the mixture was stirred at 100°C for 4 hours. To the mixture was added 8-nitro-2-trifluoromethyl-3,1-benzoxazin-4-one (3.24 g) and the solution was stirred at 100°C for additional 3 hours. To the mixture was added 1N sodium hydroxide solution (90 ml) and the resulting solution was stirred at 60°C for 1 hour. After being concentrated in vacuo, the residue was diluted with chloroform and the organic solution was washed with

saturated aqueous sodium bicarbonate solution and brine. The organic layer was dried over magnesium sulfate and the solvent was concentrated in vacuo to give 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(3-nitro-2-trifluoroacetylaminobenzoyl)-aminobenzamide as a yellow powder (10.6 g). The crude product was used for next step without further purification.

Example 5

To a solution of 4-[(indol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (160 mg) in N,N-dimethylformamide (3.0 ml) was added portionwise potassium tert-butoxide (37.3 mg) at 0°C and the mixture was stirred at 0°C for 1 hour. Methyl iodide (47.2 mg) was added to the mixture and the solution was stirred at 0°C for 1 hour. The reaction was quenched with water and then the aqueous solution was extracted with ethyl acetate. Drying, filtering and removal of solvents afforded a crude product. The crude product was purified by column chromatography (eluent; 2% methanol in chloroform) to give 4-[(1-methylindol-4-yl)-carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (65 mg) as a white syrup.

NMR (CDCl₃, δ) : 1.49-1.89 (6H, m), 2.28 (3H, s), 2.29 (3H, s), 2.32-2.42 (6H, m), 3.34 (3H, s), 3.46-3.52 (2H, m), 3.60-3.68 (2H, m), 3.78 (3H, s), 3.86 (3H, s), 3.88-3.99 (2H, m), 6.60 (1H, d, J=8Hz), 6.64 (1H, s), 6.87 (1H, d, J=8Hz), 6.91-6.98 (2H, m), 7.06 (1H, s), 7.21 (1H, d, J=3Hz), 7.27-7.32 (1H, m), 7.49 (1H, d, J=8Hz), 7.63 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz), 8.76 (1H, s)

Example 6

The following compound was obtained according to a

similar manner to that of Example 5.

4-[(1-Isopropylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.48-1.59 (2H, m), 1.57 (6H, d, J=7Hz), 1.66-1.88 (4H, m), 2.29 (6H, s), 2.32-2.40 (6H, m), 3.33 (3H, s), 3.46-3.51 (2H, m), 3.60-3.67 (2H, m), 3.79 (3H, s), 3.88-4.00 (2H, m), 4.68-4.78 (1H, m), 6.60 (1H, d, J=8Hz), 6.65 (1H, s), 6.87 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 6.97 (1H, d, J=3Hz), 7.07 (1H, s), 7.27 (1H, t, J=8Hz), 7.38 (1H, d, J=3Hz), 7.54 (1H, d, J=8Hz), 7.62 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz), 8.76 (1H, s)

Example 7

To a solution of 4-(2-amino-3-nitrobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (3.88 g) in ethanol (40 ml) were added a solution of ammonium chloride (385 mg) in water (10 ml) and iron powder (2.01 g) and the mixture was stirred at 100°C for 2 hours. The mixture was filtered through a bed of celite and the filtrate was concentrated in vacuo. The residue was diluted with ethyl acetate and the solution was washed with aqueous saturated sodium bicarbonate solution and brine. The organic layer was dried over magnesium sulfate and the solution was concentrated in vacuo to give 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide as a yellow powder (3.42 g).

NMR (CDCl₃, δ) : 1.47-1.59 (2H, m), 1.59-1.90 (4H, m), 2.29 (3H, s), 2.30 (3H, s), 2.33-2.42 (6H, m), 3.33 (3H, s), 3.47-3.50 (2H, m), 3.62-3.67 (2H, m), 3.77 (3H, s), 3.82-4.00 (2H, m), 6.57-6.68 (3H, m), 6.80-7.03 (5H, m), 8.20 (1H, d, J=7Hz), 8.44 (1H,

s)

Example 8

To a suspension of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg) in water (5 ml) was added 1N hydrochloric acid (1.3 ml) and then dicyandiamide (545 mg) was added to the stirred reaction mixture. The solution was heated under reflux for 24 hours. After cooling, aqueous sodium hydrogen carbonate was added to the mixture and extracted with ethyl acetate. The extract was washed with brine and dried over sodium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography (SiO₂, 30 g, 15% methanol in chloroform) to give 4-[2-guanidinobenzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (100 mg) as yellow amorphous.

NMR (CDCl₃, δ) : 1.36-1.83 (6H, m), 2.25 (3H, s), 2.30 (3H, s), 2.32-2.48 (6H, m), 3.34 (3H, s), 3.43-3.74 (5H, m), 3.78 (3H, s), 3.82-3.98 (1H, m), 6.56 (1H, s), 6.68 (1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.02 (1H, d, J=8Hz), 7.08-7.18 (2H, m), 7.36 (1H, d, J=8Hz), 7.97 (1H, d, J=8Hz), 8.44 (1H, d, J=8Hz)

Example 9

To a solution of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (110 mg) in anhydrous tetrahydrofuran (2 ml) was added 1,1'-thiocarbonyldiimidazole (48 mg) under nitrogen at ambient temperature and stirred at same temperature for 1 day. After being concentrated in vacuo, the residue was diluted with a mixture of chloroform and saturated sodium bicarbonate aqueous solution and the organic layer was separated. The

organic layer was washed with water and brine and the solution was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by preparative thin-layer chromatography (ethyl acetate-methanol = 1:1) to give 4-(2-mercapto-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (96 mg) as a powder.

NMR (CDCl₃, δ) : 1.48-1.62 (2H, m), 1.66-1.78 (2H, m),
1.78-1.90 (2H, m), 2.28 (3H, s), 2.31 (3H, s),
2.33-2.46 (6H, m), 3.33 (3H, s), 3.45-3.53 (2H, m),
3.60-3.70 (2H, m), 3.81 (3H, s), 3.84-4.01 (2H, m),
6.55-6.67 (2H, m), 6.86 (1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.03 (1H, s), 7.12 (2H, s-like), 7.17-
7.40 (2H, m), 7.70 (1H, s), 8.20 (1H, d, J=8Hz),
8.65 (1H, s)

Example 10

To a suspension of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (242 mg) in water (3 ml) was added cyanogen bromide (46 mg) at ambient temperature. The mixture was stirred at the same temperature for 2 hours and then allowed to stand at the same temperature overnight. To the reaction mixture was added saturated aqueous sodium bicarbonate solution and the solution was extracted with chloroform. The organic layer was washed with water, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 4-(2-amino-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (67 mg) as a powder.

NMR (CDCl₃, δ) : 1.40-1.56 (2H, m), 1.56-1.87 (4H, m),
2.23 (3H, s), 2.26 (3H, s), 2.30-2.44 (6H, m), 3.33

(3H, s), 3.41-3.53 (2H, m), 3.53-3.69 (5H, m),
3.69-3.83 (1H, m), 3.83-4.00 (1H, m), 5.54 (2H, br
peak), 6.50-6.66 (2H, m), 6.80-6.95 (2H, m), 6.95-
7.10 (2H, m), 7.69 (1H, d-like), 8.37 (1H, d-like)

5

Example 11

To 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide (120 mg) were added acetic acid (47
10 mg) and water (0.5 ml) and the suspension was stirred at
ambient temperature until a clear solution was obtained.
After being cooled to 5°C a cold solution of sodium nitrite
(15 mg) in water (0.3 ml) was added all at once to the
solution. The reaction mixture was stirred at 5°C for 5
15 minutes and then the temperature was raised to 75°C and
stirred for 10 minutes. The reaction mixture was cooled to
20°C and the solution was stirred in an ice water bath for 1
hour. To the reaction mixture were added saturated aqueous
sodium bicarbonate solution and chloroform and the organic
20 layer was separated. The organic layer was washed with
saturated aqueous sodium bicarbonate solution and brine and
dried over magnesium sulfate. The solvent was evaporated and
the residue was purified by preparative thin-layer
chromatography (chloroform-methanol = 10:1) to give 4-(1H-
25 benzotriazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide (81 mg) to give a powder.

NMR (CDCl₃, δ) : 1.46-1.62 (2H, m), 1.65-1.76 (2H, m),
1.76-1.90 (2H, m), 2.26 (3H, s), 2.32-2.45 (7H, m),
30 2.45-2.59 (2H, m), 3.35 (3H, s), 3.50-3.66 (3H, m),
3.73-3.89 (5H, m), 3.89-3.99 (1H, m), 6.57-6.65
(2H, m), 6.93 (1H, d, J=8Hz), 6.99-7.05 (2H, m),
7.53 (1H, t, J=8Hz), 8.00 (1H, d, J=8Hz), 8.10 (1H,
d, J=8Hz), 8.34 (1H, d, J=8Hz), 10.04 (1H, s)

35

Example 12

To a stirred solution of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (145 mg) in a mixture of acetonitrile and benzene [1:4(v/v)] was added methoxycarbonyl isothiocyanate (36 mg) and the reaction mixture was stirred at ambient temperature for 5 minutes. After being added 1,3-dicyclohexylcarbodiimide (73 mg) to the solution, the resulting mixture was stirred at reflux temperature for 5 hours. The reaction mixture was concentrated in vacuo and the residue was dissolved in chloroform. The solution was washed with saturated aqueous sodium bicarbonate solution, water and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by chromatography on silica gel (chromatorex) eluting with chloroform and preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 3-methoxy-4-(2-methoxycarbonylamino-1H-benzimidazol-4-yl)carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (82 mg) as a powder.

NMR (DMSO-d₆, δ) : 1.36-1.49 (2H, m), 1.49-1.62 (2H, m), 1.67-1.83 (2H, m), 2.13 (3H, s), 2.15-2.38 (9H, m), 3.20 (3H, s), 3.36-3.45 (4H, m), 3.74 (3H, s), 3.79-3.90 (4H, m), 3.90-4.03 (1H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.89 (1H, s), 6.93 (1H, d, J=8Hz), 7.03 (1H, d, J=8Hz), 7.20 (1H, t J=8Hz), 7.67 (1H, d, J=8Hz), 7.82 (1H, d, J=8Hz), 8.21-8.28 (1H, m)

Example 13

A mixture of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (90 mg) and trimethyl orthoformate (1 ml) was refluxed for 4 hours. After removing excess

reagent by evaporation, the residue was dissolved in chloroform and the solution was washed with water and saturated aqueous sodium bicarbonate solution. The organic layer was dried over magnesium sulfate and the solvent was evaporated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 4-(1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (51 mg) as a powder.

10 NMR (CDCl₃, δ) : 1.48-1.62 (2H, m), 1.67-1.78 (2H, m),
1.78-1.91 (2H, m), 2.21-2.31 (6H, m), 2.31-2.43
(6H, m), 3.35 (3H, s), 3.45-3.56 (2H, m), 3.60-3.69
(2H, m), 3.81 (1H, d-like), 3.81-3.90 (1H, m),
3.90-4.01 (1H, m), 6.54-6.65 (2H, m), 6.84-6.93
15 (1H, m), 6.93-7.07 (2H, m), 7.31-7.50 (1H, m), 7.59
(1H x 1/2, d, J=8Hz), 7.68 (1H x 1/2, d, J=8Hz),
7.98-8.33 (2H, m), 8.45-8.56 (1H x 1/2, m), 8.79
(1H x 1/2, s)

20 Example 14

To a solution of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg) in acetic acid (1 ml) was added tetramethyl orthocarbonate (66 mg) at ambient temperature and the solution was allowed to stand at the same temperature for 3 days. After being concentrated in vacuo, the residue was diluted with chloroform and saturated sodium bicarbonate aqueous solution. The organic layer was separated and washed with water and brine. The solution was dried over magnesium sulfate and the solvent was evaporated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 4-(2-methoxy-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (171 mg) as a powder.

35

NMR (CDCl₃, δ) : 1.44-1.65 (2H, m), 1.65-1.76 (2H, m),
1.76-1.90 (2H, m), 2.27 (3H, s), 2.29 (3H, s),
2.33-2.45 (6H, m), 3.34 (3H, s), 3.42-3.54 (2H, m),
3.58-3.68 (2H, m), 3.71 (3H x 2/3, s), 3.80 (3H x
1/3, s), 3.82-4.02 (2H, m), 4.20 (3H x 1/3, s),
4.28 (3H x 2/3, s), 6.53-6.68 (2H, m), 6.81-7.08
(3H, m), 7.17-7.43 (3H, m), 7.70 (1H x 1/3, d,
J=8Hz), 8.06 (1H x 2/3, d, J=8Hz), 8.22-8.31 (1H x
1/3, m), 8.54 (1H x 2/3, d, J=8Hz), 8.72 (1H x 1/3,
s), 8.90 (1H x 2/3, s)

Example 15

The following compounds were obtained according to a
similar manner to that of Example 14.

1) 4-(2-Ethoxy-1H-benzimidazol-4-yl)carbonylamino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.44-1.65 (5H, m), 1.65-1.76 (2H, m),
1.76-1.90 (2H, m), 2.25 (3H, s), 2.29 (3H, s),
2.31-2.43 (6H, m), 3.33 (3H, s), 3.44-3.53 (2H, m),
3.60-3.68 (2H, m), 3.74 (3H x 3/4, s), 3.80 (3H x
1/4, s), 3.83-4.01 (2H, m), 4.61 (2H x 1/4, q,
J=7.5Hz), 4.73 (2H x 3/4, q, J=7.5Hz), 6.55-6.66
(2H, m), 6.81-6.99 (2H, m), 7.02 (1H, s), 7.20 (1H,
t, J=8Hz), 7.35 (1H, d, J=8Hz), 7.67 (1H x 1/4, d,
J=8Hz), 8.05 (1H x 3/4, d, J=8Hz), 8.26 (1H x 1/4,
d, J=8Hz), 8.54 (1H x 3/4, d, J=8Hz), 8.70-8.78
(1H, m)

2) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-propoxy-1H-
benzimidazol-4-yl)carbonylaminobenzamide

NMR (CD₃OD, δ) : 1.06 (3H, t, J=7.5Hz), 1.42-1.56 (2H,
m), 1.56-1.71 (2H, m), 1.71-1.96 (4H, m), 2.20 (3H,

5 s), 2.25 (3H, s), 2.28-2.97 (6H, m), 3.29 (3H, s),
3.40-3.62 (4H, m), 3.70 (3H, s), 3.79-4.01 (2H, m),
4.56 (2H, t, J=7Hz), 6.67 (1H, d, J=8Hz), 6.76 (1H,
s), 6.94 (1H, s), 6.96-7.05 (2H, m), 7.13 (1H, t,
J=8Hz), 7.35 (1H, d, J=8Hz), 7.82 (1H, d, J=8Hz),
8.40 (1H, d, J=8Hz)

Example 16

10 A suspension of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-
methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-
1-yloxy]phenyl]benzamide (100 mg) in acetic acid (1 ml) was
refluxed for 8 hours. After being evaporated in vacuo, the
residue was dissolved chloroform and the solution was washed
15 with water and saturated aqueous sodium bicarbonate solution.
The organic layer was dried over magnesium sulfate and the
solvent was evaporated in vacuo. The residue was purified by
preparative thin-layer chromatography (chloroform-methanol =
10:1) to give 3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-
4-yl)carbonylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
20 carbonylpent-1-yloxy]phenyl]benzamide (84 mg) as a powder.

NMR (CDCl₃, δ) : 1.43-1.60 (2H, m), 1.60-1.75 (2H, m),
1.75-1.89 (2H, m), 2.25 (3H, s), 2.29 (3H, s),
2.31-2.43 (6H, m), 2.63 (3H, s), 3.33 (3H, s),
3.43-3.53 (2H, m), 3.58-3.68 (2H, m), 3.73-3.90
25 (4H, m), 3.90-4.00 (1H, m), 6.53-6.64 (2H, m),
6.82-6.91 (1H, m), 6.91-7.05 (2H, m), 7.22-7.33 (1H
x 2/3, m), 7.43-7.53 (1H, m), 7.80-7.90 (1H x 1/3,
m), 8.11 (1H x 2/3, d, J=8Hz), 8.23-8.31 (1H x 1/3,
m), 8.47-8.57 (1H x 2/3, m), 8.75 (1H x 1/3, s),
30 9.83 (1H x 2/3, s), 10.67 (1H x 1/3, s)

Example 17

The following compounds were obtained according to a
similar manner to that of Example 16.

1) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-trifluoromethyl-1H-benzimidazol-4-yl)carbonylaminobenzamide

5 NMR (CDCl₃, δ) : 1.43-1.62 (2H, m), 1.62-1.77 (2H, m),
1.77-1.94 (2H, m), 3.35 (3H, s), 3.43-3.58 (2H, m),
3.58-3.70 (2H, m), 3.80 (3H, s), 3.82-3.91 (1H, m),
3.91-4.01 (1H, m), 6.53-6.66 (2H, m), 6.90 (1H, d,
J=8Hz), 6.94-7.04 (2H, m), 7.48 (1H, t, J=8Hz),
7.78 (1H, br peak), 8.08 (1H, br peak), 8.41 (1H,
10 br peak)

2) 4-(2-Ethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide

15 NMR (CDCl₃, δ) : 1.41-1.75 (7H, m), 1.75-1.90 (2H, m),
2.23-2.31 (6H, m), 2.31-2.42 (6H, m), 2.99 (2H, q,
J=7.5Hz), 3.34 (3H, m), 3.44-3.52 (2H, m), 3.59-
3.67 (2H, m), 3.76-3.90 (4H, m), 3.90-4.00 (1H, m),
6.53-6.64 (2H, m), 6.83-7.04 (3H, m), 7.24-7.34
20 (1H, m), 7.44-7.55 (1H, m), 7.89 (1H x 1/3, d,
J=8Hz), 8.14 (1H x 2/3, d, J=8Hz), 8.28 (1H x 1/3,
d, J=8Hz), 8.37 (1H x 2/3, d, J=8Hz), 8.78 (1H x
1/3, s), 9.56 (1H x 2/3, s), 10.75 (1H x 1/3, s)

25 3) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-n-propyl-1H-benzimidazol-4-yl)carbonylaminobenzamide

NMR (CD₃OD, δ) : 1.06 (3H, t, J=7.5Hz), 1.47-1.60 (2H,
m), 1.60-1.74 (2H, m), 1.74-1.90 (2H, m), 1.90-2.05
30 (2H, m), 2.23 (3H, s), 2.28 (3H, s), 2.31-2.47 (6H,
m), 2.95 (2H, t, J=7.5Hz), 3.30 (3H, s), 3.48-3.61
(4H, m), 3.79 (3H, s), 3.84-3.94 (1H, m), 3.94-4.06
(1H, m), 6.70 (1H, d, J=8Hz), 6.79 (1H, s), 6.97
(1H, s), 6.99-7.08 (2H, m), 7.30 (1H, t, J=8Hz),
35 7.94 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

4) 4-(2-Isopropyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.43-1.59 (8H, m), 1.59-1.75 (2H, m),
5 1.75-1.90 (2H, m), 2.22-2.30 (6H, m), 2.30-2.42
(6H, m), 3.18-3.30 (1H, m), 3.33 (3H, s), 3.43-3.51
(2H, m), 3.60-3.67 (2H, m), 3.75-3.89 (4H, m),
3.89-4.01 (1H, m), 6.53-6.65 (2H, m), 6.87 (1H, d,
J=8Hz), 6.94 (1H, d, J=8Hz), 7.03 (1H, s), 7.30
10 (1H, t, J=8Hz), 7.45-7.56 (1H, m), 7.90 (1H x 1/3,
d, J=8Hz), 8.12 (1H x 2/3, d, J=8Hz), 8.26 (1H x
1/3, d, J=8Hz), 8.54 (1H x 2/3, d, J=8Hz), 8.77 (1H
x 1/3, s), 9.64 (1H x 2/3, s)

15 Example 18

To a solution of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (640 mg) in dichloromethane (6 ml) were added pyridine (99 mg) and
20 N-phthaloylglycyl chloride (280 mg) under nitrogen in ice water bath and the mixture was stirred at the same temperature for 2 hours. To the reaction mixture was added methanol (1 ml) at ambient temperature and stirred for additional 30 minutes. The reaction mixture was concentrated
25 in vacuo and the residue was dissolved in pyridine (6 ml). The solution was stirred at 100°C for 48 hours and the solvent was evaporated in vacuo. The residue was diluted with chloroform and saturated aqueous sodium bicarbonate solution. The organic layer was separated and washed with
30 water and brine. The organic solution was dried over magnesium sulfate and the solvent was evaporated in vacuo. The residue was chromatographed on silica gel eluting with a mixture of chloroform and methanol (50:1-25:1-10:1) to give 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-phthalimidomethyl-1H-

35

benzimidazol-4-yl)carbonylaminobenzamide (410 mg) as a powder.

NMR (CDCl₃, δ) : 1.43-1.61 (2H, m), 1.61-1.76 (2H, m),
1.76-1.89 (2H, m), 2.27 (3H, s), 2.30 (3H, s),
5 2.32-2.43 (6H, m), 3.33 (3H, s), 3.45-3.53 (2H, m),
3.59-3.67 (2H, m), 3.81-4.02 (5H, m), 5.18 (2H, s),
6.54-6.65 (2H, m), 6.77-7.03 (3H, m), 7.27-7.38
(1H, m), 7.50 (1H, d, J=8Hz), 7.64-7.77 (2H, m),
7.77-7.90 (2H, m), 8.10 (1H, d, J=8Hz), 8.46 (1H,
10 d, J=8Hz)

Example 19

The following compounds were obtained according to a similar manner to that of Example 18.

15

1) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(2-phthalimidoethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.46-1.66 (2H, m), 1.66-1.79 (2H, m),
20 1.79-1.92 (2H, m), 2.20-2.32 (6H, m), 2.32-2.43
(6H, m), 3.33 (3H, s), 3.40 (2H, t, J=7Hz), 3.44-
3.55 (2H, m), 3.55-3.68 (2H, m), 3.81-4.03 (5H, m),
4.23 (2H, t, J=7Hz), 6.55-6.69 (2H, m), 6.81-7.05
(3H, m), 7.34 (1H, t, J=8Hz), 7.53-7.62 (2H, m),
25 7.65-7.76 (2H, m), 7.76-7.91 (1H, m), 8.12 (1H, d,
J=8Hz), 8.42 (1H, d, J=8Hz)

2) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(2-pyridylmethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.46-1.62 (2H, m), 1.62-1.91 (4H, m),
2.22-2.30 (6H, m), 2.30-2.42 (6H, m), 3.34 (3H, s),
3.44-3.52 (2H, m), 3.58-3.67 (2H, m), 3.77-4.02
(5H, m), 4.47 (2H x 1/6, s), 4.52 (2H x 5/6, s),
35 6.54-6.65 (2H, m), 6.87 (1H, d, J=8Hz), 6.96 (1H,

d, J=8Hz), 7.01-7.09 (1H, m), 7.16-7.35 (overlapped in CHCl₃), 7.40 (1H x 5/6, d, J=8Hz), 7.48 (1H x 1/6, d, J=8Hz), 7.58 (1H x 5/6, d, J=8Hz), 7.63 (1H x 1/6, d, J=8Hz), 7.66-7.76 (1H, m), 7.89 (1H x 1/6, d, J=8Hz), 8.13 (1H x 5/6, d, J=8Hz), 8.30 (1H x 1/6, d, J=8Hz), 8.51 (1H x 5/6, d, J=8Hz), 8.67 (1H x 5/6, d, J=5Hz), 8.73 (1H x 1/5, s-like)

3) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(3-pyridylmethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.45-1.58 (2H, m), 1.64-1.75 (2H, m), 1.75-1.89 (2H, m), 2.25 (3H, s), 2.29 (3H, s), 2.32-2.43 (6H, m), 3.34 (3H, s), 3.43-3.53 (2H, m), 3.57-3.66 (2H, m), 3.74 (3H x 2/3, s), 3.79 (3H x 1/3, s), 3.81-3.91 (1H, m), 3.90-4.00 (1H, m), 4.32 (2H, s), 6.54-6.64 (2H, m), 6.83-6.92 (1H, m), 6.92-7.03 (2H, m), 7.23-7.36 (overlapped in CHCl₃), 7.50 (1H, d, J=8Hz), 7.66 (1H x 1/3, d, J=8Hz), 7.73 (1H x 2/3, d, J=8Hz), 7.91 (1H x 1/3, d, J=8Hz), 8.15 (1H x 2/3, d, J=8Hz), 8.21 (1H x 1/3, d, J=8Hz), 8.48-8.75 (3H, m)

4) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-pyridylmethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃-CD₃OD, δ) : 1.47-1.60 (2H, m), 1.60-1.76 (2H, m), 1.76-1.89 (2H, m), 2.26 (3H, s), 2.30 (3H, s), 2.32-2.44 (6H, m), 3.33 (3H, s), 3.46-3.54 (2H, m), 3.58-3.64 (2H, m), 3.68 (3H x 3/4, s), 3.77 (3H x 1/4, s), 3.81-4.01 (2H, m), 4.33 (2H, s), 6.55-6.65 (2H, m), 6.83-6.92 (1H, m), 6.92-7.02 (2H, m), 7.27-7.38 (3H, m), 7.53 (1H, d, J=8Hz), 8.10 (1H, d, J=8Hz), 8.45-8.55 (3H, m)

Example 20

To a mixture of 4-[(indol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (150 mg) and
5 N,N-dimethylaminopyridine (5.9 mg) in acetonitrile (10 ml) was added diethyl dicarbonate (46.6 mg) at ambient temperature. The solution was stirred at ambient temperature for a few hours and stood overnight. The resulting mixture was diluted with water and the aqueous solution was extracted
10 with ethyl acetate. Drying, filtering and removal of solvents afforded a crude product. The crude product was purified by column chromatography (eluent; 2% methanol in chloroform) to give 4-[(1-ethoxycarbonylindol-4-yl)carbonyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
15 1-yl)carbonylpent-1-yloxy]phenyl]benzamide (126 mg) as a colorless syrup.

NMR (CDCl₃, δ) : 1.49 (3H, t, J=8Hz), 1.50-1.59 (2H, m), 1.66-1.76 (2H, m), 1.79-1.88 (2H, m), 2.28 (3H, s), 2.29 (3H, s), 2.32-2.41 (6H, m), 3.33 (3H, s),
20 3.47-3.51 (2H, m), 3.60-3.66 (2H, m), 3.79 (3H, s), 3.88-4.00 (2H, m), 4.50 (2H, q, J=8Hz), 6.59 (1H, d, J=8Hz), 6.64 (1H, s), 6.87 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.05 (1H, s), 7.18 (1H, d, J=4Hz), 7.40 (1H, t, J=8Hz), 7.62 (1H, d, J=8Hz), 7.75 (1H, d, J=4Hz),
25 8.32 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz), 8.57 (1H, s)

Example 21

30 A solution of 4-[[1-(4-toluenesulfonyl)indol-4-yl]-carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (255 mg) in a mixture of 2N potassium hydroxide aqueous solution (2.5 ml) and methanol (6.0 ml) was stirred at
35 ambient temperature for 3 hours and stood overnight. The

resulting mixture was diluted with water and the solution was extracted with ethyl acetate. Drying, filtering and removal of solvents afforded a crude product. The crude product was chromatographed on silica gel (eluent; 3-8% methanol in chloroform) to give 4-[(indol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide (154 mg) as a white amorphous powder.

NMR (CDCl₃, δ) : 1.48-1.59 (2H, m), 1.66-1.88 (4H, m), 2.29 (3H, s), 2.30 (3H, s), 2.32-2.47 (6H, m), 3.33 (3H, s), 3.48-3.53 (2H, m), 3.62-3.69 (2H, m), 3.77 (3H, s), 3.84-4.00 (2H, m), 6.60 (1H, d, J=8Hz), 6.63 (1H, s), 6.88 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 6.99-7.05 (2H, m), 7.21-7.28 (1H, m), 7.35 (1H, d, J=3Hz), 7.55 (1H, d, J=8Hz), 7.61 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz), 8.73 (1H, s), 8.82-8.87 (1H, br s)

Example 22

To a solution of 4-[(1-pivaloyloxymethylindol-7-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (100 mg) in methanol (4.0 ml) was added 28% sodium methylate in methanol solution (100 mg). The solution was stirred at ambient temperature for a few hours and stood overnight. The resulting mixture was diluted with water and the solution was extracted with ethyl acetate. Drying, filtering and removal of solvents afforded 4-[(indol-7-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide (70 mg) as a slightly yellow syrup.

NMR (CDCl₃, δ) : 1.49-1.95 (6H, m), 2.27 (3H, s), 2.28 (3H, s), 2.32-2.41 (6H, m), 3.33 (3H, s), 3.47-3.51 (2H, m), 3.60-3.67 (2H, m), 3.80 (3H, m), 3.83-3.95 (2H, m), 6.56-6.65 (3H, m), 6.87 (1H, d, J=8Hz),

6.98 (1H, d, J=8Hz), 7.03 (1H, s), 7.16 (1H, t, J=8Hz), 7.32 (1H, d, J=4Hz), 7.49 (1H, d, J=8Hz), 7.83 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz), 8.78 (1H, s)

5

Example 23

The mixture of 4-[(1-tert-butoxycarbonyl-2-ethoxycarbonylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (75 mg) in trifluoroacetic acid (2.0 ml) was stirred at ambient temperature for 5 minutes. Trifluoroacetic acid was removed in vacuo and the residue was diluted with aqueous saturated sodium bicarbonate solution. The solution was extracted with ethyl acetate and the organic layer was washed with brine. Drying, filtering and removal of solvents afforded 4-[(2-ethoxycarbonylindol-4-yl)-carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (50 mg) as a yellow amorphous powder.

20 NMR (CDCl₃, δ) : 1.42 (3H, t, J=8Hz), 1.49-1.60 (2H, m), 1.68-1.90 (4H, m), 2.28 (3H, s), 2.29 (3H, s), 2.32-2.43 (6H, m), 3.33 (3H, s), 3.47-3.52 (2H, m), 3.60-3.68 (2H, m), 3.80 (3H, s), 3.87-4.00 (2H, m), 4.41 (2H, q, J=8Hz), 6.60 (1H, d, J=8Hz), 6.63 (1H, s), 6.87 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.06 (1H, s), 7.39 (1H, t, J=8Hz), 7.58-7.66 (2H, m), 7.75 (1H, s), 8.35 (1H, d, J=8Hz), 8.68 (1H, s), 9.22-9.28 (1H, br s)

30 Example 24

A mixture of 4-(3-benzyl-2-methoxymethyl-3H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg) and 5% formic acid in methanol solution (10 ml) was refluxed for 24 hours. The reaction

35

mixture was concentrated in vacuo and the residue was dissolved in chloroform. The solution was washed with saturated aqueous sodium bicarbonate solution and the organic layer was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 3-methoxy-4-(2-methoxymethyl-1H-benzimidazol-4-yl)carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (48 mg) as a powder.

NMR (CDCl₃, δ) : 1.24-1.60 (2H, m), 1.60-1.93 (6H, m), 3.33 (3H, s), 3.42-3.60 (5H, m), 3.60-3.76 (2H, m), 3.76-4.04 (5H, m), 4.77 (2H x 1/3, s), 4.83 (2H x 2/3, s), 6.51-6.67 (2H, m), 6.67-7.10 (3H, m), 7.23-7.56 (2H, m), 7.62 (1H x 2/3, d, J=8Hz), 7.85-7.96 (1H x 1/3, m), 7.17 (1H x 2/3, d, J=8Hz), 8.24-8.33 (1H x 1/3, m), 8.49 (1H x 2/3, d, J=8Hz), 8.72-8.80 (1H x 1/3, m), 9.72 (1H x 2/3, s), 10.91 (1H x 1/3, s)

Example 25

A solution of 3-methoxy-4-[1-(4-methoxybenzyl)-2-(N-methylcarbamoyl)-1H-benzimidazol-4-yl]carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (130 mg) in trifluoroacetic acid (3 ml) was stirred at 60°C for 8 hours. The reaction mixture was concentrated in vacuo and the residue was diluted with a mixture of chloroform and saturated aqueous sodium bicarbonate solution. The organic layer was separated and washed with water and brine. The solution was dried over magnesium sulfate and the solvent was evaporated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 3-methoxy-N-methyl-4-[2-(N-methylcarbamoyl)-1H-benzimidazol-4-yl]carbonylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-

carbonylpent-1-yloxy]phenyl]benzamide (45 mg) as a powder.

NMR (CDCl₃, δ) : 1.46-1.76 (4H, m), 1.76-1.90 (2H, m),
2.27 (3H, s), 2.29 (3H, s), 2.31-2.42 (6H, m), 3.19
(3H, d, J=5Hz), 3.34 (3H, s), 3.44-3.51 (2H, m),
3.58-3.65 (2H, m), 3.80-4.01 (5H, m), 6.59 (1H, d,
J=8Hz), 6.64 (1H, s), 6.86 (1H, d, J=8Hz), 6.96
(1H, d, J=8Hz), 7.09 (1H, s), 7.50 (1H, t, J=8Hz),
7.66 (1H, br peak), 7.73 (1H, d, J=8Hz), 8.25 (1H,
d, J=8Hz), 8.51 (1H, d, J=8Hz)

Example 26

To a solution of 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-phthalimidomethyl-1H-benzimidazol-4-yl)carbonylaminobenzamide (383 mg) in ethanol (5 ml) was added hydrazine monohydrate (19 mg) at ambient temperature and the solution was stirred at 60°C for 1 hour. The reaction mixture was stirred in an ice water bath for 2 hours and the precipitate was filtered off. The filtrate was evaporated in vacuo and the residue was purified by preparative thin-layer chromatography (chloroform-methanol-ammonia solution (28%) = 160:32:1) to give 4-(2-aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg) as a powder.

NMR (CDCl₃, δ) : 1.45-1.62 (2H, m), 1.64-1.75 (2H, m),
1.75-1.90 (2H, m), 2.25 (3H, s), 2.29 (3H, s),
2.31-2.48 (6H, m), 3.33 (3H, s), 3.44-3.54 (2H, m),
3.58-3.67 (2H, m), 3.71 (3H, s), 3.79-3.90 (1H, m),
3.90-4.02 (1H, m), 4.20 (2H, br peak), 6.53-6.65
(2H, m), 6.83-7.04 (3H, m), 7.22-7.32 (1H, m), 7.54
(1H, br peak), 7.98 (1H, br peak), 8.47 (1H, br
peak)

Example 27

The following compound was obtained according to a similar manner to that of Example 26.

4-[2-(2-Aminoethyl)-1H-benzimidazol-4-yl]carbonylamino-
3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.43-1.58 (2H, m), 1.58-1.90 (4H, m),
2.25 (3H, s), 2.28 (3H, s), 2.30-2.41 (6H, m),
3.08-3.16 (2H, m), 3.23-3.34 (5H, m), 3.43-3.51
(2H, m), 3.56-3.65 (2H, m), 3.76-3.88 (4H, m),
3.88-4.00 (1H, m), 6.54-6.63 (1H, m), 6.86 (1H, d, J=8Hz),
6.91-7.01 (2H, m), 7.28 (1H, t, J=8Hz),
7.60 (1H, d, J=8Hz), 8.00 (1H, br peak), 8.46 (1H, d, J=8Hz)

15

Example 28

A solution of 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(3-nitro-2-trifluoroacetylaminobenzoyl)aminobenzamide (10.5 g) in
hydrazine monohydrate (100 ml) was stirred at 60°C for 2
hours and the mixture was diluted with a mixture of water and
ethyl acetate. The organic layer was separated and washed
with saturated aqueous sodium bicarbonate solution and brine.
The organic layer was dried over magnesium sulfate and the
solution was concentrated in vacuo. The residue was purified
by silica gel column chromatography (3% methanol in
chloroform) to give 4-(2-amino-3-nitrobenzoyl)amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]benzamide as a yellow powder
(5.90 g).

30

NMR (CDCl₃, δ) : 1.47-1.58 (2H, m), 1.60-1.88 (4H, m),
2.28 (3H, s), 2.30 (3H, s), 2.33-2.41 (6H, m), 3.34
(3H, s), 3.48-3.50 (2H, m), 3.62-3.66 (2H, m), 3.78
(3H, s), 3.83-3.97 (2H, m), 6.58-6.72 (3H, m), 6.87
(1H, d, J=6Hz), 6.97 (1H, d, J=6Hz), 7.02 (1H, s),

35

7.21 (1H, d, J=6Hz), 8.12-8.15 (3H, m), 8.29-8.33 (2H, m)

Example 29

5 The solution of 4-[(2-ethoxycarbonylindol-4-yl)-
carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
(100 mg) in a mixture of 1N aqueous sodium hydroxide solution
(0.43 ml) and ethanol (4.0 ml) was stirred at ambient
10 temperature for 5.5 hours. The resulting solution was
neutralized with 1N hydrochloric acid and methanol was
removed in vacuo. The residue was diluted with water and the
aqueous layer was extracted with chloroform. Drying,
filtering and removal of solvents afforded 4-[(2-
15 carboxyindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide (70 mg) as a yellow amorphous.

NMR (DMSO-d₆, δ) : 1.38-1.58 (4H, m), 1.69-1.79 (2H,
m), 2.19 (3H, s), 2.23 (3H, s), 2.26-2.37 (6H, m),
20 3.19 (3H, s), 3.38-3.47 (3H, m), 3.70 (3H, s),
3.80-4.00 (3H, m), 6.66 (1H, d, J=8Hz), 6.83 (1H,
s), 6.90-6.97 (2H, m), 7.03 (1H, d, J=8Hz), 7.33
(1H, t, J=8Hz), 7.47 (1H, s), 7.58-7.67 (2H, m),
7.90 (1H, d, J=8Hz), 9.20 (1H, s)

Example 30

To a mixture of 4-[(2-carboxyindol-4-yl)carbonyl]amino-
3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]benzamide (60 mg), N,N-dimethyl-
30 amine hydrochloride (7.7 mg) and 1-hydroxybenzotriazole (14.5
mg) in N,N-dimethylformamide (3.0 ml) was added a solution of
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
(20.6 mg) in N,N-dimethylformamide (1.0 ml) and the mixture
was stirred at ambient temperature for 4 hours. The
35 resulting mixture was diluted with ethyl acetate and the

organic layer was washed successively with saturated aqueous sodium bicarbonate solution and brine. Drying, filtering and removal of solvents afforded 4-[(2-dimethylaminocarbonyl-indol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-
5 [5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide (60 mg) as a yellow amorphous powder.

NMR (CDCl₃, δ) : 1.50-1.61 (2H, m), 1.68-1.90 (4H, m),
2.29 (6H, s), 2.32-2.42 (6H, m), 3.18-3.27 (3H, br
s), 3.34 (3H, s), 3.43-3.52 (5H, m), 3.60-3.68 (2H,
10 m), 3.78 (3H, s), 3.87-4.00 (2H, m), 6.60 (1H, d,
J=8Hz), 6.64 (1H, s), 6.87 (1H, d, J=8Hz), 6.96
(1H, d, J=8Hz), 7.07 (1H, s), 7.34 (1H, t, J=8Hz),
7.45 (1H, s), 7.52 (1H, d, J=8Hz), 7.61 (1H, d,
J=8Hz), 8.33 (1H, d, J=8Hz), 8.62 (1H, s), 9.73-
15 9.78 (1H, br s)

Example 31

A solution of 3-methoxy-4-(2-methoxy-1H-benzimidazol-4-yl)carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-
20 methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (139 mg) in 10% hydrogen chloride in methanol (2 ml) was stirred at ambient temperature for 2 hours and 4N hydrogen chloride in 1,4-dioxane (2 ml) was added to the mixture. After being allowed to stand at ambient temperature
25 overnight, the reaction mixture was concentrated in vacuo and the residue was diluted with a mixture of chloroform and saturated aqueous sodium bicarbonate solution. The organic layer was separated and washed with water and brine. The solution was dried over magnesium sulfate and evaporated in
30 vacuo. The residue was purified by preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 4-(2-hydroxy-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (53 mg) as a powder.

35 NMR (CDCl₃, δ) : 1.44-1.65 (2H, m), 1.65-1.76 (2H, m),

1.76-1.92 (2H, m), 2.28 (3H, s), 2.30 (3H, s),
2.31-2.43 (6H, m), 3.33 (3H, s), 3.44-3.53 (2H, m),
3.58-3.69 (2H, m), 3.82 (3H, s), 3.85-4.01 (2H, m),
6.55-6.68 (2H, m), 6.87 (1H, d, J=8Hz), 6.96 (1H,
5 d, J=8Hz), 7.05 (1H, s), 7.08-7.21 (2H, m), 8.25
(1H, d, J=8Hz), 8.50 (1H, s), 8.66 (1H, s), 9.43
(1H, br s)

Example 32

10 To a solution of 4-(2-tert-butylldiphenylsiloxymethyl-1H-
benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide (296 mg) in dry tetrahydrofuran (10
ml) was added tetrabutylammonium fluoride (173 mg) in ice
15 water bath under nitrogen and the mixture was stirred at the
same temperature for 1 hour. The reaction mixture was
concentrated in vacuo and the residue was dissolved in
chloroform. The solution was washed with water and brine and
dried over magnesium sulfate. The solvent was evaporated in
20 vacuo. The residue was purified by preparative thin-layer
chromatography (chloroform-methanol = 10:1) to give 4-(2-
hydroxymethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-
methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-
1-yloxy]phenyl]benzamide (200 mg) as a powder.

25 NMR (CDCl₃, δ) : 1.37-1.58 (2H, m), 1.58-1.90 (4H, m),
2.23 (3H, s), 2.28 (3H, s), 2.30-2.44 (6H, m), 3.32
(3H, s), 3.40-3.71 (7H, m), 3.71-3.88 (1H, m),
3.88-4.01 (1H, m), 4.85 (2H, s), 6.58 (2H, s-like),
6.67-6.79 (1H, m), 6.79-6.97 (2H, m), 7.14-7.30
30 (1H, m), 7.49 (1H, br peak), 7.98 (1H, br peak),
8.36 (1H, d, J=8Hz)

Example 33

A mixture of 4-(2-amino-1H-benzimidazol-4-yl)-
35 carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-

methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
(45 mg) and acetic anhydride (0.5 ml) was stirred at ambient
temperature for 1 hour and then allowed to stand at the same
temperature overnight. The reaction mixture was concentrated
5 in vacuo and the residue was dissolved in chloroform. The
solution was washed successively with saturated aqueous
sodium bicarbonate solution, water and brine and the solution
was dried over magnesium sulfate. The solvent was evaporated
in vacuo and the residue was purified by preparative thin-
10 layer chromatography (chloroform-methanol = 10:1) to give
4-(2-acetamido-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-
N-methyl-N-[4-methyl-2-[5-(4-methyloperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]benzamide (26 mg) to give a
powder.

15 NMR (CDCl₃, δ) : 1.36-1.52 (2H, br peak), 1.58-1.82
(4H, m), 2.26 (3H, s), 2.28-2.42 (15H, m), 3.32
(3H, s), 3.43-3.51 (2H, m), 3.51-3.66 (5H, m), 3.75
(1H, br peak), 3.91 (1H, br peak), 6.55-6.65 (2H,
m), 6.84-6.95 (2H, m), 6.99 (1H, s), 7.20-7.31 (1H,
20 m), 7.54 (1H, br peak), 8.07 (1H, br peak), 8.34
(1H, d, J=8Hz)

Example 34

To a solution of 4-(2-amino-1H-benzimidazol-4-
25 yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
(90 mg) in pyridine (1 ml) was added methanesulfonyl chloride
(18 mg) in ice water bath under nitrogen and the mixture was
stirred at the same temperature for 3 hours. The reaction
30 mixture was concentrated in vacuo and the residue was
dissolved in chloroform. The organic layer was washed
successively with saturated aqueous sodium bicarbonate
solution, water and brine and the solution was dried over
magnesium sulfate. The solvent was evaporated in vacuo and
35 the residue was purified by preparative thin-layer

chromatography (chloroform-methanol = 10:1) to give 4-(2-methanesulfonylamino-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (15 mg) as a powder.

5 NMR (CDCl₃, δ) : 1.42-1.58 (2H, m), 1.62-1.95 (4H, m),
2.26 (3H, s), 2.29-2.45 (9H, m), 3.24 (3H, s), 3.34
(3H, s), 3.44-3.52 (2H, m), 3.59-3.67 (2H, m),
3.76-3.89 (4H, m), 3.89-4.00 (1H, m), 6.98 (1H, br
s), 6.54-6.65 (2H, m), 6.88 (1H, d, J=8Hz), 6.94
10 (1H, d, J=8Hz), 7.04 (1H, s), 7.23 (1H, t, J=8Hz),
7.74 (1H, d, J=8Hz), 8.11 (1H, d, J=8Hz), 8.40 (1H,
d, J=8Hz)

Example 35

15 The following compound was obtained according to a
similar manner to that of Example 34.

4-(2-Benzenesulfonylamino-1H-benzimidazol-4-yl)carbonyl-
amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
20 1-yl)carbonylpent-1-yloxy]phenyl]benzamide

 NMR (CDCl₃, δ) : 1.43-1.73 (4H, m), 1.73-1.84 (2H, m),
2.25 (3H, s), 2.30 (3H, s), 2.31-2.42 (6H, m), 3.33
(3H, s), 3.44-3.51 (2H, m), 3.59-3.66 (2H, m),
3.75-3.87 (4H, m), 3.87-4.00 (1H, m), 6.04 (2H, s),
25 6.54-6.63 (2H, m), 6.86 (1H, d, J=8Hz), 6.93 (1H,
d, J=8Hz), 7.20 (1H, t, J=8Hz), 7.48-7.56 (2H, m),
7.65 (1H, t, J=8Hz), 7.85 (1H, d, J=8Hz), 7.97 (2H,
d, J=8Hz), 8.08 (1H, d, J=8Hz), 8.37 (1H, d,
J=8Hz), 11.39 (1H, s)

30

Example 36

To a solution of 4-[(indol-4-yl)carbonyl]amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide (93 mg) in
35 dichloromethane (6.0 ml) was added N,N-dimethylmethylen-

ammonium chloride (41.7 mg) at 0°C and the mixture was stirred at ambient temperature for 1 hour. The resulting mixture was diluted with water and the aqueous solution was extracted with dichloromethane. Drying, filtering and
5 removal of solvents afforded 4-[(3-dimethylaminomethylindol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (90 mg) as a colorless syrup.

NMR (CDCl₃, δ) : 1.50-1.95 (6H, m), 2.28 (3H, s), 2.29
10 (6H, s), 2.30 (3H, s), 2.33-2.42 (6H, m), 3.34 (3H, s), 3.44-3.52 (2H, m), 3.60-3.66 (2H, m), 3.68 (3H, s), 3.84-4.00 (2H, m), 5.30 (2H, s), 6.58 (1H, d, J=8Hz), 6.62 (1H, s), 6.83 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 6.97 (1H, s), 7.10-7.26 (2H, m),
15 7.35-7.57 (3H, m), 8.23-8.31 (1H, m), 8.36-8.40 (1H, br)

Example 37

To a mixture of 4-(2-hydroxymethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (40 mg), triethylamine (31 mg), dimethyl sulfoxide (0.5 ml) and dichloromethane (0.5 ml) was added portionwise sulfur trioxide pyridine complex (29 mg) in water bath and the
25 mixture was stirred at the same temperature for 1 day. The reaction mixture was poured into saturated aqueous sodium bicarbonate solution and the solution was extracted with chloroform. The organic layer was washed with water and brine and the solution was dried over magnesium sulfate. The
30 solvent was evaporated in vacuo and the residue was purified by preparative thin-layer chromatography (chloroform-methanol = 10:1) to give 4-(2-formyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (6
35 mg) as a powder.

NMR (CDCl₃, δ) : 1.47-1.64 (2H, m), 1.64-1.90 (4H, m),
2.25 (3H, s), 2.29 (3H, s), 2.34-2.44 (6H, m),
3.23-3.38 (3H, m), 3.46-3.54 (2H, m), 3.60-3.68
(2H, m), 3.73-4.01 (5H, m), 6.54-6.66 (2H, m),
5 6.83-7.10 (3H, m), 7.47-7.63 (1H, m), 7.63-7.77
(1H, m), 8.07-8.80 (3H, m), 9.96-10.12 (1H, m),
11.85 (1H, br s)

Example 38

10 To a solution of 4-(2-amino-3-nitrobenzoyl)amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide (50 mg) in ethanol
(2 ml) was added 1N hydrochloric acid (0.16 ml) at ambient
temperature and allowed to stand at the same temperature for
15 30 minutes. After being removed the solvent under reduced
pressure, the resulting solid was dissolved in distilled
water (5 ml) and the solution was filtered through micro
filter. The filtrate was lyophilized to give 4-[(2-amino-3-
nitro)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
20 methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
dihydrochloride (48 mg) as a white powder.

NMR (DMSO-d₆, δ) : 1.37-1.51 (2H, m), 1.51-1.65 (2H,
m), 1.65-1.84 (2H, m), 2.22 (3H, s), 2.33-2.45 (2H,
m), 2.77 (3H, s), 2.81-3.46 (9H, m), 3.61 (3H, s),
25 3.80-3.91 (1H, m), 3.91-4.00 (1H, m), 4.00-4.16
(1H, m), 4.23-4.51 (1H, m), 6.65 (1H, d, J=8Hz),
6.73 (1H, t, J=8Hz), 6.82 (1H, s-like), 6.86-6.96
(2H m), 7.05 (1H, d, J=8Hz), 7.52 (1H, d, J=8Hz),
7.96 (1H, d, J=8Hz), 7.99-8.08 (2H, m), 8.21 (1H,
30 d, J=8Hz), 9.63 (1H, s)

Example 39

The following compounds were obtained according to a
similar manner to that of Example 38.

1) 4-[(Indol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-phenyl]benzamide hydrochloride

5 NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.71-1.80 (2H, m), 2.25 (3H, s), 2.37-2.42 (2H, m), 2.73 (3H, s), 2.89-3.03 (3H, m), 3.20 (3H, s), 3.31-3.48 (3H, m), 3.70 (3H, s), 3.85-4.10 (3H, m), 4.39-4.49 (1H, m), 6.67 (1H, d, J=9Hz), 6.82 (2H, s), 6.90-6.98 (2H, m), 7.04 (1H, d, J=9Hz), 7.19 (1H, t, J=8Hz), 7.51-7.58 (2H, m), 7.62 (1H, d, J=8Hz), 8.00 (1H, d, J=8Hz), 9.05 (1H, s)

15 2) 4-[(1-Methylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

20 NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.71-1.80 (2H, m), 2.25 (3H, s), 2.40 (2H, t, J=8Hz), 2.73 (3H, s), 2.90-3.05 (3H, m), 3.19 (3H, s), 3.30-3.45 (3H, m), 3.70 (3H, s), 3.87 (3H, s), 3.88-4.00 (3H, m), 4.39-4.47 (1H, m), 6.67 (1H, d, J=8Hz), 6.80 (1H, d, J=3Hz), 6.83 (1H, s), 6.91-6.97 (2H, m), 7.06 (1H, d, J=8Hz), 7.27 (1H, t, J=8Hz), 7.51 (1H, d, J=3Hz), 7.58 (1H, d, J=8Hz), 7.69 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 9.07 (1H, s)

25 3) 4-[(1-Ethoxycarbonylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

30 NMR (DMSO-d₆, δ) : 1.41 (3H, t, J=8Hz), 1.42-1.63 (4H, m), 1.71-1.81 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=8Hz), 2.74 (3H, s), 2.86-3.04 (3H, m), 3.19 (3H, s), 3.30-3.48 (3H, m), 3.67 (3H, s), 3.83-4.00 (4H, m), 4.48 (2H, q, J=8Hz), 6.66 (1H, d, J=8Hz), 6.82 (1H, s), 6.90-6.96 (2H, m), 7.05 (1H, d, J=8Hz), 7.10 (1H, d, J=4Hz), 7.47 (1H, t, J=8Hz), 7.72-7.81

(2H, m), 7.83 (1H, d, J=4Hz), 8.31 (1H, d, J=8Hz),
9.33 (1H, s)

4) 4-[(1-Isopropylindol-4-yl)carbonyl]amino-3-methoxy-N-
5 methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-
1-yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d₆, δ) : 1.41-1.67 (4H, m), 1.48 (6H, d,
J=7Hz), 1.71-1.81 (2H, m), 2.23 (3H, s), 2.39 (2H,
t, J=8Hz), 2.73 (3H, s), 2.90-3.05 (3H, m), 3.19
10 (3H, s), 3.29-3.45 (3H, m), 3.69 (3H, s), 3.84-4.00
(4H, m), 4.80-4.90 (1H, m), 6.67 (1H, d, J=8Hz),
6.80-6.87 (2H, m), 6.90-6.97 (2H, m), 7.05 (1H, d,
J=8Hz), 7.25 (1H, t, J=8Hz), 7.55 (1H, d, J=8Hz),
7.69 (1H, d, J=3Hz), 7.77 (1H, d, J=8Hz), 7.98 (1H,
15 d, J=8Hz), 9.06 (1H, s)

5) 4-[(3-Dimethylaminomethylindol-4-yl)carbonyl]amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonyl]pent-1-yloxy]phenyl]benzamide dihydrochloride

20 NMR (DMSO-d₆, δ) : 1.42-1.64 (4H, m), 1.71-1.82 (2H,
m), 2.24 (3H, s), 2.40 (2H, t, J=8Hz), 2.70 (3H,
s), 2.72 (6H, s), 2.82-3.08 (4H, m), 3.20 (3H, s),
3.38-3.54 (3H, m), 3.68 (3H, s), 3.84-4.11 (3H, m),
5.63-5.72 (2H, br s), 6.66 (1H, d, J=8Hz), 6.83
25 (1H, s), 6.90-6.98 (2H, m), 7.04 (1H, d, J=8Hz),
7.27 (1H, t, J=8Hz), 7.38-7.52 (2H, m), 7.60-7.78
(2H, m), 8.02-8.18 (2H, m)

6) 4-[(Indol-7-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-
30 methyl-2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-
yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d₆, δ) : 1.41-1.62 (4H, m), 1.70-1.81 (2H,
m), 2.23 (3H, s), 2.39 (2H, t, J=8Hz), 2.74 (3H,
s), 2.88-3.02 (3H, m), 3.20 (3H, s), 3.30-3.42 (3H,
35 m), 3.66 (3H, s), 3.85-4.03 (3H, m), 4.38-4.47 (1H,

m), 6.51 (1H, d, J=3Hz), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.89-6.98 (2H, m), 7.06 (1H, d, J=8Hz), 7.10 (1H, d, J=8Hz), 7.37 (1H, d, J=3Hz), 7.71-7.80 (3H, m), 9.35 (1H, s)

5

7) 4-[(1-Methylindol-7-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

10 NMR (DMSO-d₆, δ) : 1.40-1.80 (6H, m), 2.22 (3H, s),
2.39 (2H, t, J=8Hz), 2.72 (3H, s), 2.91-3.02 (3H, m), 3.18 (3H, s), 3.31-3.48 (3H, m), 3.63 (3H, s), 3.73 (3H, s), 3.87-4.12 (3H, m), 4.39-4.48 (1H, m), 6.51 (1H, d, J=3Hz), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.89-7.09 (3H, m), 7.20-7.29 (2H, m), 7.37 (1H, d, J=3Hz), 7.68 (1H, d, J=8Hz), 7.80 (1H, d, J=8Hz), 9.53 (1H, s)

15

8) 4-[(2-Ethoxycarbonylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

20

25 NMR (DMSO-d₆, δ) : 1.37 (3H, t, J=8Hz), 1.43-1.63 (4H, m), 1.72-1.80 (2H, m), 2.24 (3H, s), 2.40 (2H, t, J=8Hz), 2.73 (3H, s), 2.88-3.03 (3H, m), 3.20 (3H, s), 3.28-3.45 (4H, m), 3.70 (3H, s), 3.87-4.00 (3H, m), 4.37 (2H, q, J=8Hz), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.90-6.97 (2H, m), 7.07 (1H, d, J=8Hz), 7.38 (1H, t, J=8Hz), 7.56 (1H, s), 7.61-7.69 (2H, m), 7.86 (1H, d, J=8Hz), 9.27 (1H, s)

30 9) 4-[(2-Dimethylaminocarbonylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

35 NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.72-1.81 (2H, m), 2.25 (3H, s), 2.40 (2H, t, J=8Hz), 2.74 (3H, s), 2.97-3.13 (3H, m), 3.19 (3H, s), 3.30-3.49

(10H, m), 3.68 (3H, s), 3.88-4.08 (3H, m), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.90-6.97 (2H, m), 7.05 (1H, d, J=8Hz), 7.21 (1H, s), 7.30 (1H, t, J=8Hz), 7.59 (1H, d, J=8Hz), 7.65 (1H, d, J=8Hz),
5 7.90 (1H, d, J=8Hz), 9.20 (1H, s)

10) 4-[1H-Imidazo[4,5-b]pyridin-7-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride

10 NMR (DMSO-d₆, δ) : 1.39-1.67 (4H, m), 1.69-1.83 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.72 and 2.74 (Total 3H, s), 2.80-3.11 (3H, m), 3.20 (3H, s), 3.31-3.57 (3H, m), 3.76 (3H, s), 3.80-4.77 (4H, m), 6.66 (1H, d, J=8Hz), 6.82 (1H, s), 6.89-7.01
15 (2H, m), 7.05 (1H, d, J=8Hz), 7.82 (1H, d, J=7Hz), 8.28 (1H, d, J=8Hz), 8.57 (1H, d, J=7Hz), 8.79 (1H, s)

11) 4-[2-Chloro-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

20 NMR (DMSO-d₆, δ) : 1.37-1.66 (4H, m), 1.67-1.83 (2H, m), 2.21 (3H, s), 2.39 (2H, t, J=7Hz), 2.73 (3H, s), 2.80-3.09 (3H, m), 3.19 (3H, s), 3.29-3.60 (3H, m), 3.71-4.17 (6H, m), 4.43 (1H, m), 6.64 (1H, d, J=8Hz), 6.81 (1H, s), 6.87-6.99 (2H, m), 7.03 (1H, d, J=8Hz), 7.41 (1H, dd, J=8, 8Hz), 7.72 (1H, d, J=8Hz), 7.96 (1H, d, J=8Hz), 8.33 (1H, d, J=8Hz)

12) 4-(2,3-Diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride

30 NMR (DMSO-d₆, δ) : 1.38-1.52 (2H, m), 1.52-1.66 (2H, m), 1.66-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.75 (3H, s), 2.81-3.11 (3H, m), 3.18 (3H,
35

5 s), 3.28-3.50 (3H, m), 3.63 (3H, s), 3.82-3.92 (1H, m), 3.92-4.03 (1H, m), 4.45 (1H, br peak), 6.60-6.77 (2H, m), 6.83 (1H, s-like), 6.87-6.98 (2H, m), 7.05 (1H, d, J=8Hz), 7.23 (1H, d, J=8Hz), 7.50 (1H, d, J=8Hz), 7.60 (1H, d, J=8Hz), 9.31 (1H, s)

13) 4-[2-Guanidinobenzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

10 NMR (DMSO-d₆, δ) : 1.31-1.84 (6H, m), 2.22 (3H, s),
2.40 (2H, t, J=7Hz), 2.74 (3H, s), 2.80-3.11 (3H, m), 3.18 (3H, s), 3.26-3.63 (3H, m), 3.74 (3H, s),
3.80-4.20 (3H, m), 4.33-4.53 (1H, m), 6.64 (1H, d, J=8Hz), 6.82 (1H, s), 6.92 (1H, d, J=8Hz), 6.99
15 (1H, s), 7.04 (1H, d, J=8Hz), 7.33 (1H, dd, J=8, 8Hz), 7.69 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz),
8.28-8.41 (1H, m), 8.49-8.80 (3H, m)

14) 4-(1H-Benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

20 NMR (DMSO-d₆, δ) : 1.40-1.51 (2H, m), 1.51-1.64 (2H, m), 1.69-1.81 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.49 (3H, s), 2.75 (3H, d-like), 2.80-3.07 (3H, m), 3.19 (3H, s), 3.31-3.48 (3H, m), 3.73 (3H, s), 3.77-4.03 (2H, m), 4.03-4.14 (1H, m),
25 4.38-4.50 (1H, m), 6.65 (1H, d, J=8Hz), 6.81 (1H, s-like), 6.87-6.98 (2H, m), 7.04 (1H, d, J=8Hz), 7.43 (1H, t, J=8Hz), 7.83 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 8.24 (1H, br peak), 8.64 (1H, br s),
30 10.63 (1H, br peak)

15) 4-(2-Hydroxy-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

35

NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.65 (2H, m), 1.69-1.82 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.76 (3H, d-like), 2.80-3.08 (3H, m), 3.19 (3H, s), 3.27-3.45 (3H, m), 3.64 (3H, s), 3.81-3.91 (1H, m), 3.91-4.01 (1H, m), 4.01-4.14 (1H, m), 4.38-4.50 (1H, m), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.86-6.96 (2H, m), 6.96-7.12 (3H, m), 7.46 (1H, d, J=8Hz), 7.66 (1H, d, J=8Hz) 9.28 (1H, s)

10

16) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.51 (2H, m), 1.52-1.65 (2H, m), 1.65-1.82 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.68 (3H, s), 2.75 (3H, d-like), 2.79-3.09 (3H, m), 3.18 (3H, s), 3.30-3.44 (3H, m), 3.72 (3H, s), 3.80-3.90 (1H, m), 3.90-4.00 (1H, m), 4.00-4.13 (1H, m), 4.48-4.50 (1H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s-like), 6.87-6.98 (2H, m), 7.04 (1H, d, J=8Hz), 7.39 (1H, br peak), 7.77 (1H, br peak), 7.95 (1H, br peak), 10.67 (1H, br s)

15

20

17) 4-(2-Mercapto-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.65 (2H, m), 1.70-1.82 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.76 (3H, s), 3.20 (3H, s), 3.65 (3H, s), 3.80-4.06 (2H, m), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 7.05 (1H, d, J=8Hz), 7.20 (1H, t, J=8Hz), 7.30 (1H, d, J=8Hz), 7.60 (1H, d, J=8Hz), 7.64-7.71 (2H, m), 9.11 (1H, s), 9.50 (1H, s)

30

35 18) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-

1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-trifluoromethyl-1H-benzimidazol-4-yl)carbonylaminobenzamide dihydrochloride

5 NMR (DMSO-d₆, δ) : 1.39-1.53 (2H, m), 1.53-1.65 (2H, m), 1.65-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7Hz), 2.78 (3H, s), 2.82-3.07 (3H, m), 3.20 (3H, s), 3.28-3.51 (3H, m), 3.75 (3H, s), 3.80-3.91 (1H, m), 3.91-4.03 (1H, m), 4.10 (1H, br peak), 4.43 (1H, br peak), 6.66 (1H, d, J=8Hz), 6.82 (1H, s), 6.90-7.00 (2H, m), 7.04 (1H, d, J=8Hz), 7.61 (1H, t, J=8Hz), 7.94 (1H, d, J=8Hz), 8.11 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz), 10.33 (1H, br peak), 11.89 (1H, s)

15 19) 4-(2-Amino-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

20 NMR (DMSO-d₆, δ) : 1.37-1.53 (2H, m), 1.53-1.65 (2H, m), 1.70-1.84 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.75 (3H, s), 2.80-3.10 (3H, m), 3.20 (3H, s), 3.26-3.55 (3H, m), 3.66 (3H, s), 3.78-3.92 (1H, m), 3.91-4.01 (1H, m), 4.07 (1H, br s), 4.43 (1H, br s), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.87-6.98 (2H, m), 7.04 (1H, d, J=8Hz), 7.29 (1H, br peak), 7.55 (1H, br peak), 7.86 (1H, br peak), 8.20 (1H, br peak), 9.75 (1H, br peak), 10.78 (1H, br peak)

30 20) 4-(2-Acetamido-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

35 NMR (CD₃OD, δ) : 1.50-1.65 (2H, m), 1.65-1.77 (2H, m), 1.77-1.92 (2H, m), 2.28 (3H, s), 2.35 (3H, s), 2.51 (2H, t-like), 2.93 (3H, s), 2.96-3.59 (6H, m), 3.75 (3H, s), 3.81-3.95 (1H, m), 3.95-4.07 (1H, m), 4.15-4.31 (1H, m), 4.57-4.71 (1H, m), 6.70 (1H, d,

J=8Hz), 6.80 (1H, s-like), 6.94 (1H, s-like), 7.00-7.12 (2H, m), 7.47-7.58 (2H, m), 7.81 (1H, d, J=8Hz), 7.99 (1H, d, J=8Hz), 8.06 (1H, d, J=8Hz)

- 5 21) 4-(2-Methanesulfonylamino-1H-benzimidazol-4-yl)-carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

10 NMR (CD₃OD, δ) : 1.50-1.64 (2H, m), 1.64-1.77 (2H, m), 1.77-1.90 (2H, m), 2.28 (3H, s), 2.50 (2H, t, J=7Hz), 2.92 (3H, s), 2.95-3.20 (3H, m), 3.39-3.60 (5H, m), 3.76 (3H, s), 3.83-3.94 (1H, m), 3.94-4.07 (1H, m), 4.17-4.32 (1H, m), 4.60-4.72 (1H, m), 6.70 (1H, d, J=8Hz), 6.79 (1H, s), 6.92 (1H, s), 6.99-15 7.10 (2H, m), 7.30 (1H, t, J=8Hz), 7.88 (1H, d, J=8Hz), 7.95 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz)

- 22) 4-(2-Benzenesulfonylamino-1H-benzimidazol-4-yl)-carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

25 NMR (DMSO-d₆, δ) : 1.35-1.50 (2H, m), 1.50-1.65 (2H, m), 1.65-1.80 (2H, m), 2.21 (3H, s), 2.38 (2H, t, J=7.5Hz), 2.75 (3H, d, J=5Hz), 2.80-3.10 (3H, m), 3.17 (3H, s), 3.25-3.48 (3H, m), 3.73 (3H, s), 3.78-3.90 (1H, m), 3.90-4.00 (1H, m), 4.07 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.54 (1H, d, J=8Hz), 6.78-6.98 (3H, m), 7.04 (1H, d, J=8Hz), 7.19 (1H, t, J=8Hz), 7.53-7.73 (3H, m), 7.73-7.94 30 (3H, m), 8.12 (2H, d, J=8Hz), 8.25 (1H, d, J=8Hz)

- 23) 4-(1H-Benzotriazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

35 NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.66 (2H,

- 5 m), 1.70-1.83 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=8Hz), 2.77 (3H, s), 2.83-3.10 (3H, m), 3.20 (3H, s), 3.25-3.53 (3H, m), 3.77 (3H, s), 3.82-3.93 (1H, m), 3.93-4.03 (1H, m), 4.08 (1H, br peak), 4.45 (1H, br peak), 6.66 (1H, d, J=8Hz), 6.82 (1H, s), 6.92-7.01 (2H, m), 7.06 (1H, d, J=8Hz), 7.72 (1H, br peak), 8.06-8.19 (2H, m), 8.36 (1H, br peak), 10.37 (1H, br peak), 11.56 (1H, br peak)
- 10 24) 4-(2-Ethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- 15 NMR (DMSO-d₆, δ) : 1.37-1.52 (5H, m), 1.52-1.65 (2H, m), 1.68-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.75 (3H, d, J=5Hz), 2.82-3.08 (5H, m), 3.18 (3H, s), 3.30-3.60 (3H, m), 3.73 (3H, s), 3.81-3.91 (1H, m), 3.91-4.02 (1H, m), 4.10 (1H, br d, J=15Hz), 4.45 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.89-6.99 (2H, m), 7.04 (1H, d, J=8Hz), 7.30-7.41 (1H, m), 7.73 (1H, d, J=8Hz), 7.92 (1H, d, J=8Hz), 8.33 (1H, br peak), 10.49 (1H, br s)
- 20
- 25 25) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-n-propyl-1H-benzimidazol-4-yl)carbonylamino benzamide dihydrochloride
- 30 NMR (DMSO-d₆, δ) : 0.99 (3H, t, J=7.5Hz), 1.38-1.52 (2H, m), 1.52-1.67 (2H, m), 1.67-1.83 (2H, m), 1.83-1.98 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.76 (3H, d, J=4Hz), 2.82-3.08 (5H, m), 3.20 (3H, s), 3.33-3.66 (3H, m), 3.73 (3H, s), 3.81-3.93 (1H, m), 3.93-4.03 (1H, m), 4.10 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.89-6.98 (2H, m), 7.04 (1H, d, J=8Hz), 7.31-7.43 (1H, m), 7.75 (1H, br d,
- 35

J=8Hz), 7.94 (1H, br d, J=8Hz), 8.26 (1H, br peak)

26) 4-(2-Isopropyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.37-1.51 (8H, m), 1.51-1.65 (2H, m), 1.65-1.84 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.76 (3H, d, J=4Hz), 2.83-3.09 (3H, m), 3.19 (3H, s), 3.23-3.43 (4H, m), 3.73 (3H, s), 3.81-3.92 (1H, m), 3.92-4.03 (1H, m), 4.09 (1H, br d, J=15Hz), 4.45 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.90-6.97 (2H, m), 7.04 (1H, d, J=8Hz), 7.34 (1H, t, J=8Hz), 7.71 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.38 (1H, br peak)

27) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.53 (2H, m), 1.53-1.66 (2H, m), 1.66-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.76 (3H, d-like), 2.82-3.10 (3H, m), 3.19 (3H, s), 3.75 (3H, s), 3.88 (1H, br peak), 3.98 (1H, br peak), 4.04-4.15 (1H, m), 4.38-4.50 (3H, m), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.95 (2H, br peak), 7.05 (1H, d, J=8Hz), 7.43 (1H, t, J=8Hz), 7.85 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 8.33 (1H, br peak), 8.75 (2H, br peak)

28) 4-[2-(2-Aminoethyl)-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.51 (2H, m), 1.51-1.65 (1H, m), 1.65-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.74 (3H, d, J=4Hz), 2.80-3.11 (3H, m), 3.30-3.50 (5H, m), 3.92-4.02 (1H, m), 4.10 (1H, br

5 d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.90-7.00 (2H, m), 7.05 (1H, d, J=8Hz), 7.36 (1H, t, J=8Hz), 7.75 (1H, d, J=8Hz), 7.93 (1H, d, J=8Hz), 8.16-8.31 (3H, m), 8.31-8.42 (1H, m)

29) 3-Methoxy-N-methyl-4-(3-methyl-3H-benzimidazol-4-yl)carbonylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

10 NMR (CD₃OD, δ) : 1.49-1.66 (2H, m), 1.66-1.79 (2H, m), 1.79-1.94 (2H, m), 2.27 (3H, s), 2.51 (2H, t, J=7.5Hz), 2.91 (3H, s), 2.95-3.35 (3H, m), 3.35-3.61 (3H, m), 3.71 (3H, s), 3.81-4.07 (2H, m), 4.10 (3H, s), 4.23 (1H, br peak), 4.65 (1H, br peak), 15 6.69 (1H, d, J=8Hz), 6.79 (1H, s-like), 6.97 (1H, s), 7.01-7.09 (2H, m), 7.70 (1H, t, J=8Hz), 7.82 (1H, d, J=8Hz), 7.95-8.08 (2H, m), 9.34 (1H, s)

30) 3-Methoxy-4-(2-methoxymethyl-1H-benzimidazol-4-yl)-carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

20 NMR (DMSO-d₆, δ) : 1.38-1.52 (2H, m), 1.52-1.65 (2H, m), 1.65-1.82 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.75 (3H, d-like), 2.80-3.09 (3H, m), 3.20 (3H, s), 3.32-3.53 (6H, m), 3.77 (3H, s), 3.81-3.91 (1H, m), 3.91-4.00 (1H, m), 4.04-4.14 (1H, m), 25 4.40-4.50 (1H, m), 4.80 (2H, s), 6.65 (1H, d, J=8Hz), 6.80 (1H, s-like), 6.89-6.98 (2H, m), 7.04 (1H, d, J=8Hz), 7.37 (1H, t, J=8Hz), 7.74 (1H, d, J=8Hz), 30 7.94 (1H, d, J=8Hz), 8.37 (1H, d, J=8Hz), 10.55 (1H, br s)

31) 4-(1,2-Dimethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride 35

NMR (DMSO-d₆, δ) : 1.38-1.53 (2H, m), 1.53-1.65 (2H, m), 1.65-1.82 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.70 (3H, s), 2.75 (3H, d-like), 2.80-3.08 (3H, m), 3.19 (3H, s), 3.31-3.50 (3H, m), 3.85 (3H, s), 3.92-4.01 (1H, m), 4.08 (1H, br d, J=12Hz), 4.44 (1H, br d, J=12Hz), 6.66 (1H, d, J=8Hz), 6.82 (1H, s), 6.90-7.00 (2H, m), 7.05 (1H, d, J=8Hz), 7.37-7.50 (1H, m), 7.83-7.92 (1H, m), 7.96 (1H, d, J=8Hz), 8.23 (1H, br peak), 10.78 (1H, br peak)

32) 4-(1-Ethyl-2-methyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.34 (3H, t, J=7Hz), 1.40-1.53 (2H, m), 1.53-1.66 (2H, m), 1.66-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.72 (3H, s), 2.75 (3H, d-like), 2.81-3.09 (2H, m), 3.20 (3H, s), 3.30-3.52 (4H, m), 3.75 (3H, s), 4.09 (1H, br d, J=12Hz), 4.35 (2H, q, J=7Hz), 4.44 (1H, br d, J=12Hz), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.90-6.99 (2H, m), 7.04 (1H, d, J=8Hz), 7.42 (1H, t, J=8Hz), 7.90 (1H, d, J=8Hz), 7.96 (1H, d, J=8Hz), 8.25 (1H, br peak)

33) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-methyl-1-propyl-1H-benzimidazol-4-yl)carbonylamino benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 0.90 (3H, t, J=7Hz), 1.40-1.52 (2H, m), 1.52-1.65 (2H, m), 1.68-1.86 (4H, m), 2.23 (3H, s), 2.39 (2H, t, J=7Hz), 2.70 (3H, s), 2.75 (3H, d, J=5Hz), 2.80-3.07 (3H, m), 3.18 (3H, m), 3.33-3.44 (3H, m), 3.76 (3H, s), 3.86 (1H, br peak), 3.95 (1H, br peak), 4.09 (1H, br d, J=12Hz), 4.27 (1H, t, J=8Hz), 4.44 (1H, br d, J=12Hz), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.88-6.98 (2H, m), 7.05 (1H, d, J=8Hz), 7.39 (1H, t, J=8Hz), 7.87 (1H, d,

J=8Hz), 7.94 (1H, d, J=8Hz), 8.30 (1H, br peak)

34) 3-Methoxy-N-methyl-4-[2-(N-methylcarbamoyl)-1H-benzimidazol-4-yl]carbonylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.65 (2H, m), 1.70-1.83 (2H, m), 2.21 (3H, s), 2.40 (2H, t, J=7Hz), 2.75 (3H, d, J=5Hz), 2.80-3.10 (6H, m), 3.20 (3H, s), 3.33-3.50 (3H, m), 3.76 (3H, s), 3.90 (1H, br peak), 3.97 (1H, br peak), 4.09 (1H, br d, J=12Hz), 4.44 (1H, br d, J=12Hz), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.91-6.99 (2H, m), 7.05 (1H, d, J=8Hz), 7.50 (1H, t, J=8Hz), 7.79 (1H, d, J=8Hz), 8.02 (1H, d, J=8Hz), 8.31 (1H, d, J=8Hz), 8.41 (1H, q-like)

35) 4-(2-Hydroxymethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.52 (2H, m), 1.52-1.67 (2H, m), 1.67-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.78 (3H, d, J=4Hz), 2.82-3.10 (3H, m), 3.20 (3H, s), 3.75 (3H, s), 3.80-3.91 (1H, m), 3.91-4.01 (1H, m), 4.01-4.17 (1H, m), 4.36-4.52 (1H, m), 4.86 (2H, s), 6.64 (1H, d, J=8Hz), 6.81 (1H, s), 6.85-6.98 (2H, m), 7.35 (1H, t, J=8Hz), 7.72 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.24-8.37 (1H, m), 10.40 (1H, br peak)

30

36) 4-(Benzoxazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.51 (2H, m), 1.51-1.65 (2H, m), 1.65-1.82 (2H, m), 2.24 (3H, s), 2.40 (2H, t,

35

J=7.5Hz), 2.77 (3H, s), 2.81-3.10 (3H, m), 3.20 (3H, s), 3.30-3.45 (3H, m), 3.77 (3H, s), 3.81-3.91 (1H, m), 3.91-4.04 (1H, m), 4.14-4.24 (1H, m), 4.40-4.52 (1H, m), 6.65 (1H, d, J=8Hz), 6.81 (1H, s-like), 6.91-6.99 (2H, m), 7.07 (1H, d, J=8Hz), 7.65 (1H, t, J=8Hz), 8.06-8.14 (2H, m), 8.32 (1H, d, J=8Hz), 9.14 (1H, s)

37) 3-Methoxy-N-methyl-4-(2-methylimidazo[1,2-a]pyridin-4-yl)carbonylamino-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.37-1.53 (2H, m), 1.53-1.65 (2H, m), 1.68-1.84 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.46 (3H, s), 2.75 (3H, d-like), 2.80-3.08 (3H, m), 3.20 (3H, s), 3.31-3.53 (3H, m), 3.92-4.01 (1H, m), 4.09 (1H, br d, J=12Hz), 4.45 (1H, br d, J=12Hz), 6.66 (1H, d, J=8Hz), 6.83 (1H, s-like), 6.90-7.00 (2H, m), 7.06 (1H, d, J=8Hz), 7.24 (1H, br peak), 7.99 (1H, br s), 8.07-8.25 (2H, m), 8.76-8.89 (1H, m), 10.94 (1H, br peak)

38) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(2-pyridylmethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.51 (2H, m), 1.51-1.64 (2H, m), 1.67-1.82 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.75 (3H, d, J=5Hz), 2.80-3.07 (3H, m), 3.20 (3H, s), 3.33-3.44 (3H, m), 3.64 (3H, s), 3.66-4.00 (overlapped in H₂O), 4.02-4.14 (1H, m), 4.39-4.50 (1H, m), 4.67 (2H, s), 6.65 (1H, s, J=8Hz), 6.81 (1H, s), 6.85-6.96 (2H, m), 7.03 (1H, d, J=8Hz), 7.38 (1H, t, J=8Hz), 7.48-7.57 (1H, m), 7.67 (1H, d, J=8Hz), 7.76 (1H, d, J=8Hz), 7.93 (1H, d, J=8Hz), 7.98-8.09 (1H, m), 8.20-8.29 (1H, m), 8.61-8.67 (1H, m)

39) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(3-pyridylmethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.51 (2H, m), 1.51-1.65 (2H, m), 1.65-1.83 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.74 (3H, s), 2.80-3.10 (3H, m), 3.20 (3H, s), 3.29-3.52 (3H, m), 3.66 (3H, s), 3.80-3.92 (1H, m), 3.92-4.02 (1H, m), 4.10 (1H, br peak), 4.43 (1H, br peak), 4.61 (2H, s), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.89-6.99 (2H, m), 7.04 (1H, d, J=8Hz), 7.37 (1H, t, J=8Hz), 7.75 (1H, d, J=8Hz), 7.86-7.99 (2H, m), 8.24-8.35 (1H, m), 8.44 (1H, d, J=8Hz), 8.76-8.86 (1H, m), 8.99 (1H, s-like)

40) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-pyridylmethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.51 (2H, m), 1.51-1.66 (2H, m), 1.66-1.83 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.74 (3H, s), 2.80-3.09 (3H, m), 3.19 (3H, s), 3.30-3.44 (3H, m), 3.63 (3H, s), 3.65-4.00 (overlapped in H₂O), 4.08 (1H, br peak), 4.44 (1H, br peak), 4.72 (2H, s), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.87-6.95 (2H, m), 7.03 (1H, d, J=8Hz), 7.39 (1H, t, J=8Hz), 7.78 (1H, d, J=8Hz), 7.94 (1H, d, J=8Hz), 8.05 (2H, d, J=5Hz), 8.25-8.35 (1H, m), 8.88 (2H, d, J=5Hz)

41) 3-Methoxy-4-(2-methoxycarbonylamino-1H-benzimidazol-4-yl)carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.66 (2H, m), 1.66-1.81 (2H, m), 2.23 (3H, s), 2.38 (2H, t,

J=7.5Hz), 2.75 (3H, d, J=5Hz), 2.81-3.08 (3H, m),
3.19 (3H, s), 3.32-3.47 (3H, m), 3.74 (3H, s),
3.79-3.91 (4H, m), 3.91-4.02 (4H, m), 4.07 (1H, br
d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.67 (1H, d,
5 J=8Hz), 6.82 (1H, s), 6.88 (1H, s), 6.94 (1H, d,
J=8Hz), 7.05 (1H, d, J=8Hz), 7.23 (1H, t, J=8Hz),
7.69 (1H, d, J=8Hz), 7.83 (1H, d, J=8Hz), 8.25 (1H,
d, J=8Hz)

10 Preparation 33

The following compound was obtained by using 2-amino-4-nitro-1H-benzimidazole as a starting compound according to a similar manner to that of Preparation 1.

15 2-[(Methylsulfonyl)amino]-4-nitro-1H-benzimidazole
NMR (DMSO-d₆, δ) : 3.61 (3H, s), 7.16 (1H, t, J=8Hz),
7.75 (2H, br peak), 7.83 (1H, d, J=8Hz), 7.95 (1H,
d, J=8Hz)

20 Preparation 34

The following compounds were obtained according to a similar manner to that of Preparation 5.

1) N-(2-Chloroethyl)-2-nitrobenzamide
25 NMR (CDCl₃, δ) : 3.72-3.89 (4H, m), 6.27 (1H, br), 7.56
(1H, m), 7.60 (1H, dd, J=8, 8Hz), 7.70 (1H, dd,
J=8, 8Hz), 8.10 (1H, d, J=8Hz)

2) N-(3-Chloropropyl)-2-nitrobenzamide
30 NMR (CDCl₃, δ) : 2.09-2.20 (2H, m), 3.56-3.73 (4H, m),
6.15 (1H, br s), 7.50 (1H, d, J=8Hz), 7.59 (1H, dd,
J=8, 8Hz), 7.67 (1H, dd, J=8, 8Hz), 8.05 (1H, d,
J=8Hz)

35 3) N-[1-(Hydroxymethyl)cyclopentyl]-2-nitrobenzamide

NMR (CDCl₃, δ) : 1.61-2.00 (8H, m), 3.62 (1H, t, J=7Hz), 3.80 (2H, d, J=7Hz), 6.00 (1H, s), 7.52 (1H, m), 7.59 (1H, m), 7.68 (1H, t, J=8, 8Hz), 8.07 (1H, d, J=8Hz)

5

Preparation 35

The following compounds were obtained according to a similar manner to that of Preparation 8.

- 10 1) Benzyl 1-tert-butoxycarbonyl-3-tert-butyldiphenyl-silyloxymethylindole-4-carboxylate

NMR (CDCl₃, δ) : 1.09 (9H, s), 1.67 (9H, s), 5.03 (2H, s), 5.19 (2H, s), 7.27-7.43 (12H, m), 7.66-7.78 (6H, m), 8.43 (1H, d, J=8Hz)

15

- 2) 2-tert-Butyldiphenylsiloxyethyl-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 1.15 (9H, s), 5.08 (2H, s), 7.30-7.51 (7H, m), 7.70 (4H, d-like), 8.00 (1H, d, J=8Hz), 8.15 (1H, d, J=8Hz)

20

Preparation 36

To a solution of benzyl 2-hydroxymethylindole-4-carboxylate (456 mg) and imidazole (364 mg) in N,N-dimethylformamide (10 ml) was added tert-butyldiphenylsilyl chloride (802 mg) and the solution was stirred at ambient temperature for 2 hours. The resulting mixture was diluted with ethyl acetate (30 ml) and washed successively with water and brine. The organic layer was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by silica gel column chromatography (eluent; n-hexane:ethyl acetate = 15:1) to give benzyl 2-tert-butyldiphenylsilyloxymethylindole-4-carboxylate (810 mg).

25

30

35

NMR (CDCl₃, δ) : 1.09 (9H, s), 4.93 (2H, s), 5.42 (2H, s), 6.89 (1H, s), 7.19 (1H, t, J=8Hz), 7.30-7.54

(12H, m), 7.68 (4H, d, J=8Hz), 7.92 (1H, d, J=8Hz),
8.32-8.37 (1H, br)

Preparation 37

5 The following compound was obtained by using methyl 2-formyl-1-methoxymethoxyindole-4-carboxylate as a starting compound according to a similar manner to that of Preparation 10.

10 Methyl 2-hydroxymethylindole-4-carboxylate
NMR (DMSO-d₆, δ) : 3.88 (3H, s), 4.67 (2H, d, J=6Hz),
5.37 (1H, t, J=6Hz), 6.80 (1H, s), 7.13 (1H, t,
J=8Hz), 7.60 (1H, d, J=8Hz), 7.69 (1H, d, J=8Hz)

15 Preparation 38

The following compounds were obtained according to a similar manner to that of Preparation 12.

- 1) 1-tert-Butoxycarbonyl-2-ethoxycarbonylindoline-4-carboxylic acid
20 NMR (DMSO-d₆, δ) : 1.22 (3H, t, J=8Hz), 1.45 (9H, s),
3.37 (1H, dd, J=8, 16Hz), 3.78 (1H, dd, J=10, 16Hz), 4.17 (2H, q, J=8Hz), 4.90 (1H, dd, J=8, 10Hz), 7.32 (1H, t, J=8Hz), 7.51 (1H, d, J=8Hz),
25 7.94-8.03 (1H, m)
- 2) 1-tert-Butoxycarbonyl-3-tert-butyldiphenylsilyloxy-methylindole-4-carboxylic acid
NMR (CDCl₃, δ) : 1.08 (9H, s), 1.68 (9H, s), 5.10 (2H, s), 7.28-7.41 (7H, m), 7.65-7.70 (4H, m), 7.77 (1H, s), 7.88 (1H, d, J=9Hz), 8.52 (1H, d, J=9Hz)
- 30 3) 1-tert-Butoxycarbonyl-2-phthalimidomethylindole-4-carboxylic acid
NMR (CDCl₃, δ) : 1.69 (9H, s), 5.15 (2H, s), 6.86 (1H,
- 35

s), 7.37 (1H, t, J=8Hz), 7.80 (1H, d, J=8Hz), 7.89-8.03 (4H, m), 8.28 (1H, d, J=8Hz)

4) 1-tert-Butoxycarbonyl-2-methylindole-4-carboxylic acid
5 NMR (CDCl₃, δ) : 1.70 (9H, s), 2.67 (3H, s), 7.13 (1H, s), 7.30 (1H, t, J=9Hz), 8.03 (1H, d, J=9Hz), 8.40 (1H, d, J=9Hz)

Preparation 39

10 To a solution of benzyl 1-tert-butoxycarbonyl-2-tert-butyl-
diphenylsilyloxymethylindole-4-carboxylate (762 mg) in
5.0% formic acid-methanol (20.0 ml) was added 10% palladium
on carbon (100 mg) and the mixture was stirred under nitrogen
atmosphere at ambient temperature for 2 hours. The resulting
15 solution was filtered through a bed of celite and the
filtrate was concentrated in vacuo. The residue was diluted
with chloroform (10 ml) and the solution was washed
successively with water and brine. The organic layer was
dried over magnesium sulfate and concentrated in vacuo to
20 give 1-tert-butoxycarbonyl-2-tert-butyl-
diphenylsilyloxy-methylindole-4-carboxylic acid (597 mg).

NMR (CDCl₃, δ) : 1.13 (9H, s), 1.49 (9H, s), 5.05 (2H, s),
7.29-7.43 (7H, m), 7.57 (1H, s), 7.71 (4H, d, J=8Hz),
8.08 (1H, d, J=8Hz), 8.44 (1H, d, J=8Hz)

25

Preparation 40

To a solution of benzyl 1-tert-butoxycarbonylindole-6-
carboxylate (1.27 g) in 5.0% formic acid-methanol (20.0 ml)
was added 10% palladium on carbon (1.27 g) and the mixture
30 was stirred under nitrogen atmosphere at ambient temperature
for 4 hours and stand overnight. The resulting solution was
filtered through a bed of celite and the filtrate was
concentrated in vacuo. The residue was diluted with
chloroform and the solution was washed successively with
35 water and brine. The organic layer was dried over magnesium

sulfate and concentrated in vacuo. The residue was triturated with ether:n-hexane (1:5) to give 1-tert-butoxycarbonylindoline-6-carboxylic acid (761 mg).

5 NMR (DMSO-d₆, δ) : 1.50 (9H, s), 3.11 (2H, t, J=11Hz),
3.96 (2H, t, J=11Hz), 7.28 (1H, d, J=8Hz), 7.54
(1H, d, J=8Hz), 8.20-8.30 (1H, br)

Preparation 41

10 To a solution of benzyl 1-tert-butoxycarbonylindole-6-carboxylate (450 mg) in 5.0% formic acid-methanol (10.0 ml) was added 10% palladium on carbon (153 mg) and the mixture was stirred under nitrogen atmosphere at ambient temperature for 4 hours. The resulting solution was filtered through a bed of celite and the filtrate was concentrated in vacuo.

15 The residue was diluted with chloroform and the solution was washed successively with water and brine. The organic layer was dried over magnesium sulfate and concentrated in vacuo. The residue was triturated with ether:n-hexane (1:5) to give 1-tert-butoxycarbonylindole-6-carboxylic acid (302 mg).

20 NMR (DMSO-d₆, δ) : 1.67 (9H, s), 6.80 (1H, d, J=3Hz),
7.69 (1H, d, J=8Hz), 7.83 (1H, d, J=8Hz), 7.86 (1H,
d, J=3Hz), 8.75 (1H, s)

Preparation 42

25 The following compound was obtained by using 3-methoxy-4-methoxycarbonyl-N-[4-methyl-2-(4-phthalimidobut-1-yloxy)]-phenylbenzamide as a starting compound according to a similar manner to that of Preparation 15.

30 3-Methoxy-4-methoxycarbonyl-N-methyl-N-[4-methyl-2-(4-phthalimidobut-1-yloxy)]phenylbenzamide

NMR (CDCl₃, δ) : 1.77-1.91 (4H, m), 2.23 (3H, s), 3.31
(3H, s), 3.69 (3H, s), 3.77 (2H, t, J=7.5Hz), 3.81
(3H, s), 3.82-3.99 (2H, m), 6.40-6.50 (2H, m), 6.82
35 (1H, d, J=8Hz), 6.88 (1H, d, J=8Hz), 6.90 (1H, s),

7.55 (1H, d, J=8Hz), 7.65-7.73 (2H, m), 7.81-7.88
(2H, m)

Preparation 43

5 The following compound was obtained according to a
similar manner to that of Preparation 42.

3-Methoxy-4-nitro-N-methyl-N-[4-methyl-2-(5-
phthalimidopent-1-yloxy)]phenylbenzamide

10 NMR (CDCl₃, δ) : 1.46-1.58 (2H, m), 1.70-1.91 (4H, m),
2.27 (3H, s), 3.30 (3H, s), 3.74 (2H, t, J=7.5Hz),
3.79 (3H, s), 3.85-3.95 (2H, m), 6.57-6.63 (2H, m),
6.82 (1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.06 (1H,
s), 7.61 (1H, d, J=8Hz), 7.68-7.75 (2H, m), 7.80-
15 7.85 (2H, m)

Preparation 44

The following compounds were obtained according to a
similar manner to that of Preparation 17.

20

1) 2-Benzyloxymethyl-1-tert-butoxycarbonylindoline-4-
carboxylic acid

25 NMR (CDCl₃, δ) : 1.50 (9H, s), 3.45-3.59 (3H, m), 3.67
(1H, dd, J=4, 10Hz), 4.52 (2H, s), 4.57-4.67 (1H,
br), 7.21-7.38 (7H, m), 7.70 (1H, d, J=8Hz)

2) 3-Formylindole-4-carboxylic acid

30 NMR (DMSO-d₆, δ) : 7.31 (1H, t, J=8Hz), 7.73 (2H, d,
J=8Hz), 8.27 (1H, d, J=3Hz), 10.45 (1H, s)

3) 2-Hydroxymethylindole-4-carboxylic acid

35 NMR (CDCl₃, δ) : 4.65 (2H, s), 5.27-5.40 (1H, br), 6.82
(1H, s), 7.10 (1H, t, J=8Hz), 7.57 (1H, d, J=8Hz),
7.67 (1H, d, J=8Hz)

- 4) 4-Carboxy-3-methoxy-N-methyl-N-(4-methyl-2-nitrophenyl)benzamide

NMR (CDCl₃, δ) : 2.24 (3H, s), 3.43 (3H, s), 3.97 (3H, s), 6.80 (1H, d, J=8Hz), 7.09 (1H, s), 7.20 (1H, d, J=8Hz), 7.34 (1H, d, J=8Hz), 7.60 (1H, s), 7.89 (1H, d, J=8Hz)

- 5) 4-Carboxy-3-methoxy-N-methyl-N-[2-(4-tert-butoxycarbonylaminobut-1-yl)oxy-4-methyl]phenylbenzamide

NMR (CDCl₃, δ) : 1.45 (9H, s), 1.60-1.72 (2H, m), 1.76-1.87 (2H, m), 2.26 (3H, s), 3.19 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.81-3.97 (2H, m), 3.89 (3H, s), 6.57-6.62 (2H, m), 7.85 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 7.07 (1H, s), 7.90 (1H, s)

- 6) 4-Carboxy-N-[2-[4,4-dimethyl(2,5-oxazolinyl)]phenyl]-N-methyl-3-methoxybenzamide

NMR (CDCl₃, δ) : 1.36 (3H, s), 1.37 (3H, s), 3.37 (3H, s), 3.81 (3H, s), 4.02-4.18 (2H, m), 7.01-7.18 (3H, m), 7.21-7.39 (2H, m), 7.79 (1H, m), 7.91 (1H, d, J=8Hz)

- 7) 4-Carboxy-3-methoxy-N-methyl-N-[2-(morpholin-4-yl)phenyl]benzamide

NMR (CDCl₃, δ) : 2.28-2.43 (2H, m), 2.78-2.92 (2H, m), 3.52 (3H, s), 3.59-3.85 (7H, m), 6.86 (1H, d, J=8Hz), 7.04 (1H, s), 7.08-7.18 (2H, m), 7.21 (1H, d, J=8Hz), 7.29 (1H, dd, J=8, 8Hz), 7.92 (1H, d, J=8Hz)

- 8) 4-Carboxy-3-methoxy-N-methyl-N-[2-(1-pyrrolyl)phenyl]benzamide

NMR (CDCl₃, δ) : 3.51 (3H, s), 3.88 (3H, s), 6.25 (2H, s), 6.32-6.41 (2H, m), 6.53 (1H, d, J=8Hz), 6.60 (1H, s), 7.11 (1H, m), 7.26-7.52 (3H, m), 7.75 (1H,

d, J=8Hz)

- 9) 4-Carboxy-3-methoxy-N-methyl-N-(2-piperidinophenyl)-benzamide

5 NMR (CDCl₃, δ) : 1.41-1.71 (6H, m), 2.20-2.36 (2H, m),
2.68-2.83 (2H, m), 3.52 (3H, s), 3.79 (3H, s), 6.83
(1H, d, J=8Hz), 6.96-7.30 (5H, m), 7.92 (1H, d,
J=8Hz)

- 10 10) 4-Carboxy-N-methyl-3-methoxy-N-[2-(4-methyl-1-piperazinyl)phenyl]benzamide

15 NMR (DMSO-d₆, δ) : 2.22 (3H, s), 2.23-2.50 (6H, m),
2.70-2.91 (2H, m), 3.36 (3H, s), 3.50 (3H, s), 6.81
(1H, s), 6.86-7.00 (2H, m), 7.10 (1H, m), 7.19 (1H,
m), 7.32-7.46 (2H, m)

- 11) 4-Carboxy-3-methoxy-N-methyl-N-[2-(2,5-oxazolyl)phenyl]-benzamide

20 NMR (CDCl₃, δ) : 3.46 (3H, s), 3.78 (3H, s), 6.77-6.85
(2H m), 7.23-7.46 (4H, m), 7.75-7.91 (3H, m)

- 12) 4-Carboxy-N-methyl-3-methoxy-N-[2-(2,5-oxazolinyl)-phenyl]benzamide

25 NMR (CDCl₃, δ) : 3.41 (3H, s), 3.82 (3H, s), 4.02-4.17
(2H, m), 4.32-4.50 (2H, m), 7.00-7.08 (2H, m), 7.15
(1H, d, J=8Hz), 7.28 (1H, dd, J=8, 8Hz), 7.38 (1H,
dd, J=8, 8Hz), 7.77 (1H, m), 7.89 (1H, d, J=8Hz)

- 13) 4-Carboxy-N-methyl-3-methoxy-N-[2-(3H,4H,5H-2,6-oxazinyl)phenyl]benzamide

30 NMR (CDCl₃, δ) : 1.94-2.08 (2H, m), 3.42 (3H, s), 3.58
(2H, t, J=7Hz), 3.78 (3H, s), 4.29-4.41 (2H, m),
6.84-7.37 (4H, m), 7.48-7.60 (2H, m), 7.90 (1H, d,
J=8Hz)

- 14) N-[2-(1-(Aza-3-oxaspiro[4.4]non-1-en-2-yl)phenyl)-4-carboxy-3-methoxy-N-methylbenzamide

NMR (CDCl₃, δ) : 1.61-1.77 (4H, m), 1.82-2.04 (4H, m),
3.38 (3H, s), 3.82 (3H, s), 4.16-4.28 (2H, m),
7.06-7.16 (3H, m), 7.22-7.39 (2H, m), 7.77 (1H, m),
7.92 (1H, d, J=8Hz)

- 15) 2-Carbamoyl-1-(4-methoxybenzyl)-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 3.69 (3H, s), 5.89 (3H, s), 6.86
(2H, d, J=8Hz), 7.23 (2H, d, J=8Hz), 7.30 (1H, t,
J=8Hz), 7.66 (1H, d, J=8Hz), 7.72 (1H, d, J=8Hz),
7.89 (1H, br s), 9.30 (1H, br peak)

- 16) 2-(N,N-Dimethylcarbamoyl)-1-(4-methoxybenzyl)-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 2.91 (3H, s), 3.05 (3H, s), 3.71
(3H, s), 5.49 (2H, s), 6.89 (2H, d, J=8Hz), 7.22
(2H, d, J=8Hz), 7.42 (1H, t, J=8Hz), 7.82 (1H, d,
J=8Hz), 7.92 (1H, d, J=8Hz)

- 17) 2-[1-(Benzyloxycarbonyl)-4-piperidyl]-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 1.69-1.88 (2H, m), 1.95-2.08 (2H,
m), 2.99 (2H, br peak), 3.24-3.59 (1H, m), 4.08-
4.19 (2H, m), 5.10 (2H, s), 7.23-7.47 (6H, m), 7.79
(1H, d, J=8Hz), 7.85 (1H, d, J=8Hz)

- 18) 2-(N-tert-Butoxycarbonylaminomethyl)-1-methyl-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 1.36 (9H, s), 4.01 (3H, s), 4.77
(2H, d, J=5Hz), 7.16-7.31 (2H, m), 8.11 (1H, d,
J=8Hz), 8.23 (1H, d, J=8Hz)

- 19) 2-(N-tert-Butoxycarbonylaminomethyl)-3-methyl-3H-

benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 1.40 (9H, s), 3.96 (3H, s), 4.65 (2H, d, J=6Hz), 7.52 (1H, t, J=8Hz), 7.77 (1H, br peak), 7.87 (1H, d, J=8Hz), 7.95 (1H, d, J=8Hz)

5

20) 2-Methylthio-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 2.68 (3H, s), 7.22 (1H, t, J=8Hz), 7.70 (1H, d, J=8Hz), 7.76 (1H, d, J=8Hz)

10 21) 2-Methylsulfonyl-1H-benzimidazole-4-carboxylic acid

NMR (DMSO-d₆, δ) : 3.56 (3H, s), 7.50 (1H, t, J=8Hz), 8.01 (1H, d, J=8Hz), 8.09 (1H, br peak)

22) 2-Sulfamoyl-1H-benzimidazole-4-carboxylic acid

15 NMR (DMSO-d₆, δ) : 7.46 (1H, t, J=8Hz), 7.89-8.02 (3H, m), 8.07 (1H, d, J=8Hz)

23) 2-Methyl-1H-pyrazolo[1,5-b][1,2,4]triazole-7-carboxylic acid

20 NMR (DMSO-d₆, δ) : 2.43 (3H, s), 7.82 (1H, s)

24) 2-(4-Pyridyl)-1H-benzimidazole-4-carboxylic acid

25 NMR (DMSO-d₆, δ) : 7.37 (1H, t, J=8Hz), 7.89 (1H, d, J=8Hz), 8.00 (1H, d, J=8Hz), 8.27 (1H, d, J=6Hz), 8.74 (1H, d, J=6Hz)

25) 2-(3-Pyridyl)-1H-benzimidazole-4-carboxylic acid

30 NMR (DMSO-d₆, δ) : 7.40 (1H, t, J=8Hz), 7.72 (1H, t, J=8Hz), 7.90 (1H, d, J=8Hz), 8.01 (1H, d, J=8Hz), 8.77-8.80 (2H, m), 9.52 (1H, s)

26) 2-(2-Pyridyl)-1H-benzimidazole-4-carboxylic acid

35 NMR (DMSO-d₆, δ) : 7.39 (1H, t, J=8Hz), 7.57 (1H, t, J=7Hz), 7.87 (1H, d, J=8Hz), 7.98-8.06 (2H, m), 8.35 (1H, d, J=8Hz), 8.79 (1H, d, J=4Hz)

- 27) 2-Dimethylaminomethyl-1H-benzimidazole-4-carboxylic acid
NMR (DMSO-d₆, δ) : 2.89 (6H, s), 4.68 (2H, s), 7.39
(1H, t, J=8Hz), 7.90 (1H, d, J=8Hz), 8.00 (1H, d,
J=8Hz)
- 5
- 28) 2-(4-Methylpiperazin-1-yl)methyl-1H-benzimidazole-4-
carboxylic acid
NMR (DMSO-d₆, δ) : 2.50 (3H, s), 2.72 (4H, br s), 2.88
(4H, br s), 3.87 (2H, s), 7.27 (1H, t, J=8Hz), 7.77
10 (1H, d, J=8Hz), 7.82 (1H, d, J=8Hz)
- 29) 2-(4-Dimethylaminopiperidino)methyl-1H-benzimidazole-4-
carboxylic acid
NMR (DMSO-d₆, δ) : 1.90 (2H, br s), 2.10-2.20 (2H, m),
15 2.52-2.70 (7H, m), 3.15-3.50 (6H, m), 7.30 (1H, t,
J=8Hz), 7.82 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz)
- 30) 2-Morpholinomethyl-1H-benzimidazole-4-carboxylic acid
NMR (DMSO-d₆, δ) : 2.50 (4H, br s), 3.88 (4H, br s),
20 4.72 (2H, s), 7.40 (1H, t, J=8Hz), 7.90 (1H, d,
J=8Hz), 7.99 (1H, d, J=8Hz)
- 31) 2H-1,4-Benzoxazin-3-one-8-carboxylic acid
NMR (DMSO-d₆, δ) : 4.62 (2H, s), 6.96-7.05 (2H, m),
25 7.30 (1H, d, J=8Hz)

Preparation 45

The following compounds were obtained according to a
similar manner to that of Preparation 25.

30

- 1) 4-Amino-3-methoxy-N-methyl-N-[2-(5-tert-butoxycarbonyl-
aminopent-1-yl)oxy-4-methyl]phenylbenzamide
NMR (CDCl₃, δ) : 1.42 (9H, s), 1.42-1.59 (4H, m), 1.70-
1.81 (2H, m), 2.28 (9H, s), 3.06-3.17 (2H, m), 3.31
35 (3H, s), 3.61 (3H, s), 3.76-3.93 (2H, m), 3.87 (2H,

s), 4.69 (1H, br), 6.43 (1H, d, J=8Hz), 6.57-6.63 (2H, m), 6.80-6.85 (2H, m), 6.89 (1H, d, J=8Hz)

2) tert-Butyl 4-amino-1H-benzimidazole-1-carboxylate

5 NMR (CDCl₃, δ) : 1.68 (9H, s), 4.37 (2H, s), 6.61 (1H, d, J=8Hz), 7.16 (1H, dd, J=8, 8Hz), 7.32 (1H, d, J=8Hz), 8.30 (1H, s)

3) 2-[4,4-Dimethyl(2,5-oxazolinyl)]phenylamine

10 NMR (CDCl₃, δ) : 1.39 (6H, s), 4.00 (2H, s), 6.06-6.28 (2H, br), 6.60-6.73 (2H, m), 7.19 (1H, m), 7.68 (1H, d, J=8Hz)

4) 4-Amino-N-[2-[4,4-dimethyl(2,5-oxazolinyl)]phenyl]-N-methyl-3-methoxybenzamide

15 NMR (CDCl₃, δ) : 1.32 (6H, s), 3.38 (3H, s), 3.58 (3H, s), 3.86 (2H, br s), 4.07 (2H, s), 6.41 (1H, d, J=8Hz), 6.78-6.90 (2H, m), 7.12 (1H, d, J=8Hz), 7.22 (1H, dd, J=8, 8Hz), 7.35 (1H, dd, J=8, 8Hz),
20 7.79 (1H, d, J=8Hz)

5) 4-Amino-3-methoxy-N-methyl-N-[2-(morpholin-4-yl)]phenyl]benzamide

25 NMR (CDCl₃, δ) : 2.52-2.68 (2H, m), 2.77-2.93 (2H, m), 3.46 (3H, s), 3.59 (3H, s), 3.63-3.82 (4H, m), 6.45 (1H, d, J=8Hz), 6.84 (1H, s), 6.86-6.96 (2H, m), 7.04 (1H, dd, J=8, 8Hz), 7.12-7.23 (2H, m)

6) 2-(4-Methyl-1-piperazinyl)phenylamine

30 NMR (CDCl₃, δ) : 2.37 (3H, s), 2.46-2.75 (4H, m), 2.88-3.04 (4H, m), 3.94 (2H, br s), 6.75 (2H, dd, J=8, 8Hz), 6.92 (1H, dd, J=8, 8Hz), 7.02 (1H, d, J=8Hz)

7) 2-(2,5-Oxazolyl)phenylamine

35 NMR (CDCl₃, δ) : 6.70-6.83 (2H, m), 7.16-7.28 (2H, m),

7.64 (1H, s), 7.86 (1H, m)

8) 2-(1-Aza-3-oxaspiro[4.4]non-1-en-2-yl)phenylamine

5 NMR (CDCl₃, δ) : 1.52-2.06 (8H, m), 4.12 (2H, s), 6.10
(2H, br s), 6.59-6.73 (2H, m), 7.18 (1H, dd, J=8,
8Hz), 7.66 (1H, d, J=8Hz)

9) 4-Amino-2-methoxy-1H-benzimidazole

10 NMR (DMSO-d₆, δ) : 4.01 (3H, s), 4.84 (2H, s), 6.30
(1H, d, J=8Hz), 6.47 (1H, br peak), 6.74 (1H, t,
J=8Hz)

10) 4-Amino-2-[2-(dimethylamino)ethyl]-1H-benzimidazole

15 NMR (CDCl₃, δ) : 2.39 (6H, s), 2.71 (2H, t, J=5Hz),
3.07 (2H, t, J=5Hz), 4.23 (2H, br peak), 6.50 (1H,
d, J=8Hz), 6.84 (1H, d, J=8Hz), 7.00 (1H, t, J=8Hz)

11) 4-Amino-1-(tert-butoxycarbonyl)-2-[[2-[N-(tert-butoxy-
carbonyl)-N-methylamino]ethyl]amino]-1H-benzimidazole

20 NMR (CDCl₃, δ) : 1.45 (9H, s), 1.69 (9H, s), 2.93 (3H,
s), 3.51-3.65 (2H, m), 3.65-3.88 (2H, m), 6.57 (1H,
d, J=8Hz), 6.88 (1H, t, J=8Hz), 7.02 (1H, d, J=8Hz)

12) 4-Amino-1-(tert-butoxycarbonyl)-2-[[2-[(tert-butoxy)-
carbonylamino]ethyl]methylamino]-1H-benzimidazole

25 NMR (CDCl₃, δ) : 1.35 (9H, s), 1.70 (9H, s), 3.03 (3H,
s), 3.41-3.55 (2H, m), 3.61-3.77 (2H, m), 6.14 (1H,
br peak), 6.59 (1H, d, J=8Hz), 6.91-7.10 (2H, m)

13) 4-Amino-1-(tert-butoxycarbonyl)-2-[[2-(dimethylamino)-
ethyl]amino]-1H-benzimidazole

30 NMR (CDCl₃, δ) : 1.69 (9H, s), 2.43 (6H, br s), 2.80
(2H, br peak), 2.69-2.81 (2H, m), 6.56 (1H, d,
J=8Hz), 6.85 (1H, t, J=8Hz), 7.06 (1H, d, J=8Hz)

- 14) 4-Amino-2-[[2-(dimethylamino)ethyl]methylamino]-1H-benzimidazole

NMR (DMSO-d₆, δ) : 2.20 (6H, s), 2.45 (2H, t, J=5Hz),
3.11 (3H, s), 3.52 (2H, t, J=5Hz), 6.18 (1H, d,
J=8Hz), 6.44 (1H, d, J=8Hz), 6.60 (1H, t, J=8Hz)

Preparation 46

The following compounds were obtained according to a similar manner to that of Preparation 31.

- 1) Methyl-2-carbamoyl-1-(4-methoxybenzyl)-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 3.74 (3H, s), 4.04 (3H, s), 5.65 (1H, br s), 5.98 (2H, s), 6.80 (2H, d, J=8Hz), 7.15 (1H, d, J=8Hz), 7.40 (1H, t, J=8Hz), 7.63 (1H, d, J=8Hz), 7.86 (1H, br s), 8.03 (1H, d, J=8Hz)

- 2) Methyl 2-(N,N-dimethylcarbamoyl)-1-(4-methoxybenzyl)-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 3.10 (3H, s), 3.19 (3H, s), 3.76 (3H, s), 4.01 (3H, s), 5.54 (2H, s), 6.80 (2H, d, J=8Hz), 7.12 (2H, d, J=8Hz), 7.35 (1H, t, J=8Hz), 7.58 (1H, d, J=8Hz), 7.99 (1H, d, J=8Hz)

Preparation 47

The following compounds were obtained by using ethyl 2-phthalimidomethyl-1H-benzimidazole-4-carboxylate as a starting compound according to a similar manner to that of Preparation 31.

- 1) Ethyl 1-methyl-2-phthalimidomethyl-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.26 (3H, t, J=7.5Hz), 3.95 (3H, s), 4.35 (2H, q, J=7.5Hz), 5.20 (2H, s), 7.32 (1H, t, J=8Hz), 7.52 (1H, d, J=8Hz), 7.70-7.80 (2H, m),

7.84-7.95 (3H, m)

2) Ethyl 3-methyl-2-phthalimidomethyl-3H-benzimidazole-4-carboxylate

5 NMR (CDCl₃, δ) : 1.45 (3H, t, J=7.5Hz), 4.06 (3H, s),
4.43 (2H, q, J=7.5Hz), 5.15 (2H, s), 7.21 (1H, t,
J=8Hz), 7.71-7.80 (3H, m), 7.85 (1H, d, J=8Hz),
7.87-7.94 (2H, m)

10 Preparation 48

To a solution of benzyl 2-tert-butyldiphenylsilyloxymethylindole-4-carboxylate (805 mg) and N,N-dimethylaminopyridine (189 mg) in acetonitrile (15 ml) was added di-tert-butyl dicarbonate (507 mg) and the mixture
15 was stirred at ambient temperature for 20 minutes. The resulting mixture was diluted with ethyl acetate (30 ml) and washed successively with saturated sodium bicarbonate aqueous solution and brine. The organic layer was dried over magnesium sulfate and concentrated in vacuo. The residue was
20 purified by silica gel column chromatography (eluent; n-hexane:ethyl acetate = 15:1) to give benzyl 1-tert-butoxycarbonyl-2-tert-butyldiphenylsilyloxymethylindole-4-carboxylate (762 mg).

25 NMR (CDCl₃, δ) : 1.12 (9H, s), 1.47 (9H, s), 5.01 (2H, s), 5.47 (2H, s), 7.27-7.50 (12H, m), 7.55 (1H, s), 7.71 (4H, d, J=8Hz), 8.01 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)

Preparation 49

30 The following compounds were obtained according to a similar manner to that of Preparation 48.

1) Methyl 2-benzyloxymethyl-1-tert-butoxycarbonylindoline-4-carboxylate

35 NMR (CDCl₃, δ) : 1.50 (9H, s), 3.42-3.53 (3H, m), 3.67

(1H, dd, J=4, 12Hz), 3.89 (3H, s), 4.50 (2H, s),
4.56-4.67 (1H, br), 7.20-7.38 (7H, m), 7.60 (1H, d,
J=8Hz)

- 5 2) Benzyl 1-tert-butoxycarbonyl-3-formylindole-4-
 carboxylate

NMR (CDCl₃, δ) : 1.68 (9H, s), 5.43 (2H, s), 7.32-7.49
(6H, m), 7.92 (1H, d, J=8Hz), 8.37 (1H, s), 8.49
(1H, d, J=8Hz), 10.47 (1H, s)

10

- 3) Benzyl 1-tert-butoxycarbonyl-2-formylindole-4-
carboxylate

NMR (CDCl₃, δ) : 1.72 (9H, s), 5.45 (2H, s), 7.34-7.45
(3H, m), 7.47-7.57 (3H, m), 8.07 (1H, d, J=9Hz),
8.10 (1H, s), 8.43 (1H, d, J=9Hz), 10.40 (1H, s)

15

- 4) Benzyl 1-tert-butoxycarbonyl-2-methylindole-4-
carboxylate

NMR (CDCl₃, δ) : 1.70 (9H, s), 2.61 (3H, s), 5.40 (2H,
s), 7.07 (1H, s), 7.22-7.28 (1H, m), 7.31-7.43 (3H,
m), 7.47-7.52 (2H, m), 7.97 (1H, d, J=8Hz), 8.35
(1H, d, J=8Hz)

20

- 5) Benzyl 1-tert-butoxycarbonylindole-6-carboxylate

NMR (CDCl₃, δ) : 1.67 (9H, s), 5.39 (2H, s), 6.61 (1H,
d, J=3Hz), 7.32-7.42 (3H, m), 7.46-7.50 (2H, m),
7.59 (1H, d, J=8Hz), 7.78 (1H, d, J=3Hz), 7.97 (1H,
d, J=8Hz) 8.86 (1H, s)

25

- 6) 3-Methoxy-4-methoxycarbonyl-N-methyl-N-[4-methyl-2-[4-
(tert-butoxycarbonylamino)but-1-yloxy]]phenylbenzamide

NMR (CDCl₃, δ) : 1.42 (9H, s), 1.57-1.70 (2H, m), 1.72-
1.86 (2H, m), 2.23 (3H, s), 3.19 (2H, t, J=7.5Hz),
3.31 (3H, s), 3.69 (3H, s), 3.78-3.96 (2H, m), 3.82
(3H, s), 6.55-6.60 (2H, m), 6.87 (1H, d, J=8Hz),

35

6.91 (1H, d, J=8Hz), 6.94 (1H, s), 7.57 (1H, d, J=8Hz)

7) tert-Butyl 4-nitro-1H-benzimidazole-1-carboxylate

5 NMR (CDCl₃, δ) : 1.72 (9H, s), 7.53 (1H, dd, J=8, 8Hz),
8.20 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz), 8.61 (1H, s)

8) 1-(tert-Butoxycarbonyl)-2-[[2-[N-(tert-butoxycarbonyl)-
10 N-methylamino]ethyl]amino]-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 1.41 (9H, s), 1.71 (9H, s), 2.95 (3H, s),
3.60 (1H, t-like, J=5Hz), 3.84 (1H, q-like, J=5Hz), 7.04 (1H, t, J=8Hz), 7.31 (1H, d, J=8Hz),
8.00 (1H, d, J=8Hz)

9) 1-(tert-Butoxycarbonyl)-2-[[2-[(tert-butoxy)carbonyl-
amino]ethyl]methylamino]-4-nitro-1H-benzimidazole

15 NMR (CDCl₃, δ) : 1.39 (9H, s), 1.69 (9H, s), 3.11 (3H, s),
3.50 (2H, q-like, J=5Hz), 3.80 (2H, t-like, J=5Hz),
20 6.01 (1H, br peak), 7.11 (1H, t, J=8Hz),
7.88 (1H, d, J=8Hz), 8.03 (1H, d, J=8Hz)

10) 1-(tert-Butoxycarbonyl)-2-[[2-(dimethylamino)ethyl]-
amino]-4-nitro-1H-benzimidazole

25 NMR (CDCl₃, δ) : 1.72 (9H, s), 2.55 (6H, br peak), 3.00 (2H, br peak),
3.92 (2H, br peak), 7.07 (1H, t, J=8Hz), 7.84 (1H, br peak), 7.90 (1H, d, J=8Hz),
8.01 (1H, d, J=8Hz)

30 Preparation 50

The following compounds were obtained according to a similar manner to that of Example 1.

1) 3-Methoxy-4-methoxycarbonyl-N-methyl-N-(4-methyl-2-
35 nitrophenyl)benzamide

NMR (CDCl₃, δ) : 2.35 (3H, s), 3.37 (3H, s), 3.70 (3H, s), 3.80 (3H, s), 6.80 (1H, d, J=8Hz), 6.91 (1H, s), 7.11 (1H, d, J=8Hz), 7.30 (1H, d, J=8Hz), 7.54 (1H, d, J=8Hz), 7.63 (1H, s)

5

- 2) 3-Methoxy-4-methoxycarbonyl-N-[4-methyl-2-(4-phthalimidobut-1-yloxy)]phenylbenzamide

NMR (CDCl₃, δ) : 1.87-1.96 (4H, m), 2.31 (3H, s), 3.78 (2H, m), 3.91 (3H, s), 3.98 (3H, s), 4.10 (2H, m), 6.70 (1H, s), 6.79 (1H, d, J=8Hz), 7.35 (1H, d, J=8Hz), 7.62 (1H, s), 7.66-7.74 (2H, m), 7.79-7.87 (2H, m), 7.96 (1H, d, J=8Hz), 8.33 (1H, d, J=8Hz), 8.52 (1H, s)

10

- 15 3) 3-Methoxy-4-nitro-N-[4-methyl-2-(5-phthalimidopent-1-yloxy)]phenylbenzamide

NMR (CDCl₃, δ) : 1.48-1.60 (2H, m), 1.70-1.82 (2H, m), 1.85-1.95 (2H, m), 2.31 (3H, s), 3.69 (2H, t, J=7.5Hz), 4.02 (3H, s), 4.04 (2H, t, J=7.5Hz), 6.71 (1H, s), 6.81 (1H, d, J=8Hz), 7.39 (1H, d, J=8Hz), 7.66-7.72 (2H, m), 7.76-7.74 (2H, m), 7.98 (1H, d, J=8Hz), 8.32 (1H, d, J=8Hz), 8.52 (1H, s)

20

Preparation 51

25 The following compounds were obtained according to a similar manner to that of Example 7.

- 1) 4-Amino-2-phthalimidomethyl-1H-benzimidazole

NMR (DMSO-d₆, δ) : 4.95 (2H, s), 5.10 (2H, br s), 6.30 (1H, d, J=8Hz), 6.57 (1H, d, J=8Hz), 6.82 (1H, t, J=8Hz), 7.83-8.02 (4H, m)

30

- 2) 4-Amino-2-(2-phthalimidoethyl)-1H-benzimidazole

NMR (DMSO-d₆, δ) : 3.11 (2H, br peak), 4.00 (2H, br peak), 5.01 (2H, br peak), 6.28 (1H, br peak), 6.60

35

(1H, br peak), 6.80 (1H, br peak), 7.85 (4H, br peak)

- 5 3) 4-Amino-2-tert-butyldiphenylsiloxymethyl-1H-benzimidazole

NMR (CD₃OD, δ) : 1.09 (9H, s), 4.91 (2H, s), 6.53 (1H, d, J=8Hz), 6.88 (1H, d, J=8Hz), 6.99 (1H, t, J=8Hz), 7.32-7.50 (6H, m), 7.66-7.75 (4H, m)

- 10 4) 4-Amino-2-[(tert-butoxy)carbonylamino]-1H-benzimidazole
NMR (CDCl₃, δ) : 1.69 (9H, s), 6.58 (1H, d, J=8Hz), 6.90 (1H, t, J=8Hz), 7.06 (1H, d, J=8Hz)

- 15 5) 4-Amino-2-[(methylsulfonyl)amino]-1H-benzimidazole
NMR (DMSO-d₆, δ) : 3.40 (3H, s), 4.90 (2H, s), 6.41-6.51 (1H, m), 6.65 (2H, s), 6.71-6.81 (2H, m)

- 20 6) 4-Amino-2-methoxymethyl-1H-benzimidazole
NMR (DMSO-d₆, δ) : 3.32 (3H x 2/3, s), 3.37 (3H x 1/3, s), 4.55 (2H x 2/3, s), 4.60 (2H x 1/3, s), 5.65 (2H, s), 6.26-6.40 (1H, m), 6.62 (1H x 2/3, d, J=8Hz), 6.73-6.90 (1H + 1H x 1/3, m)

- 25 7) 4-Amino-2-(n-propyl)-1H-benzimidazole
NMR (CDCl₃, δ) : 1.00 (3H, t, J=7.5Hz), 1.83 (2H, m), 2.86 (2H, t, J=7.5Hz), 4.26 (2H, br s), 6.50 (1H, d, J=8Hz), 6.80 (1H, d, J=8Hz), 7.00 (1H, t, J=8Hz)

- 30 8) 4-Amino-2-isopropyl-1H-benzimidazole
NMR (CDCl₃, δ) : 1.43 (6H, d, J=7.5Hz), 3.23 (1H, m), 4.28 (1H, br s), 6.50 (1H, d, J=8Hz), 6.80 (1H, br peak), 7.01 (1H, t, J=8Hz), 8.84 (1H, br s)

- 35 9) 4-Amino-2-(3-pyridyl)-1H-benzimidazole
NMR (DMSO-d₆, δ) : 5.30 (2H, br s), 6.38 (1H, d,

141

J=8Hz), 6.73 (1H, d, J=8Hz), 6.93 (1H, t, J=8Hz), 7.56 (1H, dd, J=5, 8Hz), 8.43 (1H, d, J=8Hz), 8.93 (1H, d, J=5Hz), 9.30 (1H, s)

- 5 10) 4-Amino-2-(N,N-dimethylaminomethyl)-1H-benzimidazole

NMR (CDCl₃, δ) : 2.41 (6H, s), 3.83 (2H, s), 6.50 (1H, d, J=8Hz), 6.83 (1H, d, J=8Hz), 7.04 (1H, t, J=8Hz)

- 10 11) 4-Amino-2-(1-imidazolyl)methyl-1H-benzimidazole

NMR (DMSO-d₆, δ) : 5.16 (2H, s), 5.38 (2H, s), 6.33 (1H, d, J=8Hz), 6.67 (1H, d, J=8Hz), 6.86 (1H, t, J=8Hz), 6.91 (1H, s), 7.23 (1H, s), 7.77 (1H, s)

- 15 12) 4-Amino-2-[(4-methylpiperazin-1-yl)methyl]-1H-
benzimidazole

NMR (DMSO- d_6 , δ) : 2.15 (3H, s), 2.20-2.49 (8H, m),
3.61 (2H x 2/3, s), 3.65 (2H x 1/3, s), 5.10 (2H x
2/3, s), 5.20 (2H x 1/3, s), 6.28 (1H x 2/3, d,
J=8Hz), 6.33 (1H x 1/3, d, J=8Hz), 6.60 (1H x 2/3,
d, J=8Hz), 6.73-6.86 (1H + 1H x 1/3, m)

- 13) 4-Amino-2-(morpholin-4-ylmethyl)-1H-benzimidazole

NMR (CDCl₃, δ) : 2.50-2.60 (4H, m), 3.68-3.76 (4H, m), 3.79 (2H, s), 4.28 (2H, br peak), 6.51 (1H, d, J=8Hz), 6.83 (1H, d, J=8Hz), 7.03 (1H, t, J=8Hz)

- 14) 4-Amino-2-(pyrrolidin-1-ylmethyl)-1H-benzimidazole

NMR (CDCl₃, δ) : 1.83-2.00 (2H, m), 2.71-2.83 (2H, m), 4.03 (2H, s), 6.50 (1H, d, J=8Hz), 6.85 (1H, d, J=8Hz), 7.03 (1H, t, J=8Hz)

- 15) 4-Amino-2-(piperidinomethyl)-1H-benzimidazole

NMR (CDCl₃, δ) : 1.47-1.63 (2H, m), 1.73-1.86 (4H, m),
2.73 (4H, br peak), 4.01 (2H, s), 4.26 (2H, br

142

peak), 6.51 (1H, d, J=8Hz), 6.90 (1H, d, J=8Hz),
7.05 (1H, t, J=8Hz)

- 16) 4-Amino-2-[2-(4-methylpiperazin-1-yl)ethyl]-1H-benzimidazole

NMR (DMSO-d₆, δ) : 2.18 (3H, s), 2.22-2.53 (8H, m),
2.72 (2H, t, J=7.5Hz), 2.91 (2H, t, J=7.5Hz), 5.02
(2H, br peak), 6.29 (1H, d, J=8Hz), 6.61 (1H, d,
J=8Hz), 6.80 (1H, t, J=8Hz)

- 17) 4-Amino-2-(4-methylpiperazin-1-yl)-1H-benzimidazole

NMR (DMSO-d₆, δ) : 2.22 (3H, s), 2.36-2.46 (4H, m),
3.36-3.49 (4H, m), 4.70 (2H x 2/3, br s), 4.82 (2H
x 1/3, br s), 6.20 (1H x 1/3, d, J=8Hz), 6.23 (1H x
2/3, d, J=8Hz), 6.45 (1H x 2/3, d, J=8Hz), 6.52 (1H
x 1/3, d, J=8Hz), 6.62 (1H x 2/3, t, J=8Hz), 6.68
(1H x 1/3, t, J=8Hz)

- 18) 4-Amino-2-dimethylamino-1H-benzimidazole

NMR (DMSO-d₆, δ) : 3.21 (6H, s), 5.73 (2H, br peak),
6.46 (1H, d, J=8Hz), 6.57 (1H, d, J=8Hz), 6.93 (1H,
t, J=8Hz)

- 19) 4-Amino-2-(1-imidazolyl)-1H-benzimidazole

NMR (DMSO-d₆, δ) : 5.20 (2H, br s), 6.42 (1H, d,
J=8Hz), 6.72 (1H, br peak), 6.94 (1H, t, J=8Hz),
7.20 (1H, s), 7.85 (1H, s), 8.40 (1H, s)

- 20) 4-Amino-2-(1,2,4-tetrazol-1-yl)-1H-benzimidazole

NMR (DMSO-d₆, δ) : 5.31 (2H, br peak), 6.43 (1H, d,
J=8Hz), 6.85 (1H, br peak), 6.95 (1H, t, J=8Hz),
8.42 (1H, s), 9.34 (1H, s)

- 21) 4-Amino-2-[(2-methoxyethyl)amino]-1H-benzimidazole

NMR (DMSO-d₆, δ) : 3.30 (3H, s), 3.45-3.60 (4H, m),

143

6.32 (1H, d, J=8Hz), 6.50 (1H, d, J=8Hz), 6.75 (1H, t, J=8Hz), 7.55 (1H, br peak)

Preparation 52

5 The following compound was obtained according to a similar manner to that of Example 14.

Methyl 2-formyl-1H-benzimidazole-4-carboxylate

10 NMR (DMSO-d₆, δ) : 3.97 (3H, s), 7.50 (1H, t, J=8Hz),
8.03 (1H, d, J=8Hz), 8.11 (1H, d, J=8Hz), 10.06
(1H, s)

Preparation 53

15 The following compound was obtained according to a similar manner to that of Example 16.

2-Methoxymethyl-4-nitro-1H-benzimidazole

20 NMR (CDCl₃, δ) : 3.59 (3H, s), 4.85 (2H, s), 7.40 (1H, t, J=8Hz), 8.08 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz)

Preparation 54

To a solution of ethyl 2,3-diaminobenzoate (4.72 g) and pyridine (2.49 g) in dichloroethane (50 ml) was added chloroacetyl chloride (3.11 g) in chloroform (10 ml) at -70°C and the reaction mixture was stood overnight. After the reaction mixture was concentrated in vacuo, the residue was diluted with ethanol (50 ml). To the solution was added p-toluenesulfonic acid (249 mg) and the reaction mixture was refluxed for 2 hours. After the reaction mixture was concentrated in vacuo, the residue was diluted with ethyl acetate and saturated sodium hydrogen carbonate aqueous solution. The organic layer was separated and washed with brine. The solution was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by silica gel column chromatography (n-hexane:ethyl acetate = 4:1) to

25

30

35

give ethyl 2-chloromethyl-1H-benzimidazole-4-carboxylate .
(2.98 g) .

5 NMR (CDCl₃, δ) : 0.94 (3H, t, J=7Hz), 4.47 (2H, q,
J=7Hz), 4.86 (2H, s), 7.32 (1H, t, J=8Hz), 7.93
(2H, d, J=8Hz)

Preparation 55

The following compounds were obtained according to a
similar manner to that of Preparation 54.

10

1) 4-Nitro-2-(n-propyl)-1H-benzimidazole

NMR (CDCl₃, δ) : 1.08 (3H, t, J=7.5Hz), 1.94 (2H, m),
2.99 (2H, t, J=7.5Hz), 7.34 (1H, t, J=8Hz), 8.03
(1H, d, J=8Hz), 8.13 (1H, d, J=8Hz)

15

2) 2-Isopropyl-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 1.51 (6H, d, J=7.5Hz), 3.32 (1H, m),
7.34 (1H, t, J=8Hz), 8.06 (1H, d, J=8Hz), 8.12 (1H,
d, J=8Hz)

20

3) 4-Nitro-2-(3-pyridyl)-1H-benzimidazole

NMR (DMSO-d₆, δ) : 7.47 (1H, t, J=8Hz), 7.63 (1H, dd,
J=5, 8Hz), 8.17 (2H, d, J=8Hz), 8.69 (1H, dd, J=8,
2Hz), 8.74 (1H, dd, J=5, 2Hz), 9.47 (1H, d, J=2Hz)

25

4) 2-(2-Chloroethyl)-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 3.49 (2H, t, J=7Hz), 4.04 (2H, t,
J=7Hz), 7.39 (1H, t, J=8Hz), 8.06 (1H, d, J=8Hz),
8.17 (1H, d, J=8Hz)

30

5) Ethyl 2-(3-pyridyl)-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.47 (3H, t, J=7Hz), 4.48 (3H, q,
J=7Hz), 7.36 (1H, t, J=8Hz), 7.46-7.50 (1H, m),
7.93 (1H, d, J=8Hz), 8.03 (1H, d, J=8Hz), 8.42 (1H,
35 d, J=8Hz), 8.73 (1H, d, J=3Hz), 9.32 (1H, s)

145

6) Ethyl 2-(2-pyridyl)-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.48 (3H, t, J=7Hz), 4.50 (2H, q, J=7Hz), 7.31-7.40 (2H, m), 7.86 (1H, t, J=8Hz), 7.97 (1H, d, J=8Hz), 8.03 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz), 8.68 (1H, d, J=3Hz)

Preparation 56

The following compounds were obtained according to a similar manner to that of Example 26.

1) 3-Methoxy-4-methoxycarbonyl-N-methyl-N-[4-methyl-2-(4-aminobut-1-yloxy)]phenylbenzamide

2) 3-Methoxy-N-methyl-4-nitro-N-[2-(5-tert-butoxycarbonyl-aminopent-1-yl)oxy-4-methyl]phenylbenzamide

NMR (CDCl₃, δ) : 1.44 (9H, s), 1.47-1.68 (4H, m), 1.73-1.87 (2H, m), 2.29 (3H, s), 3.10-3.18 (2H, m), 3.31 (3H, s), 3.79 (3H, s), 3.84-3.95 (2H, m), 6.58-6.62 (2H, m), 6.88 (1H, d, J=8Hz), 6.95 (1H, d, J=8Hz), 7.09 (1H, s), 7.62 (1H, d, J=8Hz)

3) Ethyl 2-aminomethyl-1-methyl-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.45 (3H, t, J=7.5Hz), 3.84 (3H, s), 4.20 (2H, s), 4.49 (2H, q, J=7.5Hz), 7.31 (1H, t, J=8Hz), 7.51 (1H, d, J=8Hz), 7.93 (1H, d, J=8Hz)

4) Ethyl 2-aminomethyl-3-methyl-3H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.43 (3H, t, J=7.5Hz), 3.93 (3H, s), 4.15 (2H, s), 4.43 (2H, q, J=7.5Hz), 7.25 (1H, t, J=8Hz), 7.76 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz)

Preparation 57

The following compounds were obtained according to a

similar manner to that of Example 30.

1) Methyl 2-carbamoyl-1H-benzimidazole-4-carboxylate

5 NMR (DMSO-d₆, δ) : 3.94 (3H, s), 7.36 (1H, t, J=8Hz),
7.82-7.92 (2H, m), 7.97 (1H, d, J=8Hz), 8.26 (1H,
br s)

2) Methyl 2-(N,N-dimethylcarbamoyl)-1H-benzimidazole-4-carboxylate

10 NMR (CDCl₃, δ) : 3.21 (3H, s), 3.80 (3H, s), 4.02 (3H,
s), 7.38 (1H, t, J=8Hz), 8.04 (2H, d, J=8Hz)

3) 2-(N,N-Dimethylcarbamoyl)-4-nitro-1H-benzimidazole

15 NMR (DMSO-d₆, δ) : 3.10 (3H, s), 3.33 (3H, s), 7.51
(3H, t, J=8Hz), 8.09-8.25 (2H, m)

Preparation 58

To a suspension of 3-formylindole-4-carboxylic acid (390 mg) and potassium carbonate (285 mg) in N,N-dimethylformamide
20 (10 ml) was added benzyl bromide (353 mg) at ambient temperature and the mixture was stirred for 4 hours. The resulting mixture was diluted with ethyl acetate and water, then the layers were separated. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in
25 vacuo. The residual solid was triturated with diethyl ether:n-hexane (1:6) to give benzyl 3-formylindole-4-carboxylate (520 mg).

30 NMR (DMSO-d₆, δ) : 5.37 (2H, s), 7.28-7.48 (7H, m),
7.62 (1H, d, J=8Hz), 7.75 (1H, d, J=8Hz), 8.36 (1H, s), 10.22 (1H, s)

Preparation 59

The following compounds were obtained according to a similar manner to that of Preparation 58.

147

1) Benzyl 2-hydroxymethylindole-4-carboxylate

NMR (CDCl₃, δ) : 4.88 (1H, d, J=6Hz), 5.44 (2H, s),
7.03 (1H, s), 7.22 (1H, t, J=8Hz), 7.32-7.43 (3H,
m), 7.47-7.52 (2H, m), 7.56 (1H, d, J=8Hz), 7.94
(1H, d, J=8Hz), 8.52-8.58 (1H, br)

2) Benzyl 2-methylindole-4-carboxylate

NMR (CDCl₃, δ) : 2.48 (3H, s), 5.44 (2H, s), 6.88 (1H,
s), 7.13 (1H, t, J=8Hz), 7.30-7.52 (6H, m), 7.90
(1H, d, J=8Hz), 8.03-8.10 (1H, br)

Preparation 60

To a solution of benzyl 1-tert-butoxycarbonyl-3-formylindole-4-carboxylate (363 mg) in methanol (15 ml) was added sodium borohydride (109 mg) at 0°C and the mixture was stirred for 5 minutes. The resulting mixture was diluted with water and the solution was neutralized with 1N hydrochloric acid. The solution was extracted with ethyl acetate, and then the organic solution was washed successively with saturated sodium bicarbonate aqueous solution and brine, dried over magnesium sulfate and concentrated in vacuo to afford benzyl 1-tert-butoxycarbonyl-3-hydroxymethylindole-4-carboxylate (365 mg).

NMR (CDCl₃, δ) : 1.67 (9H, s), 4.11 (1H, t, J=8Hz),
4.72 (2H, d, J=8Hz), 5.43 (2H, s), 7.30-7.51 (6H, m), 7.71 (1H, s), 7.92 (1H, d, J=8Hz), 8.49 (1H, d, J=8Hz)

Preparation 61

The following compound was obtained according to a similar manner to that of Preparation 60.

Benzyl 1-tert-butoxycarbonyl-2-hydroxymethylindole-4-carboxylate

NMR (CDCl₃, δ) : 1.73 (9H, s), 3.58 (1H, t, J=9Hz),

148

4.82 (2H, d, J=9Hz), 5.42 (2H, s), 7.30-7.42 (5H, m), 7.47-7.50 (2H, m), 8.00 (1H, d, J=8Hz), 8.22 (1H, d, J=8Hz)

5 Preparation 62

To a solution of 3-methoxy-4-methoxycarbonylbenzoic acid (800 mg) in dichloromethane (15 ml) was added oxalyl chloride (0.664 ml) and 1 drop of N,N-dimethylformamide and the mixture was stirred at ambient temperature for 2 hours.

10 After being removed a solvent by evaporation, residual acid chloride in dichloromethane (5 ml) was added to a mixture of 2-[4,4-dimethyl(2,5-oxazolinyl)]phenylamine (724 mg) and triethylamine (770 mg) in dichloromethane (15 ml) at 0°C and the mixture was stirred at ambient temperature for 2 hours.

15 After being removed a solvent by evaporation, the residue was diluted with ethyl acetate and washed successively with 1N hydrochloric acid, saturated aqueous sodium hydrogencarbonate and brine and dried over sodium sulfate. The solvent was removed by rotary evaporation to give N-[2-[4,4-dimethyl(2,5-oxazolinyl)]phenyl]-3-methoxy-4-methoxycarbonylbenzamide
20 (1.46 g).

NMR (CDCl₃, δ) : 1.44 (6H, s), 3.94 (3H, s), 4.01 (3H, s), 4.12 (3H, s), 7.16 (1H, dd, J=8, 8Hz), 7.53 (1H, dd, J=8, 8Hz), 7.72 (1H, d, J=8Hz), 7.78 (1H, s), 7.85-7.94 (2H, m), 8.92 (1H, d, J=8Hz)
25

Preparation 63

The following compounds were obtained according to a similar manner to that of Preparation 62.

30

1) N-[2-[4,4-Dimethyl(2,5-oxazolinyl)]phenyl]-3-methoxy-4-nitrobenzamide

NMR (CDCl₃, δ) : 1.43 (6H, s), 4.08 (3H, s), 4.13 (2H, s), 7.18 (1H, dd, J=8, 8Hz), 7.56 (1H, dd, J=8, 8Hz), 7.78 (1H, d, J=8Hz), 7.86-8.00 (3H, m), 8.91
35

(1H, d, J=8Hz)

- 2) 3-Methoxy-4-methoxycarbonyl-N-[2-(morpholin-4-yl)phenyl]benzamide

5 NMR (CDCl₃, δ) : 2.87-2.98 (4H, m), 3.82-3.90 (4H, m),
3.91 (3H, s), 4.00 (3H, s), 7.09-7.18 (1H, m),
7.20-7.30 (1H, m), 7.39 (1H, d, J=8Hz), 7.66 (1H,
s), 7.92 (1H, d, J=8Hz), 8.54 (1H, d, J=8Hz), 9.58
(1H, s)

10

- 3) 3-Methoxy-N-[2-(morpholin-4-yl)phenyl]-4-nitrobenzamide
NMR (CDCl₃, δ) : 2.88-3.08 (4H, br), 3.82-3.99 (4H,
br), 4.08 (3H, s), 7.12-7.36 (3H, m), 7.46 (1H, m),
7.82 (1H, s), 7.97 (1H, d, J=8Hz), 8.48 (1H, m)

15

- 4) 3-Methoxy-4-methoxycarbonyl-N-[2-(1-pyrrolyl)phenyl]-
benzamide

20 NMR (CDCl₃, δ) : 3.90 (3H, s), 3.92 (3H, s), 6.42-6.49
(2H, m), 6.82-6.90 (2H, m), 7.04-7.12 (1H, m),
7.18-7.32 (2H, m), 7.39 (1H, d, J=8Hz), 7.48 (1H,
dd, J=8, 8Hz), 7.70-7.83 (2H, m), 8.62 (1H, d,
J=8Hz)

- 5) 3-Methoxy-4-methoxycarbonyl-N-(2-piperidinophenyl)-
benzamide

25 NMR (CDCl₃, δ) : 1.54-1.81 (6H, m), 2.79-2.90 (4H, m),
3.92 (3H, s), 4.01 (3H, s), 7.04-7.15 (1H, m), 7.20
(1H, dd, J=8, 8Hz), 7.42 (1H, d, J=8Hz), 7.66 (3H,
s), 7.93 (1H, d, J=8Hz), 8.54 (1H, d, J=8Hz), 9.68
30 (1H, s)

- 6) 3-Methoxy-4-nitro-N-(2-piperidinophenyl)benzamide

35 NMR (CDCl₃, δ) : 1.56-1.82 (6H, m), 2.78-2.92 (4H, m),
4.06 (3H, s), 7.08-7.29 (3H, m), 7.44 (1H, m), 7.77
(1H, s), 7.97 (1H, d, J=8Hz), 8.51 (1H, d, J=8Hz),

150

9.71 (1H, s)

- 7) 3-Methoxy-4-methoxycarbonyl-N-[2-(4-methyl-1-piperazinyl)phenyl]benzamide
5 NMR (CDCl₃, δ) : 2.38 (3H, s), 2.48-2.79 (4H, br),
2.91-3.02 (4H, m), 3.94 (3H, s), 4.02 (3H, s), 7.11
(1H, dd, J=8, 8Hz), 7.18-7.30 (2H, m), 7.42 (1H, d,
J=8Hz), 7.65 (1H, s), 7.92 (1H, d, J=8Hz), 8.54
(1H, d, J=8Hz), 9.58 (1H, s)
- 10 8) 3-Methoxy-N-[2-(4-methyl-1-piperazinyl)phenyl]-4-nitrobenzamide
NMR (CDCl₃, δ) : 2.42 (3H, s), 2.53-2.80 (4H, m), 2.92-
3.06 (4H, m), 4.07 (3H, s), 7.11-7.20 (1H, m),
15 7.21-7.32 (2H, m), 7.45 (1H, m), 7.79 (1H, s), 7.98
(1H, d, J=8Hz), 8.52 (1H, d, J=8Hz)
- 9) 3-Methoxy-4-methoxycarbonyl-N-[2-(2,5-oxazolyl)phenyl]-benzamide
20 NMR (CDCl₃, δ) : 1.63 (1H, br s), 3.94 (3H, s), 4.04
(3H, s), 7.22 (1H, dd, J=8, 8Hz), 7.31 (1H, s),
7.53 (1H, m), 7.72-7.82 (3H, m), 7.94 (1H, d,
J=8Hz), 8.09 (1H, m), 8.98 (1H, d, J=8Hz)
- 25 10) 3-Methoxy-4-methoxycarbonyl-N-[2-(2,5-oxazolinyl)-phenyl]benzamide
NMR (CDCl₃, δ): 3.92 (3H, s), 4.02 (3H, s), 4.20 (9H, t,
J=8Hz), 4.44 (9H, t, J=8Hz), 7.14 (1H, dd, J=8, 8Hz),
7.54 (1H, dd, J=8, 8Hz), 7.69 (1H, d, J=8Hz), 7.75
30 (1H, s), 7.87-7.96 (2H, m), 8.95 (1H, d, J=8Hz)
- 11) 3-Methoxy-4-methoxycarbonyl-N-[2-(3H,4H,5H-2,6-oxazinyl)phenyl]benzamide
NMR (CDCl₃, δ) : 1.98-2.11 (2H, m), 3.71 (2H, t,
35 J=7Hz), 3.93 (3H, s), 4.00 (3H, s), 4.43 (2H, t,

151

J=7Hz), 7.09 (1H, dd, J=8, 8Hz), 7.47 (1H, dd, -J=8, 8Hz), 7.56 (1H, d, J=8Hz), 7.70 (1H, s), 7.89 (1H, d, J=8Hz), 7.93 (1H, d, J=8Hz), 8.89 (1H, d, J=8Hz)

- 5 12) N-[2-(1-Aza-3-oxaspiro[4.4]non-1-en-2-yl)phenyl]-3-methoxy-4-methoxycarbonylbenzamide

NMR (CDCl₃, δ) : 1.70-1.84 (4H, m), 1.85-2.09 (4H, m),
3.93 (3H, s), 4.00 (3H, s), 4.26 (2H, s), 7.15 (1H, dd, J=8, 8Hz), 7.52 (1H, dd, J=8, 8Hz), 7.69 (1H, d, J=8Hz), 7.77 (1H, s), 7.84 (1H, d, J=8Hz), 7.91 (1H, d, J=8Hz), 8.94 (1H, d, J=8Hz)

Preparation 64

To a solution of N-[2-[4,4-dimethyl(2,5-oxazolinyl)]-phenyl]-3-methoxy-4-methoxycarbonylbenzamide (1.45 g) in N,N-dimethylformamide (18 ml) was added portionwise sodium hydride (167 mg) at 0°C and the mixture was stirred at 0°C for 30 minutes. Methyl iodide (0.283 ml) was added to the mixture and the solution was stirred at 0°C for 1 hour. The reaction was quenched with water and then the aqueous solution was extracted with ethyl acetate. The organic layer was washed with brine and dried over sodium sulfate. The solvent was evaporated in vacuo to give N-[2-[4,4-dimethyl(2,5-oxazolinyl)]phenyl]-N-methyl-3-methoxy-4-methoxycarbonylbenzamide (1.5 g).

NMR (CDCl₃, δ) : 1.35 (3H, s), 1.36 (3H, s), 3.33 (3H, s), 3.63 (3H, s), 4.00-4.14 (2H, m), 6.93-7.09 (3H, m), 7.18-7.36 (2H, m), 7.57 (1H, d, J=8Hz), 7.78 (1H, d, J=8Hz)

Preparation 65

The following compounds were obtained according to a similar manner to that of Preparation 64.

- 35 1) N-[2-[4,4-Dimethyl(2,5-oxazolinyl)]phenyl]-N-methyl-3-

152

methoxy-4-nitrobenzamide

NMR (CDCl₃, δ) : 1.38 (3H, s), 1.39 (3H, s), 3.35 (3H, s), 3.71 (3H, s), 4.02-4.16 (2H, m), 7.00-7.10 (2H, m), 7.21 (1H, s), 7.23-7.39 (2H, m), 7.63 (1H, d, J=8Hz), 7.80 (1H, m)

2) 3-Methoxy-4-methoxycarbonyl-N-methyl-N-[2-(morpholin-4-yl)phenyl]benzamide

NMR (CDCl₃, δ) : 2.32-2.48 (2H, m), 2.78-2.91 (2H, m), 3.50 (3H, s), 3.61 (3H, s), 3.62-3.80 (4H, m), 3.82 (3H, s), 6.86 (1H, d, J=8Hz), 6.91 (1H, s), 7.02-7.32 (4H, m), 7.59 (1H, d, J=8Hz)

3) 3-Methoxy-N-methyl-N-[2-(morpholin-4-yl)phenyl]-4-nitrobenzamide

NMR (CDCl₃, δ) : 2.28-2.45 (2H, m), 2.77-2.92 (2H, m), 3.50 (3H, s), 3.57-3.82 (7H, m), 6.87 (1H, d, J=8Hz), 7.02 (1H, s), 7.04-7.18 (2H, m), 7.19-7.33 (2H, m), 7.61 (1H, d, J=8Hz)

4) 3-Methoxy-4-methoxycarbonyl-N-methyl-N-[2-(1-pyrrolyl)phenyl]benzamide

NMR (CDCl₃, δ) : 3.43 (3H, s), 3.66 (3H, s), 3.81 (3H, s), 6.25 (2H, s), 6.38-6.51 (3H, m), 6.56 (1H, s), 7.12 (1H, m), 7.21-7.51 (4H, m)

5) 3-Methoxy-4-methoxycarbonyl-N-methyl-N-(2-piperidinophenyl)benzamide

NMR (CDCl₃, δ) : 1.43-1.72 (6H, m), 2.29-2.44 (2H, m), 2.70-2.84 (2H, m), 3.50 (3H, s), 3.60 (3H, s), 3.81 (3H, s), 6.84 (1H, d, J=8Hz), 6.90 (1H, s), 6.98-7.09 (2H, m), 7.15 (1H, dd, J=8, 8Hz), 7.22 (1H, d, J=8Hz), 7.59 (1H, d, J=8Hz)

6) 3-Methoxy-N-methyl-4-nitro-N-(2-piperidinophenyl)-

benzamide

NMR (CDCl₃, δ) : 1.42-1.76 (6H, m), 2.23-2.41 (2H, m),
2.70-2.87 (2H, m), 3.53 (3H, s), 3.68 (3H, s), 6.87
(1H, d, J=8Hz), 7.01 (1H, s), 7.03-7.14 (2H, m),
5 7.20 (1H, dd, J=8, 8Hz), 7.27 (1H, m), 7.64 (1H, d,
J=8Hz)

7) N-Methyl-3-methoxy-4-methoxycarbonyl-N-[2-(4-methyl-1-
piperazinyl)phenyl]benzamide

10 NMR (CDCl₃, δ) : 2.34 (3H, s), 2.39-2.61 (6H, m), 2.82-
2.99 (2H, m), 3.49 (3H, s), 3.61 (3H, s), 3.82 (3H,
s), 6.81-6.93 (2H, m), 7.00-7.11 (2H, m), 7.12-7.29
(2H, m), 7.60 (1H, d, J=8Hz)

15 8) N-Methyl-3-methoxy-N-[2-(4-methyl-1-piperazinyl)phenyl]-
4-nitrobenzamide

NMR (CDCl₃, δ) : 2.34 (3H, s), 2.35-2.62 (6H, m), 2.85-
3.00 (2H, m), 3.52 (3H, s), 3.68 (3H, s), 6.89 (1H,
d, J=8Hz), 7.01 (1H, s), 7.04-7.32 (4H, m), 7.64
20 (1H, d, J=8Hz)

9) 3-Methoxy-4-methoxycarbonyl-N-methyl-N-[2-(2,5-
oxazolyl)phenyl]benzamide

25 NMR (CDCl₃, δ) : 3.45 (3H, s), 3.60 (3H, s), 3.80 (3H,
s), 6.69 (1H, d, J=8Hz), 6.72 (1H, s), 7.22-7.42
(4H, m), 7.48 (1H, d, J=8Hz), 7.76 (1H, s), 7.89
(1H, d, J=8Hz)

30 10) N-Methyl-3-methoxy-4-methoxycarbonyl-N-[2-(2,5-
oxazolyl)phenyl]benzamide

NMR (CDCl₃, δ) : 3.40 (3H, s), 3.67 (3H, s), 3.81 (3H,
s), 4.02-4.12 (2H, m), 4.30-4.46 (2H, m), 6.88-6.98
(2H, m), 7.12 (1H, d, J=8Hz), 7.24 (1H, dd, J=8,
8Hz), 7.35 (1H, dd, J=8, 8Hz), 7.57 (1H, d, J=8Hz),
35 7.76 (1H, d, J=8Hz)

- 11) N-Methyl-3-methoxy-4-methoxycarbonyl-N-[2-(3H, 4H, 5H-2, 6-oxaziny)phenyl]benzamide

5 NMR (CDCl₃, δ) : 1.92-2.08 (2H, m), 3.40 (3H, s), 3.58 (2H, t, J=7Hz), 3.64 (3H, s), 3.82 (3H, s), 4.27-4.40 (2H, m), 6.93-7.00 (1H, m), 7.01-7.09 (2H, m), 7.16-7.28 (2H, m), 7.51-7.65 (2H, m)

- 12) N-[2-(1-Aza-3-oxaspiro[4.4]non-1-en-2-yl)phenyl]-3-methoxy-4-methoxycarbonyl-N-methylbenzamide

10 NMR (CDCl₃, δ) : 1.59-1.77 (4H, m), 1.80-2.05 (4H, m), 3.36 (3H, s), 3.65 (3H, s), 3.80 (3H, s), 4.15-4.27 (2H, m), 6.93-7.03 (2H, m), 7.06 (1H, d, J=8Hz), 7.18-7.35 (2H, m), 7.58 (1H, d, J=8Hz), 7.75 (1H, m)

15

Preparation 66

To a solution of 2-(1-pyrrolyl)nitrobenzene (1.11 g) in ethanol (30 ml) were added iron powder (1.65 g) and acetic acid (3.54 g) and the mixture was refluxed for 1 hour. The reaction mixture was filtered through a bed of celite and the filtrate was concentrated in vacuo. The residue was diluted with a mixture of ethyl acetate and saturated aqueous sodium bicarbonate solution and the mixture was filtered through a bed of celite again. The organic layer was separated and washed with water and brine. The solution was dried over sodium sulfate and the solvent was evaporated in vacuo to give 2-(1-pyrrolyl)phenylamine (860 mg).

20

25

NMR (CDCl₃, δ) : 3.72 (2H, br s), 6.31-6.40 (2H, m), 6.72-6.90 (4H, m), 7.10-7.24 (2H, m)

30

Preparation 67

The following compounds were obtained according to a similar manner to that of Preparation 66.

35

- 1) 4-Amino-3-methoxy-N-methyl-N-(2-piperidinophenyl)-

155

benzamide

5 NMR (CDCl₃, δ) : 1.43-1.71 (6H, m), 2.54-2.68 (2H, m),
2.71-2.86 (2H, m), 3.46 (3H, s), 3.57 (3H, s), 3.86
(2H, s), 6.41 (1H, d, J=8Hz), 6.83 (1H, s), 6.86-
7.00 (3H, m), 7.07-7.19 (2H, m)

2) 4-Amino-N-methyl-3-methoxy-N-[2-(4-methyl-1-
piperazinyl)phenyl]benzamide

10 NMR (CDCl₃, δ) : 2.35 (3H, s), 2.41-2.61 (4H, m), 2.66-
2.80 (2H, m), 2.84-3.00 (2H, m), 3.46 (3H, s), 3.58
(3H, s), 3.88 (2H, s), 6.42 (1H, d, J=8Hz), 6.82
(1H, s), 6.86-7.06 (3H, m), 7.09-7.21 (2H, m)

3) 2-(2,5-Oxazolinyl)phenylamine

15 NMR (CDCl₃, δ) : 4.10 (2H, t, J=8Hz), 4.34 (2H, t,
J=8Hz), 6.60-6.76 (2H, m), 7.20 (1H, dd, J=8, 8Hz),
7.71 (1H, d, J=8Hz)

4) 2-(3H,4H,5H-2,6-Oxazinyl)phenylamine

20 NMR (CDCl₃, δ) : 1.91-2.02 (2H, m), 3.63 (2H, t,
J=7Hz), 4.35 (2H, t, J=7Hz), 6.20 (2H, br s), 6.56-
6.68 (2H, m), 7.12 (1H, dd, J=8, 8Hz), 7.70 (1H, d,
J=8Hz)

25 5) 4-Amino-2-(N,N-dimethylcarbamoyl)-1H-benzimidazole

30 NMR (DMSO-d₆, δ) : 3.06 (3H x 3/5, s), 3.09 (3H x 2/5,
s), 3.65 (3H x 3/5, s), 3.19 (3H x 2/5, s), 5.35
(2H x 3/5, s), 5.48 (2H x 2/5, s), 6.36 (1H x 3/5,
d, J=8Hz), 6.45 (1H x 2/5, d, J=8Hz), 6.66 (1H x
3/5, d, J=8Hz), 6.88-7.02 (1H + 1H x 2/5, m), 7.15
(1H, br peak)

Preparation 68

35 To a solution of 3-nitro-1,2-phenylenediamine (1.0 g)
and triethylamine (793 mg) in dichloromethane (3 ml) under

nitrogen was added portionwise phthalylglycyl chloride (1.61 g) in ice water bath and the mixture was stirred at the same temperature for 2 hours. The reaction mixture was washed with saturated sodium bicarbonate aqueous solution, dried over magnesium sulfate and evaporated in vacuo. To the resulting crude product was added polyphosphoric acid (5 ml) and stirred at 130°C for 3 hours. After the mixture was cooled at ambient temperature, ammonia solution (28%) was added to the reaction mixture in ice water bath. The precipitate was collected by vacuum filtration to give 2-phthalimidomethyl-4-nitro-1H-benzimidazole (1.35 g).

NMR (DMSO-d₆, δ) : 5.13 (2H, s), 7.36 (1H, t, J=8Hz), 7.86-8.05 (5H, m), 8.11 (1H, d, J=8Hz)

15 Preparation 69

The following compound was obtained according to a similar manner to that of Preparation 68.

4-Nitro-2-(2-phthalimidoethyl)-1H-benzimidazole

20 NMR (DMSO-d₆, δ) : 3.28 (2H, t, J=7.5Hz), 4.05 (2H, t, J=7.5Hz), 7.34 (1H, t, J=8Hz), 7.78-7.90 (4H, m), 7.96 (1H, d, J=8Hz), 8.08 (1H, d, J=8Hz)

Preparation 70

25 To a solution of 2-chloro-4-nitro-1H-benzimidazole (300 mg) in N-methyl-2-pyrrolidone (4 ml) was added imidazole (517 mg) and the mixture was stirred at 80°C for 8 hours. The reaction mixture was poured into brine and extracted with a mixture of chloroform and methanol. The organic layer was dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed on silica gel eluting with a mixture of chloroform and methanol (100-0 - 30-1) to give 2-(1-imidazolyl)-4-nitro-1H-benzimidazole (175 mg).

35 NMR (DMSO-d₆, δ) : 7.21 (1H, s), 7.46 (1H, t, J=8Hz), 8.05-8.19 (3H, m), 8.70 (1H, s)

Preparation 71

The following compound was obtained according to a similar manner to that of Preparation 70.

- 5 4-Nitro-2-(1,2,4-tetrazol-1-yl)-1H-benzimidazole
NMR (DMSO-d₆, δ) : 7.50 (1H, t, J=8Hz), 8.06 (1H, br
peak), 8.17 (1H, d, J=8Hz), 8.50 (1H, s), 9.53 (1H,
s)

10 Preparation 72

A mixture of 2-chloro-4-nitro-1H-benzimidazole (300 mg) and N,N-dimethylethylenediamine (2 ml) were stirred at 80°C for 8 hours. The reaction mixture was poured into saturated sodium bicarbonate aqueous solution and extracted with a mixture of chloroform and methanol. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was washed with diisopropyl ether to give 2-[[2-(dimethylamino)ethyl]amino]-4-nitro-1H-benzimidazole (113 mg).

- 20 NMR (CDCl₃, δ) : 2.65 (3H, s), 2.90-2.98 (2H, m), 3.55-
3.64 (2H, m), 7.11 (1H, t, J=8Hz), 7.69 (1H, d,
J=8Hz), 7.87 (1H, d, J=8Hz)

Preparation 73

25 The following compounds were obtained according to a similar manner to that of Preparation 72.

- 1) 2-(4-Methylpiperazin-1-yl)-4-nitro-1H-benzimidazole
NMR (CDCl₃, δ) : 2.42 (3H, s), 2.58-2.72 (4H, m), 3.63-
30 3.80 (4H, m), 7.19 (1H, t, J=8Hz), 7.68 (1H, d,
J=8Hz), 7.85 (1H, d, J=8Hz), 9.44 (1H, br s)
- 2) 2-Dimethylamino-4-nitro-1H-benzimidazole
NMR (CDCl₃, δ) : 3.30 (6H, s), 7.21 (1H, t, J=8Hz),
35 7.76 (1H, d, J=8Hz), 7.86 (1H, d, J=8Hz)

- 3) 2-[(2-Aminoethyl)methylamino]-4-nitro-1H-benzimidazole
NMR (CDCl₃, δ) : 3.09 (2H, t, J=5Hz), 3.26 (3H, s),
3.60 (2H, t, J=5Hz), 7.15 (1H, t, J=8Hz), 7.61 (1H,
d, J=8Hz), 7.83 (1H, d, J=8Hz)

5

- 4) 2-[(2-Methylamino)ethyl]amino-4-nitro-1H-benzimidazole
NMR (CDCl₃, δ) : 2.68 (3H, s), 3.09 (2H, t-like,
J=5Hz), 3.59 (2H, t-like, J=5Hz), 7.02 (1H, t,
J=8Hz), 7.51 (1H, d, J=8Hz), 7.74 (1H, d, J=8Hz)

10

- 5) 2-[[2-(Dimethylamino)ethyl]methylamino]-4-nitro-1H-
benzimidazole
NMR (CDCl₃, δ) : 2.58 (6H, s), 2.81-2.91 (2H, m), 3.34
(3H, s), 3.50-3.60 (2H, m), 7.09 (1H, t, J=8Hz),
7.70 (1H, d, J=8Hz), 7.83 (1H, d, J=8Hz)

15

- 6) 2-[(2-Methoxyethyl)amino]-4-nitro-1H-benzimidazole
NMR (CDCl₃, δ) : 3.50 (3H, s), 3.60-3.73 (4H, m), 5.60
(1H, br peak), 7.15 (1H, t, J=8Hz), 7.63 (1H, d,
J=8Hz), 7.83 (1H, d, J=8Hz)

20

Preparation 74

To a solution of ethyl 2-chloromethyl-1H-benzimidazole-
4-carboxylate (250 mg) in dichloroethane (2.5 ml) was added
morpholine (183 mg) under ice bath cooling and the reaction
mixture was stirred at ambient temperature for 15 hours. To
the reaction mixture was added morpholine (91 mg) and stirred
at 80°C for 6 hours. After the reaction mixture was
concentrated in vacuo, the residue was diluted with
chloroform and saturated sodium hydrogen carbonate aqueous
solution. The organic layer was separated and washed with
brine. The solution was dried over magnesium sulfate and
concentrated in vacuo. The residue was purified by silica
gel column chromatography (methanol:chloroform = 1:9) to give
ethyl 2-morpholinomethyl-1H-benzimidazole-4-carboxylate (219

35

159

mg).

NMR (CDCl₃, δ) : 1.47 (3H, t, J=7Hz), 2.57-2.60 (4H, m), 3.75-3.77 (4H, m), 3.86 (2H, s), 4.47 (2H, q, J=8Hz), 7.28 (1H, t, J=8Hz), 7.90 (1H, d, J=8Hz), 7.91 (1H, d, J=8Hz)

Preparation 75

The following compounds were obtained according to a similar manner to that of Preparation 74.

10

1) 2-(N,N-Dimethylaminomethyl)-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 2.36 (6H, s), 3.81 (2H, s), 7.35 (1H, t, J=8Hz), 8.03 (1H, d, J=8Hz), 8.15 (1H, d, J=8Hz)

15

2) 2-(1-Imidazolyl)methyl-4-nitro-1H-benzimidazole

NMR (DMSO-d₆, δ) : 5.53 (2H, s), 6.91 (1H, s), 7.28 (1H, s), 7.40 (1H, t, J=8Hz), 7.80 (1H, s), 8.08 (1H, d, J=8Hz), 8.14 (1H, d, J=8Hz)

20

3) 2-[(4-Methylpiperazin-1-yl)methyl]-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 2.40 (3H, s), 2.49-2.82 (8H, m), 3.91 (2H, s), 7.35 (1H, t, J=8Hz), 8.04 (1H, d, J=8Hz), 8.16 (1H, d, J=8Hz)

25

4) 2-(Morpholin-4-ylmethyl)-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 2.55-2.69 (4H, m), 3.73-3.85 (4H, m), 3.89 (2H, s), 7.35 (1H, t, J=8Hz), 8.05 (1H, d, J=8Hz), 8.15 (1H, d, J=8Hz)

30

5) 4-Nitro-2-(pyrrolidin-1-ylmethyl)-1H-benzimidazole

NMR (CDCl₃, δ) : 1.96 (4H, br s), 2.86 (4H, br s), 4.19 (2H, s), 7.36 (1H, t, J=8Hz), 8.03 (1H, d, J=8Hz), 8.16 (1H, d, J=8Hz)

35

6) 4-Nitro-2-(piperidinomethyl)-1H-benzimidazole

NMR (CDCl₃, δ) : 1.58 (2H, br peak), 1.73 (4H, br peak), 2.65 (4H, br peak), 4.00 (2H, br s), 7.36 (1H, t, J=8Hz), 8.03 (1H, d, J=8Hz), 8.17 (1H, d, J=8Hz)

5

7) 2-[2-(Dimethylamino)ethyl]-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 2.50 (6H, s), 2.83 (2H, t, J=7Hz), 3.18 (2H, t, J=8Hz), 7.30 (1H, t, J=8Hz), 8.00 (1H, d, J=8Hz), 8.11 (1H, d, J=8Hz)

10

8) 2-[2-(4-Methylpiperazin-1-yl)ethyl]-4-nitro-1H-benzimidazole

NMR (CDCl₃, δ) : 2.41 (3H, s), 2.70 (8H, br peak), 2.89 (2H, t, J=5Hz), 3.18 (2H, t, J=5Hz), 7.32 (1H, t, J=8Hz), 8.01 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz)

15

9) Ethyl 2-dimethylaminomethyl-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.44 (3H, t, J=7Hz), 2.33 (6H, s), 3.77 (2H, s), 4.45 (2H, q, J=7Hz), 7.27 (1H, t, J=8Hz), 7.88-7.92 (2H, m)

20

10) Ethyl 2-(4-methylpiperazin-1-yl)methyl-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.47 (3H, t, J=7Hz), 2.51 (4H, br s), 2.62 (4H, br s), 3.86 (2H, s), 4.47 (2H, q, J=8Hz), 7.26 (1H, t, J=8Hz), 7.90 (2H, d, J=8Hz)

25

11) Ethyl 2-(4-dimethylaminopiperidino)methyl-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.46 (3H, t, J=7Hz), 1.61 (2H, dt, J=2, 8Hz), 1.82 (1H, br s), 2.12-2.23 (4H, m), 2.28 (6H, s), 2.92-2.98 (2H, m), 3.82 (2H, s), 4.48 (2H, q, J=7Hz), 7.28 (1H, t, J=8Hz), 7.89 (2H, d, J=8Hz)

30

35

Preparation 76

A suspension of 2-methyl-4-nitrobenzimidazole (2.2 g) in 1,4-dioxane (35 ml) was treated with triethylamine (2.51 g) and di-tert-butyl dicarbonate (5.42 g). After 15 minutes, to the reaction mixture was added N,N-dimethylaminopyridine (catalytic amount). The solution was stirred a further 20 hours and concentrated in vacuo and the residue was dissolved in ethyl acetate. The solution was washed with 1N hydrochloric acid, water, saturated aqueous sodium hydrogen carbonate and brine. The organic layer was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by silica gel column chromatography (chloroform) to give tert-butyl 2-methyl-4-nitro-1H-benzimidazole-1-carboxylate (3.0 g).

NMR (CDCl₃, δ) : 1.73 (9H, s), 2.93 (3H, s), 7.41 (1H, dd, J=8, 8Hz), 8.13 (1H, d, J=8Hz), 8.29 (1H, d, J=8Hz)

Preparation 77

The following compounds were obtained according to a similar manner to that of Preparation 76.

1) Ethyl 2-(N-tert-butoxycarbonylaminomethyl)-1-methyl-1H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.40-1.50 (12H, m), 3.34 (3H, s), 4.49 (2H, q, J=7.5Hz), 4.70 (2H, d, J=7Hz), 5.48 (1H, br peak), 7.33 (1H, t, J=8Hz), 7.53 (1H, d, J=8Hz), 7.94 (1H, d, J=8Hz)

2) Ethyl 2-(N-tert-butoxycarbonylaminomethyl)-3-methyl-3H-benzimidazole-4-carboxylate

NMR (CDCl₃, δ) : 1.44 (3H, t, J=7.5Hz), 1.48 (9H, s), 3.93 (3H, s), 4.44 (2H, q, J=7.5Hz), 4.64 (2H, d, J=5Hz), 5.51 (1H, br peak), 7.25 (1H, t, J=8Hz), 7.77 (1H, d, J=8Hz), 7.88 (1H, d, J=8Hz)

162

- 3) 2-[(tert-Butoxy)carbonylamino]-4-nitro-1H-benzimidazole
NMR (DMSO-d₆, δ) : 1.68 (9H, s), 7.11 (1H, t, J=8Hz),
7.84-8.00 (3H, m)

5 Preparation 78

To a solution of N-(2-chloroethyl)-2-nitrobenzamide (3.63 g) in acetonitrile (100 ml) was slowly added 40% potassium fluoride on alumina (10 g). This slurry was stirred at ambient temperature for 24 hours. The potassium fluoride on alumina was filtered through a bed of celite, washed with ethyl acetate. The solvent was evaporated to give 2-(2,5-oxazolinyl)nitrobenzene (3.0 g).

NMR (CDCl₃, δ) : 4.09 (2H, t, J=8Hz), 4.45 (2H, t, J=8Hz), 7.58-7.69 (2H, m), 7.80-7.89 (2H, m)

15

Preparation 79

The following compound was obtained according to a similar manner to that of Preparation 78.

20 2-(2-Nitrophenyl)-4H,5H,6H-1,3-oxazine

NMR (CDCl₃, δ) : 1.96-2.11 (2H, m), 3.56-3.68 (2H, m), 4.25-4.37 (2H, m), 7.54 (1H, dd, J=8, 8Hz), 7.61 (1H, dd, J=8, 8Hz), 7.71 (1H, d, J=8Hz), 7.85 (1H, d, J=8Hz)

25

Preparation 80

The following compound was obtained according to a similar manner to that of Example 84.

30 Benzyl 2-formylindole-4-carboxylate

NMR (DMSO-d₆, δ) : 5.45 (2H, s), 7.30-7.55 (6H, m), 7.73-7.82 (2H, m), 7.88 (1H, d, J=8Hz), 9.93 (1H, s)

35 Preparation 81

The following compound was obtained according to a similar manner to that of Example 107.

4-Methyl-1-(2-nitrophenyl)piperazine

5 NMR (CDCl₃, δ) : 2.36 (3H, s), 2.52-2.61 (4H, m), 3.03-3.12 (4H, m), 7.03 (1H, dd, J=8, 8Hz), 7.15 (1H, d, J=8Hz), 7.48 (1H, dd, J=8, 8Hz), 7.75 (1H, d, J=8Hz)

10 Preparation 82

To a solution of indole-4-carboxylic acid (500 mg) in methanol (9 ml) and conc. hydrochloric acid (1.0 ml) was added a portion of sodium cyanoborohydride (487 mg) at 0°C and the mixture was stirred at ambient temperature for 1
15 hour. The suspension was diluted with water (10 ml) and then the clear solution was neutralized with 2N sodium hydroxide aqueous solution. Methanol was removed and the aqueous solution was diluted with dioxane (15 ml) and 1N sodium hydroxide aqueous solution (10 ml). To the mixture was added
20 portionwise di-tert-butyl dicarbonate (813 mg) and the solution was stirred at ambient temperature for 2 hours. The solution was neutralized with 1N hydrochloric acid and diluted with ethyl acetate (30 ml). The resulting solution was washed with brine, dried over magnesium sulfate and
25 concentrated in vacuo. The solid was triturated with diethyl ether:n-hexane (1:9) to give 1-tert-butoxycarbonylindoline-4-carboxylic acid (727 mg).

NMR (DMSO-d₆, δ) : 1.50 (9H, s), 3.36 (2H, t, J=9Hz),
3.92 (2H, t, J=9Hz), 7.27 (1H, t, J=8Hz), 7.49 (1H,
30 d, J=8Hz), 7.80-7.94 (1H, br)

Preparation 83

To a solution of methyl trans-2-[β -(dimethylamino)vinyl]-3-nitrobenzoate (3.16 g) in
35 tetrahydrofuran (20 ml) and pyridine (3.57 ml) was added

dropwise benzyloxyacetyl chloride (4.2 g) at ambient temperature and the mixture was refluxed for 3 hours. The resulting mixture was diluted with ethyl acetate and the solution was washed successively with water, saturated sodium bicarbonate aqueous solution and brine. Drying, filtering and removal of solvents afforded a crude product. The crude product was purified by column chromatography to give 2-[β -(dimethylamino)- α -(benzyloxyacetyl)vinyl]-3-nitrobenzoate (3.32 g).

10 NMR (CDCl₃, δ) : 2.70-2.80 (6H, br s), 3.80 (3H, s), 4.08-4.16 (2H, br s), 4.50 and 4.51 (Total 2H, s), 7.28-7.38 (5H, m), 7.49 (1H, t, J=8Hz), 7.79 (1H, d, J=8Hz), 7.80 (1H, s), 7.89 (1H, d, J=8Hz)

15 Preparation 84

A solution of 2-[β -(dimethylamino)- α -(benzyloxyacetyl)vinyl]-3-nitrobenzoate (3.31 g) and p-toluenesulfonic acid hydrate (632 mg) in 1,4-dioxane (15 ml) and water (5 ml) was refluxed for 24 hours. The resulting mixture was evaporated in vacuo and the residue was diluted with ethyl acetate. The organic layer was washed successively with water and brine. Drying, filtering and removal of solvents afforded a crude product as a dark-red oil. The crude product was purified by silica gel column chromatography (eluent; hexane:ethyl acetate = 3:1) to give 3-benzyloxymethyl-5-nitroisocoumarin (1.0 g).

25 NMR (CDCl₃, δ) : 4.40 (2H, s), 4.69 (2H, s), 7.29-7.42 (5H, m), 7.43 (1H, s), 7.62 (1H, t, J=8Hz), 8.47 (1H, d, J=8Hz), 8.60 (1H, d, J=8Hz),

30

Preparation 85

A solution of 3-benzyloxymethyl-5-nitroisocoumarin (850 mg) in methanol (40.0 ml) was treated with aqueous titanium trichloride (11.2 ml), added as a single portion. After stirring 2 hours at ambient temperature, water (100 ml) and chloroform (120 ml) were added. The whole was carefully basified with saturated sodium bicarbonate aqueous solution and the organic layer was separated. The aqueous layer was further extracted with chloroform (120 ml) and the combined extract was washed with water, dried over magnesium sulfate, and concentrated to afford 3-benzyloxymethyl-5-aminoisocoumarin (806 mg).

NMR (CDCl₃, δ) : 3.89-4.00 (2H, br), 4.38 (2H, s), 4.69 (2H, s), 6.50 (1H, s), 7.01 (1H, d, J=9Hz), 7.29 (1H, t, J=9Hz), 7.30-7.40 (5H, m), 7.73 (1H, d, J=9Hz)

Preparation 86

A mixture of 5-amino-3-benzyloxymethylisocoumarin (800 mg) and sodium methylate in methanol (768 mg) was stirred at ambient temperature for 20 minutes. After removal of solvents, water (40 ml) was added to the residue and the whole was extracted with chloroform. The extract was washed with water, dried over magnesium sulfate, and evaporated to dryness to leave a crude product, which was purified by silica gel column chromatography with n-hexane:ethyl acetate (6:1) as an eluent to afford methyl 2-benzyloxymethylindole-4-carboxylate (660 mg).

NMR (CDCl₃, δ) : 3.98 (3H, s), 4.57 (2H, s), 4.78 (2H, s), 7.08 (1H, s), 7.22 (1H, t, J=8Hz), 7.29-7.39 (5H, m), 7.54 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.50-8.56 (1H, br)

Preparation 87

To a solution of methyl 2-benzyloxymethylindole-4-carboxylate (650 mg) in methanol (27 ml) and concentrated

hydrochloric acid (3.0 ml) was added sodium cyanoborohydride (968 mg) with ice-bath stirring and the mixture was stirred at ambient temperature for 2.5 hours. The resulting mixture was diluted with water (30 ml) and basified with saturated sodium bicarbonate aqueous solution. The mixture was extracted with ethyl acetate (40 ml) and the organic layer was washed successively with water and brine, dried over magnesium sulfate and concentrated in vacuo. The residue was purified by silica gel column chromatography (eluent; n-hexane:ethyl acetate = 6:1) to give methyl 2-benzyloxymethylindoline-4-carboxylate (550 mg).

NMR (CDCl₃, δ) : 3.02 (1H, dd, J=8, 17Hz), 3.41-3.58 (3H, m), 3.88 (3H, s), 4.08-4.18 (1H, m), 4.33-4.41 (1H, br), 4.55 (2H, s), 6.74 (1H, d, J=8Hz), 7.08 (1H, t, J=8Hz), 7.28-7.39 (6H, m)

Preparation 88

The Vilsmeier reagent was prepared by the dropwise addition of phosphoryl oxychloride (0.75 mg) to cooled N,N-dimethylformamide (20.0 ml) under constant stirring. A solution of methyl indole-4-carboxylate in N,N-dimethylformamide (12.0 ml) was added to the above solution at 0°C and the solution was stirred for 30 minutes. The mixture was diluted with water (40 ml) and the solution was neutralized with saturated sodium bicarbonate aqueous solution and extracted with ethyl acetate (60 ml). The organic layer was washed successively with water and brine, dried over magnesium sulfate and concentrated in vacuo. The solid was triturated with diethyl ether (7 ml) to give methyl 3-formylindole-4-carboxylate (792 mg).

NMR (CDCl₃, δ) : 4.00 (3H, s), 7.32 (1H, t, J=9Hz), 7.64 (1H, d, J=9Hz), 7.83 (1H, d, J=9Hz), 8.06 (1H, d, J=3Hz), 9.78-9.88 (1H, br), 10.53 (1H, s)

Preparation 89

To a solution of 2,2,6,6-tetramethylpiperidine (1.97 g) in tetrahydrofuran (24.0 ml) was added dropwise n-butyllithium (6.8 ml, 1.64M solution in n-hexane) at -40°C and the mixture was stirred at 0°C for 30 minutes. A solution of methyl 1-methoxymethoxyindole-4-carboxylate (1.64 g) in tetrahydrofuran (12.0 ml) was added to the above solution at -60°C and the solution was stirred at the same temperature for 30 minutes. A solution of N,N-dimethylformamide (662 mg) in tetrahydrofuran (9.0 ml) was added to the reaction mixture at -60°C and the solution was stirred at the same temperature for 2 hours. The temperature was raised to -30°C and the reaction was quenched with saturated ammonium chloride aqueous solution. The aqueous solution was extracted with ethyl acetate (80 ml) and the organic layer was washed with brine. Drying, filtering and removal of solvents afforded a crude product. The crude product was purified by silica gel column chromatography (eluent; n-hexane:ethyl acetate = 9:1) to give methyl 2-formyl-1-methoxymethoxyindole-4-carboxylate (1.15 g).

NMR (CDCl₃, δ) : 3.67 (3H, s), 4.01 (3H, s), 5.34 (2H, s), 7.50 (1H, t, J=9Hz), 7.77 (1H, d, J=9Hz), 7.81 (1H, s), 7.98 (1H, d, J=9Hz), 9.98 (1H, s)

Preparation 90

To a solution of benzyl 1-tert-butoxycarbonyl-2-hydroxymethylindole-4-carboxylate (465 mg), phthalimide (179 mg) and triphenylphosphine (640 mg) in tetrahydrofuran (15.0 ml) was added diethyl azodicarboxylate (425 mg) and the mixture was stirred at ambient temperature for 1 hour. The resulting mixture was concentrated in vacuo and the residue was chromatographed on silica gel with n-hexane:ethyl acetate (6:1). The solid was triturated with methanol to give benzyl 1-tert-butoxycarbonyl-2-phthalimidomethylindole-4-carboxylate (440 mg).

NMR (CDCl₃, δ) : 1.72 (9H, s), 5.27 (2H, s), 5.29 (2H,

168

s), 6.95 (1H, s), 7.21-7.34 (6H, m), 7.75-7.81 (2H, m), 7.89-7.94 (2H, m), 7.96 (1H, d, J=8Hz), 8.37 (1H, d, J=8Hz)

5 Preparation 91

The mixture of tert-butyl 2-methyl-4-nitro-1H-benzimidazole-1-carboxylate (3.0 g) and 10% palladium on carbon (300 mg) in methanol (25 ml) and 1,4-dioxane (60 ml) was hydrogenated under atmospheric pressure at ambient
10 temperature for 11 hours. The reaction mixture was filtered through a bed of celite and the filtrate was concentrated in vacuo to give crude product. The solid was washed with diisopropyl ether/n-hexane (1/2) to give 4-amino-2-methyl-1H-benzimidazole-1-carboxylate (2.0 g).

15 NMR (CDCl₃, δ) : 1.70 (9H, s), 2.79 (3H, s), 4.28 (2H, br), 6.58 (1H, d, J=8Hz), 7.06 (1H, dd, J=8, 8Hz), 7.25 (1H, d, J=8Hz)

Preparation 92

20 A solution of N-[(1-hydroxymethyl)cyclopentyl]-2-nitrobenzamide (2.1 g) in thionyl chloride (5.8 ml) was stirred at ambient temperature for 1 hour. To the reaction mixture was added diethyl ether and the resulting precipitate was filtered. The collected precipitate was dissolved in
25 ethyl acetate and 1N sodium hydroxide. The organic layer was washed with brine and dried over magnesium sulfate and concentrated to give 4-aza-3-(2-nitrophenyl)-2-oxaspiro[4.4]-non-3-ene (1.95 g).

30 NMR (CDCl₃, δ) : 1.62-1.79 (4H, m), 1.82-2.10 (4H, m), 4.26 (2H, s), 7.54-7.67 (2H, m), 7.80 (1H, d, J=8Hz), 7.88 (1H, d, J=8Hz)

Preparation 93

To a suspension of methyl 2-formyl-1H-benzimidazole-4-
35 carboxylate (460 mg) in a mixture of water (3.2 ml) and

t-butyl alcohol (12 ml) were added 2-methyl-2-butene (700 mg) and sodium dihydrogenphosphate (387 mg) in water bath. To the mixture was added portionwise sodium chlorite (901 mg) and stirred for 1 day at same temperature. The reaction mixture was cooled in an ice bath, adjusted to pH 4 with 1N hydrochloric acid and the precipitate was collected by vacuum filtration. The precipitate was washed with ethyl acetate and methanol to give methyl 2-carboxy-1H-benzimidazole-4-carboxylate (400 mg).

10 MASS (ES-) (m/z) : 219

Preparation 94

To a solution of 3-ethoxycarbonyl-1,2-phenylenediamine (790 mg) in tetrahydrofuran (10 ml) was added 1,1'-thiocarbonyldiimidazole (1.02 g) in ice water bath and the mixture was stirred for 20 hours at ambient temperature. The reaction solvent was concentrated in vacuo and the residue was washed with chloroform and collected by vacuum filtration to give ethyl 2-mercapto-1H-benzimidazole-4-carboxylate (665 mg).

20 NMR (DMSO-d₆, δ) : 1.33 (3H, t, J=7.5Hz), 4.42 (2H, q, J=7.5Hz), 7.23 (1H, t, J=8Hz), 7.38 (1H, d, J=8Hz), 7.66 (1H, d, J=8Hz)

25 Preparation 95

To a solution of ethyl 2-mercapto-1H-benzimidazole-4-carboxylate (500 mg) were added iodomethane (351 mg) and potassium carbonate (622 mg) at ambient temperature and the mixture was stirred for 20 hours at same temperature. The reaction mixture was poured into water and extracted with chloroform. The organic layer was washed with water and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was washed with diisopropyl ether to give ethyl 2-methylthio-1H-benzimidazole-4-carboxylate (310 mg).

35 NMR (CDCl₃, δ) : 1.44 (3H, t, J=7.5Hz), 3.80 (3H, s),

170

4.44 (2H, q, J=7.5Hz), 7.24 (1H, t, J=8Hz), 7.77-
7.88 (2H, m)

Preparation 96

5 To a solution of ethyl 2-methylthio-1H-benzimidazole-4-
carboxylate (200 mg) in dichloromethane (4 ml) was added
dropwise a solution of m-chloroperoxybenzoic acid (292 mg)
in dichloromethane (4 ml) in ice water bath under nitrogen
and the mixture was stirred for 4 hours at the same
10 temperature. The reaction mixture was poured into water and
extracted with dichloromethane. The organic layer was washed
with water, dried over magnesium sulfate and evaporated in
vacuo. The residue was purified by preparative thin-layer
chromatography (chloroform:ethyl acetate:methanol = 16:1:1) to
15 give ethyl 2-methylsulfonyl-1H-benzimidazole-4-carboxylate
(102 mg).

NMR (CDCl₃, δ) : 1.48 (3H, t, J=7.5Hz), 3.42 (3H, s),
4.50 (2H, q, J=7.5Hz), 7.49 (1H, t, J=8Hz), 8.06-
8.16 (2H, m)

20

Preparation 97

To a suspension of ethyl 2-mercapto-1H-benzimidazole-4-
carboxylate (950 mg) in 20% acetic acid aqueous solution (30
ml) at 0°C was bubbled chlorine for 30 minutes. The
25 resulting crude product was collected by vacuum filtration
and added portionwise to ammonia aqueous solution (28%, 10
ml) in ice water bath. The reaction mixture was stirred at
ambient temperature for 1 hour and adjusted to pH 5 with 1N
hydrochloric acid. The precipitate was collected by vacuum
30 filtration to give ethyl 2-sulfamoyl-1H-benzimidazole-4-
carboxylate (775 mg).

NMR (DMSO-d₆, δ) : 1.38 (3H, t, J=7.5Hz), 4.43 (2H, q,
J=7.5Hz), 7.47 (1H, t, J=8Hz), 7.97 (1H, d, J=8Hz),
8.04 (1H, d, J=8Hz)

35

Preparation 98

Thionyl chloride (8.3 ml) was added dropwise to 2-hydroxymethyl-4-nitro-1H-benzimidazole (1.15 g) at 0°C and the suspension was then heated at reflux for 3 hours. The
5 excess of thionyl chloride was removed in vacuo, and the residue was poured into ice and adjusted to pH 7 with saturated sodium bicarbonate aqueous solution. The precipitate was collected by vacuum filtration and washed
10 water to give 2-chloromethyl-4-nitro-1H-benzimidazole (1.32 g).

NMR (DMSO-d₆, δ) : 4.98 (2H, s), 7.39 (1H, t, J=8Hz),
8.09 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz)

Preparation 99

15 To a solution of 2,4-dihydroxy-8-methylquinazoline (1.00 g) in mixture of 2-methyl-2-propanol (20 ml) and water (20 ml) was added potassium permanganate (3.59 g) and magnesium sulfate (1.37 g) and the reaction mixture was stirred at 90°C for 15 hours. After the reaction mixture was filtered
20 through a bed of celite, the filtrate was diluted with water. The solution was adjusted to pH 4 with 1N hydrochloric acid. The formed precipitate was collected by vacuum filtration to give 2,4-dihydroxyquinazoline-8-carboxylic acid (390 mg).

25 NMR (DMSO-d₆, δ) : 7.31 (1H, t, J=8Hz), 8.19 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz)

Preparation 100

To a solution of ethyl 3-aminopyrazole-4-carboxylate (5.00 g) in carbon tetrachloride (70 ml) was added triethyl
30 orthoacetate (6.53 g) and the reaction mixture was stirred at 90°C for 3 hours. After the reaction mixture was concentrated in vacuo, the residue was purified by silica gel (Chromatorex, Fuji Silysia Chemical Ltd.) column chromatography (n-hexane:ethyl acetate = 1:2) to give ethyl
35 3-(1-aza-2-ethoxyprop-1-enyl)pyrazole-4-carboxylate (4.94 g).

172

NMR (CDCl₃, δ) : 1.27-1.36 (6H, m), 1.92 (3H, s), 4.22-4.36 (4H, m), 7.91 (1H, s)

Preparation 101

5 To a solution of ethyl 3-(1-aza-2-ethoxyprop-1-enyl)pyrazole-4-carboxylate (4.90 g) in N,N-dimethylformamide (50 ml) was added hydroxylamine hydrochloride (15.1 g) and stirred at ambient temperature for 3 hours. The reaction mixture was concentrated in vacuo and the residue was diluted
10 with water. The solution was adjusted to pH 7 with 1N hydrochloric acid. The formed precipitate was collected by vacuum filtration to give ethyl 3-[[1-(hydroxyimino)ethyl]-amino]pyrazole-4-carboxylate (1.52 g).

15 NMR (DMSO-d₆, δ) : 1.27 (3H, t, J=8Hz), 2.20 (3H, s), 4.22 (2H, q, J=8Hz), 8.16 (1H, s), 8.07 (1H, s), 9.82 (1H, s)

Preparation 102

To a solution of ethyl 3-[[1-(hydroxyimino)ethyl]amino]-
20 pyrazole-4-carboxylate (1.50 g) and pyridine (1.12 g) in N,N-dimethylformamide (20 ml) was added dropwise p-toluenesulfonyl chloride (1.39 g) under ice bath cooling and stirred at ambient temperature for 3 hours. The reaction mixture was diluted with ethyl acetate and water. The
25 organic layer was separated and washed with saturated sodium hydrogen carbonate aqueous solution and brine. The solution was dried over magnesium sulfate and concentrated in vacuo to give ethyl 3-[1-(p-toluenesulfonyloxyimino)ethyl]amino-1H-pyrazole-4-carboxylate (2.14 g).

30 NMR (CDCl₃, δ) : 1.42 (3H, t, J=8Hz), 2.30 (3H, s), 2.43 (3H, s), 4.38 (2H, q, J=8Hz), 7.33 (2H, d, J=8Hz), 7.90-7.93 (3H, m), 9.10 (1H, s)

Preparation 103

35 A solution of ethyl 3-[1-(p-toluenesulfonyloxyimino)-

ethylamino-1H-pyrazole-4-carboxylate (1.00 g) and pyridine (240 mg) in ethanol (20 ml) was refluxed for 3 hours. The reaction mixture was concentrated in vacuo and the residue was diluted with water. The solution was adjusted to pH 4 with 1N hydrochloric acid. The formed precipitate was collected by vacuum filtration to give ethyl 2-methyl-1H-pyrazolo[1,5-b][1,2,4]triazole-7-carboxylate (240 mg).

NMR (DMSO-d₆, δ) : 1.29 (3H, t, J=8Hz), 2.45 (3H, s), 4.23 (2H, q, J=8Hz), 7.85 (1H, s)

Preparation 104

To a solution of ethyl 3-aminophthalic acid (1.00 g) in ethanol (10 ml) was added formamidine hydrochloride (444 mg) and refluxed for 12 hours. After cooling, the formed precipitate was collected by vacuum filtration and the precipitate was washed with ethanol to give 4-hydroxyquinazoline-5-carboxylic acid (504 mg).

NMR (DMSO-d₆, δ) : 7.52-7.62 (2H, m), 7.72 (1H, t, J=8Hz), 8.13 (1H, s)

Preparation 105

To a solution of ethyl 3-nitro-2-aminobenzoate (700 mg) in N,N-dimethylaniline (7 ml) was added dropwise isonicotinoyl chloride (2.96 g) at 120°C for 3 hours. After cooling, the reaction mixture was diluted with ethyl acetate and saturated sodium hydrogen carbonate aqueous solution. The organic layer was separated and washed with brine. The solution was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by silica gel column chromatography (n-hexane:ethyl acetate = 9:1) to give ethyl 3-nitro-2-(4-pyridyl)carbonylaminobenzoate (200 mg).

NMR (CDCl₃, δ) : 1.42 (3H, t, J=7Hz), 4.43 (2H, q, J=7Hz), 7.26 (1H, s), 7.40 (1H, t, J=7Hz), 7.82-7.84 (2H, m), 8.15 (1H, d, J=8Hz), 8.31 (1H, d, J=8Hz), 8.85 (1H, d, J=7Hz)

Preparation 106

To a solution of ethyl 3-nitro-2-(4-pyridyl)-
carbonylaminobenzoate (200 mg) in ethanol (2 ml) was added
iron (177 mg) and acetic acid (381 mg) at ambient temperature
5 and the reaction mixture was stirred at 60°C for 2 hours.
After the reaction mixture was filtered through a bed of
celite, the filtrate was concentrated in vacuo. The residue
was diluted with chloroform and saturated sodium hydrogen
carbonate aqueous solution. The organic layer was separated
10 and washed with brine. The solution was dried over magnesium
sulfate and concentrated in vacuo to give ethyl 2-(4-
pyridyl)-1H-benzimidazole-4-carboxylate (150 mg).

NMR (CDCl₃, δ) : 1.48 (3H, t, J=7Hz), 4.40 (2H, q,
J=7Hz), 7.38 (1H, t, J=8Hz), 7.90-8.00 (3H, m),
15 8.05 (1H, d, J=8Hz), 8.82 (1H, d, J=6Hz)

Preparation 107

To a solution of ethyl 3-amino-2-hydroxybenzoate (500
mg), sodium hydrogen carbonate (927 mg) and
20 benzyltributylammonium bromide (983 mg) in chloroform (10 ml)
was added dropwise chloroacetyl chloride (374 mg) in
chloroform (3 ml) under ice bath cooling. The reaction
mixture was stirred at ambient temperature for 1 hour and at
60°C for 2 hours. To the reaction mixture was added dropwise
25 chloroacetyl chloride (312 mg) in chloroform (3 ml) under ice
bath cooling and stirred at 60°C for 2 hours. After the
reaction mixture was concentrated in vacuo, the residue was
diluted with chloroform and saturated sodium hydrogen
carbonate aqueous solution. The organic layer was separated
30 and washed with brine. The solution was dried over magnesium
sulfate and concentrated in vacuo to give ethyl 2H-1,4-
benzoxazin-3-one-8-carboxylate (460 mg).

NMR (CDCl₃, δ) : 1.38 (3H, t, J=7Hz), 4.36 (2H, q,
J=8Hz), 4.71 (2H, s), 6.96-7.00 (2H, m), 7.47-7.52
35 (1H, m), 9.04 (1H, s)

Example 40

The following compounds were obtained according to a similar manner to that of Example 1.

- 5 1) 4-(Imidazo[1,5-a]pyridine-1-carbonyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
- NMR (CDCl₃, δ) : 1.48-1.58 (2H, m), 1.65-1.88 (4H, m),
2.27 (3H, s), 2.29 (3H, s), 2.32-2.40 (6H, m), 3.33
10 (3H, s), 3.46-3.51 (2H, m), 3.59-3.67 (2H, m), 3.80
(3H, s), 3.81-3.99 (2H, m), 6.58 (1H, d, J=8Hz),
6.62 (1H, s), 6.74-6.87 (2H, m), 6.93 (1H, d,
J=8Hz), 7.00 (1H, s), 7.02-7.09 (1H, m), 8.01 (1H,
d, J=8Hz), 8.03 (1H, s), 8.29 (1H, d, J=9Hz), 8.36
15 (1H, d, J=8Hz), 9.60 (1H, s)

- 2) 4-[(1-tert-Butoxycarbonyl-2-ethoxycarbonylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
20 benzamide
- NMR (CDCl₃, δ) : 1.28 (3H, t, J=8Hz), 1.52 (9H, s),
1.55-1.88 (6H, m), 2.28 (3H, s), 2.31 (3H, s),
2.32-2.43 (6H, m), 3.32 (3H, s), 3.42-3.53 (3H, m),
3.60-3.67 (2H, m), 3.78 (3H, s), 3.80-4.00 (3H, m),
25 4.19 (2H, q, J=8Hz), 4.82-4.91 (1H, m), 6.59 (1H,
d, J=8Hz), 6.62 (1H, s), 6.84 (1H, d, J=8Hz), 6.92
(1H, d, J=8Hz), 7.02 (1H, s), 7.20 (1H, d, J=8Hz),
7.26-7.32 (1H, m), 8.02-8.10 (1H, m), 8.20 (1H, d,
J=8Hz), 8.42 (1H, s)

- 30 3) 4-[(1-tert-Butoxycarbonylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
- NMR (CDCl₃, δ) : 1.48-1.60 (2H, m), 1.56 (9H, s),
35 1.66-1.88 (4H, m), 2.28 (3H, s), 2.30 (3H, s),

176

2.31-2.42 (6H, m), 3.31 (3H, s), 3.40 (2H, t, J=9Hz), 3.47-3.51 (2H, m), 3.59-3.67 (2H, m), 3.78 (3H, s), 3.84-4.04 (4H, m), 6.58 (1H, d, J=8Hz), 6.62 (1H, s), 6.84 (1H, d, J=8Hz), 6.92 (1H, d, J=8Hz), 7.02 (1H, s), 7.20 (1H, t, J=8Hz), 7.23-7.29 (1H, m), 8.01 (1H, s), 8.23 (1H, d, J=8Hz), 8.38 (1H, s)

5

- 4) 4-[(2-Benzyloxymethyl-1-tert-butoxycarbonylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

10

NMR (CDCl₃, δ) : 1.50 (9H, s), 1.51-1.58 (2H, m), 1.66-1.87 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.41 (6H, m), 3.32 (3H, s), 3.41-3.53 (5H, m), 3.59-3.67 (3H, m), 3.76 (3H, s), 3.83-3.99 (2H, m), 4.48 (2H, s), 4.57-4.65 (1H, br), 6.58 (1H, d, J=8Hz), 6.63 (1H, s), 6.84 (1H, d, J=8Hz), 6.92 (1H, d, J=8Hz), 7.02 (1H, s), 7.19-7.30 (8H, m), 8.26 (1H, d, J=8Hz), 8.38 (1H, s)

15

20

- 5) 4-[(1-tert-Butoxycarbonyl-3-tert-butylidiphenylsilyloxy-methylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

25

NMR (CDCl₃, δ) : 0.90 (9H, s), 1.50-1.60 (2H, m), 1.67 (9H, s), 1.68-1.89 (4H, m), 2.22 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.33 (3H, s), 3.46-3.51 (2H, m), 3.60-3.67 (5H, m), 3.87-3.98 (2H, m), 4.88 (2H, s), 6.53 (1H, d, J=8Hz), 6.62 (1H, s), 6.83 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 6.98 (1H, s), 7.21-7.41 (8H, m), 7.50-7.59 (5H, m), 8.16 (1H, s), 8.26 (1H, d, J=8Hz), 8.34 (1H, d, J=8Hz)

30

35

- 6) 4-[(1-tert-Butoxycarbonyl-2-tert-butylidiphenylsilyloxy-

methylinol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

5 NMR (CDCl₃, δ) : 0.90 (9H, s), 1.50-1.60 (2H, m), 1.67 (9H, s), 1.68-1.89 (4H, m), 2.22 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.33 (3H, s), 3.46-3.51 (2H, m), 3.60-3.67 (5H, m), 3.87-3.98 (2H, m), 4.88 (2H, s), 6.53 (1H, d, J=8Hz), 6.62 (1H, s), 6.83 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 6.98 (1H, s), 7.21-
10 7.41 (8H, m), 7.50-7.59 (5H, m), 8.16 (1H, s), 8.26 (1H, d, J=8Hz), 8.34 (1H, d, J=8Hz)

7) 4-[(1-tert-Butoxycarbonyl-2-phthalimidomethylinol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

15 NMR (CDCl₃, δ) : 1.46-1.57 (2H, m), 1.64-1.85 (4H, m), 1.73 (9H, s), 2.25 (3H, s), 2.30 (3H, s), 2.32-2.41 (6H, m), 3.30 (3H, s), 3.44-3.51 (2H, m), 3.60-3.67 (2H, m), 3.72 (3H, s), 3.81-3.96 (2H, m), 5.29 (2H, s), 6.54 (1H, d, J=8Hz), 6.59 (1H, s), 6.78 (1H, s), 6.83 (1H, d, J=8Hz), 6.98 (1H, s), 7.32 (1H, t, J=8Hz), 7.54 (1H, d, J=8Hz), 7.77-7.81 (2H, m), 7.88-7.93 (2H, m), 8.03 (1H, s), 8.18 (1H, d, J=8Hz),
20 8.31 (1H, d, J=8Hz), 8.42 (1H, s)

8) 4-[(1-tert-Butoxycarbonyl-2-methylinol-4-yl)carbonyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

30 NMR (CDCl₃, δ) : 1.48-1.60 (2H, m), 1.65-1.90 (4H, m), 1.69 (9H, s), 2.28 (3H, s), 2.29 (3H, s), 2.32-2.43 (6H, m), 2.62 (3H, s), 3.33 (3H, s), 3.47-3.53 (2H, m), 3.60-3.68 (2H, m), 3.78 (3H, s), 3.86-4.00 (2H, m), 6.59 (1H, d, J=8Hz), 6.63 (1H, s), 6.83-6.98 (3H, m), 7.05 (1H, s), 7.24-7.31 (1H, m), 7.57 (1H,
35

d, J=8Hz), 8.28-8.37 (2H, m), 8.53 (1H, s)

- 9) 4-[(1-tert-Butoxycarbonylindolin-6-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.50-1.63 (2H, m), 1.58 (9H, s), 1.67-1.88 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.12 (2H, t, J=8Hz), 3.32 (3H, s), 3.46-3.51 (2H, m), 3.60-3.67 (2H, m), 3.77 (3H, s), 3.85-3.99 (2H, m), 4.03 (2H, t, J=8Hz), 6.58 (1H, d, J=8Hz), 6.63 (1H, s), 6.84 (1H, d, J=8Hz), 6.90 (1H, d, J=8Hz), 7.00 (1H, s), 7.21 (1H, d, J=8Hz), 7.45 (1H, d, J=8Hz), 8.24 (1H, d, J=8Hz), 8.47 (1H, s)

- 10) 4-[(1-tert-Butoxycarbonylindol-6-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.48-1.59 (2H, m), 1.67-1.90 (4H, m), 1.70 (9H, s), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.33 (3H, s), 3.47-3.52 (2H, m), 3.60-3.67 (2H, m), 3.79 (3H, s), 3.83-4.00 (2H, m), 6.58 (1H, d, J=8Hz), 6.61 (1H, d, J=3Hz), 6.63 (1H, s), 6.86 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 7.03 (1H, s), 7.62 (1H, d, J=9Hz), 7.72 (1H, d, J=9Hz), 7.75 (1H, d, J=3Hz), 8.30 (1H, d, J=8Hz), 8.60 (1H, s), 8.67 (1H, s)

- 11) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(quinolin-8-yl)-carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.48-1.59 (2H, m), 1.65-1.76 (2H, m), 1.78-1.89 (2H, m), 2.24 (3H, s), 2.26 (3H, s), 2.31-2.41 (6H, m), 3.32 (3H, s), 3.43-3.50 (2H, m), 3.58-3.66 (2H, m), 3.83-3.99 (2H, m), 3.88 (3H, s),

179

6.54-6.64 (2H, m), 6.87 (1H, d, J=8Hz), 6.97 (1H, d, J=8Hz), 7.03 (1H, s), 7.50 (1H, dd, J=8, 7Hz), 7.70 (1H, t, J=8Hz), 7.87 (1H, d, J=8Hz), 8.31 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz), 8.88 (1H, d, J=7Hz), 8.98 (1H, m)

5

- 12) 4-(3-Hydroxy-1H-indazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

10 NMR (CDCl₃, δ) : 1.47-1.58 (2H, m), 1.66-1.78 (2H, m), 1.78-1.89 (2H, m), 2.27 (3H, s), 2.37 (3H, s), 2.43-2.54 (6H, m), 3.33 (3H, s), 3.52-3.57 (2H, m), 3.64-3.72 (2H, m), 3.79 (3H, s), 3.85-4.00 (2H, m), 6.54-6.65 (2H, m), 6.80-7.02 (3H, m), 7.34-7.49
15 (2H, m), 7.58 (1H, d, J=8Hz), 8.26 (1H, d, J=8Hz), 9.32 (1H, br)

- 13) 3-Methoxy-N-methyl-N-(4-methyl-2-benzyloxyphenyl)-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide

20 NMR (CDCl₃, δ) : 1.48 (9H, s), 2.23 (3H, s), 3.39 (3H, s), 3.61 (3H, s), 4.52 (2H, d, J=7Hz), 4.87 (1H, d, J=12Hz), 5.03 (1H, d, J=12Hz), 5.60 (1H, br), 6.60-6.70 (2H, m), 6.85-7.00 (3H, m), 7.21-7.40 (6H, m),
25 7.48 (1H, d, J=8Hz), 8.10 (1H, d, J=8Hz), 8.47 (1H, d, J=8Hz)

- 14) N-(2,4-Dimethylphenyl)-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonylaminomethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide

30 NMR (CDCl₃, δ) : 1.50 (9H, s), 2.14 (3H, s), 2.25 (3H, s), 3.37 (3H, s), 3.74 (3H, s), 4.57 (2H, d, J=7Hz), 5.64 (1H, br), 6.86-6.99 (7H, m), 7.31 (1H, t, J=8Hz), 7.51 (1H, br), 8.10 (1H, br), 8.47 (1H, br)
35

- 15) 3-Methoxy-N-(2-methoxy-4-methylphenyl)-N-methyl-4-[2-(tert-butoxycarbonylaminomethyl)-1H-benzimidazol-4-yl]-carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.49 (9H, s), 2.28 (3H, s), 3.34 (3H, s), 3.71 (3H, s), 3.77 (3H, s), 4.57 (2H, d, J=7Hz), 5.69 (1H, br), 6.56-6.63 (2H, m), 6.86-6.98 (3H, m), 7.29 (1H, t, J=8Hz), 7.50 (1H, d, J=8Hz), 8.09 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz)

- 16) 3-Methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonylphenylmethoxy]phenyl]-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide

MASS (m/z) : 776

- 17) 3-Methoxy-N-methyl-N-[4-methyl-2-[3-(4-methylpiperazin-1-yl)carbonylprop-1-yloxy]phenyl]-4-[2-(tert-butoxycarbonylaminomethyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.50 (9H, s), 2.06-2.17 (2H, m), 2.26 (3H, s), 2.31-2.39 (4H, m), 2.50 (2H, t, J=7.5Hz), 3.33 (3H, s), 3.43-3.50 (2H, m), 3.52-3.70 (2H, m), 3.81 (3H, s), 3.85-4.06 (2H, m), 4.58 (2H, m), 6.60-6.68 (2H, m), 6.89-7.05 (3H, m), 7.33 (1H, t, J=8Hz), 7.51 (1H, d, J=8Hz), 8.12 (1H, d, J=8Hz), 8.51 (1H, d, J=8Hz)

- 18) 3-Methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonylbut-1-yloxy]phenyl]-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.51 (9H, s), 1.64-1.73 (2H, m), 1.76-1.88 (2H, m), 2.26 (6H, s), 2.28 (3H, s), 2.32-2.47 (6H, m), 3.33 (3H, s), 3.43-3.51 (2H, m), 3.58-3.68 (2H, m), 3.76-4.00 (5H, m), 4.60 (2H, m), 5.83 (1H,

181

br), 6.41 (1H, d, J=8Hz), 6.54-6.64 (2H, m), 6.78-7.03 (3H, m), 7.42 (1H, d, J=8Hz), 8.10 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz)

- 5 19) N-[2-(4-Ethoxycarbonylpent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(3-nitro-2-phthalimidomethylcarbonylaminophenyl)carbonylaminobenzamide

10 NMR (CDCl₃, δ) : 1.24 (3H, t, J=7.5Hz), 1.44-1.58 (2H, m), 1.64-1.77 (2H, m), 1.77-1.90 (2H, m), 2.28 (3H, s), 2.33 (2H, t, J=7.5Hz), 3.34 (3H, s), 3.72 (3H, s), 3.82-4.00 (2H, m), 4.11 (2H, q, J=7.5Hz), 4.49 (2H, s), 6.60-6.66 (2H, m), 6.88 (1H, d, J=8Hz), 6.98 (1H, s), 7.38 (1H, t, J=8Hz), 7.72-7.90 (6H, m), 8.02 (1H, d, J=8Hz), 8.40 (1H, s)

15

- 20) 4-[2-Carbamoyl-1-(4-methoxybenzyl)-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

20 NMR (CDCl₃, δ) : 1.47-1.63 (2H, m), 1.63-1.77 (2H, m), 1.77-1.91 (2H, m), 2.24 (3H, s), 2.29 (3H, s), 2.31-2.42 (6H, m), 3.33 (3H, s), 3.43-3.53 (2H, m), 3.58-3.68 (2H, m), 3.75 (3H, s), 3.80-3.90 (4H, m), 3.90-4.00 (1H, m), 5.98 (2H, s), 6.02 (1H, br s), 6.55-6.65 (2H, m), 6.81 (2H, d, J=8Hz), 6.88 (1H, d, J=8Hz), 6.98 (1H, d, J=8Hz), 7.10 (1H, s), 7.21 (2H, d, J=8Hz), 7.48 (1H, t, J=8Hz), 7.62 (1H, d, J=8Hz), 7.84 (1H, s), 8.24 (1H, d, J=8Hz), 8.51 (1H, d, J=8Hz)

30

- 21) 4-[2-(N,N-Dimethylcarbamoyl)-1-(4-methoxybenzyl)-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

35 NMR (CDCl₃, δ) : 1.49-1.64 (2H, m), 1.64-1.78 (2H, m),

182

1.78-1.90 (2H, m), 2.27 (3H, s), 2.30 (3H, s),
2.33-2.44 (6H, m), 3.13 (3H, s), 3.19 (3H, s), 3.34
(3H, s), 3.46-3.54 (2H, m), 3.60-3.69 (2H, m),
3.74-3.81 (6H, m), 3.81-4.01 (2H, m), 5.55 (2H, s),
5 6.58 (1H, d, J=8Hz), 6.64 (1H, s), 6.80-6.90 (3H,
m), 6.84 (1H, d, J=8Hz), 7.06 (1H, s), 7.16 (2H, d,
J=8Hz), 7.44 (1H, t, J=8Hz), 7.56 (1H, d, J=8Hz),
8.24 (1H, d, J=8Hz), 8.47 (1H, d, J=8Hz)

10 22) 4-[2-[1-(Benzyloxycarbonyl)-4-piperidyl]-1H-
benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.37-1.49 (2H, m), 1.49-1.61 (2H,
15 m), 1.67-1.79 (2H, m), 1.79-1.92 (2H, m), 2.07-2.35
(14H, m), 3.07 (2H, br peak), 3.15-3.29 (4H, m),
3.29-3.46 (4H, m), 3.68 (3H, s), 3.84 (1H, br
peak), 3.94 (1H, br peak), 4.11-4.22 (2H, m), 5.11
(2H, s), 6.63 (1H, d, J=8Hz), 6.81 (1H, s), 6.88-
20 6.97 (2H, m), 7.04 (1H, d, J=8Hz), 7.26-7.45 (6H,
m), 7.70 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.43
(1H, d, J=8Hz)

23) 4-[2-(N-tert-Butoxycarbonylaminomethyl)-1-methyl-1H-
25 benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.44-1.56 (11H, m), 1.64-1.75 (2H, m),
1.74-1.88 (2H, m), 2.26 (6H, s), 2.30-2.40 (6H, m),
30 3.33 (3H, s), 3.43-3.50 (2H, m), 3.58-3.66 (2H, m),
3.81 (3H, s), 3.83-4.00 (5H, m), 4.68 (1H, d,
J=5Hz), 5.90 (1H, br peak), 6.54-6.64 (2H, m), 6.88
(1H, d, J=8Hz), 6.95 (1H, d, J=8Hz), 7.09 (1H, s),
7.41 (1H, t, J=8Hz), 7.51 (1H, d, J=8Hz), 8.18 (1H,
35 d, J=8Hz), 8.53 (1H, d, J=8Hz)

24) 4-[2-(N-tert-Butoxycarbonylaminomethyl)-3-methyl-3H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

5 NMR (CDCl₃, δ) : 1.40-1.61 (2H, m), 1.61-1.91 (4H, m),
2.22-2.32 (6H, m), 2.32-2.44 (6H, m), 3.33 (3H, s),
3.43-3.55 (2H, m), 3.58-3.69 (2H, m), 3.74 (3H, s),
3.82 (3H, s), 3.87-4.03 (2H, m), 4.62 (2H, d,
J=5Hz), 5.55 (1H, br peak), 6.60 (1H, d, J=8Hz),
10 6.66 (1H, s), 6.80-6.90 (1H, m), 6.96 (1H, d,
J=8Hz), 7.04 (1H, s), 7.23-7.32 (1H, m), 7.43 (1H,
d, J=8Hz), 7.84 (1H, d, J=8Hz), 8.28 (1H, d,
J=8Hz), 8.34 (1H, s)

15 25) 4-(2-Methylthio-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO₆, δ) : 1.35-1.49 (2H, m), 1.49-1.62 (2H, m),
1.67-1.80 (2H, m), 2.12 (3H, s), 2.15-2.34 (9H, m),
20 2.85 (3H, s), 3.19 (3H, s), 3.37-3.46 (4H, m), 3.73
(3H, s), 3.89 (1H, br peak), 3.96 (1H, br peak),
6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.89-6.99 (2H,
m), 7.03 (1H, d, J=8Hz), 7.28 (1H, t, J=8Hz), 7.61
(1H, d, J=8Hz), 7.87 (1H, d, J=8Hz), 8.39 (1H, d,
25 J=8Hz)

26) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-methylsulfonyl-1H-benzimidazol-4-yl)carbonylaminobenzamide

30 NMR (CDCl₃, δ) : 1.46-1.63 (2H, m), 1.63-1.78 (2H, m),
1.78-1.90 (2H, m), 2.24 (3H, s), 2.29 (3H, s),
2.33-2.44 (6H, m), 3.33 (3H, s), 3.41 (3H, s),
3.44-3.53 (2H, m), 3.60-3.69 (2H, m), 3.76-3.90
(4H, m), 3.90-4.02 (1H, m), 6.53-6.64 (2H, m), 6.88
35 (1H, d, J=8Hz), 6.92-7.02 (2H, m), 7.49 (1H, t,

J=8Hz), 7.77 (1H, d, J=8Hz), 8.09 (1H, d, J=8Hz),
8.39 (1H, d, J=8Hz)

- 27) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-sulfamoyl-1H-
benzimidazol-4-yl)carbonylaminobenzamide

NMR (DMSO-d₆, δ) : 1.36-1.50 (2H, m), 1.50-1.64 (2H,
m), 1.68-1.81 (2H, m), 2.19 (3H, s), 2.23 (3H, s),
2.34-2.38 (6H, m), 3.19 (3H, s), 3.39-3.50 (4H, m),
3.76 (3H, s), 3.87 (1H, br peak), 3.96 (1H, br
peak), 6.64 (1H, d, J=8Hz), 6.83 (1H, s), 6.89 (1H,
s), 6.95 (1H, d, J=8Hz), 7.03 (1H, d, J=8Hz), 7.50
(1H, t, J=8Hz), 7.82 (1H, d, J=8Hz), 7.98-8.10 (3H,
m), 8.36 (1H, d, J=8Hz)

- 28) 4-(2,4-Dihydroxyquinazolin-8-yl)carbonylamino-3-methoxy-
N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.38-1.52 (2H, m), 1.52-1.63 (2H,
m), 1.70-1.80 (2H, m), 2.22 (3H, s), 2.23 (3H, s),
2.32-2.43 (6H, m), 3.18 (3H, s), 3.40-3.46 (4H, m),
3.63 (3H, s), 3.80-4.00 (2H, m), 6.65 (1H, d,
J=8Hz), 6.82 (1H, s), 6.90-6.93 (2H, m), 7.05 (1H,
d, J=8Hz), 7.30 (1H, t, J=8Hz), 7.53 (1H, d,
J=8Hz), 8.12 (1H, d, J=8Hz), 8.23 (1H, d, J=8Hz)

- 29) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-methyl-1H-
pyrazolo[1,5-b][1,2,4]triazol-7-yl)carbonylamino-
benzamide

NMR (DMSO-d₆, δ) : 1.40-1.50 (2H, m), 1.50-1.62 (2H,
m), 1.70-1.80 (2H, m), 2.23 (6H, s), 2.29-2.37 (6H,
m), 2.39 (3H, s), 3.19 (3H, s), 3.43-3.47 (4H, m),
3.64 (3H, s), 3.43-3.47 (4H, m), 3.64 (3H, s),
3.80-4.00 (4H, m), 6.63 (1H, d, J=8Hz), 6.78-6.90

185

(3H, m), 7.01 (1H, d, J=8Hz), 7.73 (1H, d, J=8Hz),
8.05 (1H, s), 8.88 (1H, s)

- 5 30) 4-(4-Hydroxyquinazolin-5-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.40-1.55 (2H, m), 1.65-1.80 (4H, m),
2.27 (3H, s), 2.30 (3H, s), 2.33-2.43 (6H, m), 3.33
(3H, s), 3.50 (2H, t, J=7Hz), 3.60-3.67 (5H, m),
10 3.77-3.97 (2H, m), 6.60-6.65 (2H, m), 6.90-6.95
(2H, m), 7.02 (1H, s), 7.50-7.53 (1H, m), 7.77-7.80
(1H, m), 7.92 (1H, s), 7.98 (1H, s), 8.33 (1H, d,
J=8Hz)

- 15 31) 4-(2-Dimethylaminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.50-1.60 (2H, m), 1.60-1.78 (2H, m),
1.78-1.90 (2H, m), 2.25 (3H, s), 2.27 (3H, s),
20 2.33-2.40 (12H, m), 3.33 (3H, s), 3.48 (2H, t,
J=7Hz), 3.62 (2H, t, J=7Hz), 3.75-4.00 (7H, m),
6.55-6.62 (2H, m), 6.85 (1H, d, J=8Hz), 6.94 (1H,
d, J=8Hz), 7.01 (1H, s), 7.33 (1H, t, J=8Hz), 7.57
(1H, d, J=8Hz), 8.15 (1H, d, J=8Hz), 8.50 (1H, d,
25 J=8Hz)

- 32) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-methylpiperazin-1-yl)methyl-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.50-1.60 (2H, m), 1.67-1.78 (2H, m),
1.78-1.86 (2H, m), 2.26-2.38 (12H, m), 2.48 (4H, br
s), 2.62 (4H, br s), 3.32 (3H, s), 3.47 (2H, t,
J=7Hz), 3.62 (2H, t, J=7Hz), 3.80-4.00 (7H, m),
30 6.54-6.63 (2H, m), 6.86 (1H, d, J=8Hz), 6.93-7.03
35

(2H, m), 7.28-7.37 (1H, m), 7.58 (1H, d, J=8Hz),
8.15 (1H, d, J=8Hz), 8.52 (1H, d, J=8Hz)

33) 4-[2-(4-Dimethylaminopiperidino)methyl-1H-benzimidazol-
4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-
(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide

NMR (CDCl₃, δ) : 1.50-1.93 (8H, m), 2.13-2.40 (17H, m),
2.92-3.00 (2H, m), 3.32 (3H, s), 3.48 (2H, t,
J=7Hz), 3.62 (2H, t, J=7Hz), 3.82-4.00 (7H, m),
6.54-6.62 (2H, m), 6.86 (1H, d, J=8Hz), 6.93-7.05
(2H, m), 7.35 (1H, t, J=8Hz), 7.59 (1H, d, J=8Hz),
8.15 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz)

34) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-morpholinomethyl-
1H-benzimidazol-4-yl)carbonylamino benzamide

NMR (CDCl₃, δ) : 1.50-1.60 (2H, m), 1.66-1.77 (2H, m),
1.78-1.90 (2H, m), 2.25 (3H, s), 2.28 (3H, s),
2.35-2.40 (6H, m), 2.53-2.62 (4H, m), 3.33 (3H, s),
3.50 (2H, t, J=7Hz), 3.63 (2H, t, J=7Hz), 3.73-3.77
(4H, m), 3.81-4.01 (7H, m), 6.56-6.63 (2H, m), 6.87
(1H, d, J=8Hz), 6.96-7.07 (2H, m), 7.28-7.38 (1H,
m), 7.50-7.60 (1H, m), 8.17 (1H, d, J=8Hz), 8.52
(1H, d, J=8Hz)

Example 41

The following compound was obtained according to a
similar manner to that of Example 4.

1) N-[2-[4,4-Dimethyl(2,5-oxazolinyl)]phenyl]-3-methoxy-N-
methyl-4-(3-nitro-2-trifluoroacetylaminobenzoyl)-
aminobenzamide

NMR (CDCl₃, δ) : 1.36 (3H, s), 1.37 (3H, s), 3.39 (3H,
s), 3.63 (3H, s), 4.10 (2H, s), 6.71-7.60 (6H, m),

7.77 (1H, d, J=8Hz), 7.95 (1H, d, J=8Hz), 8.09 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz)

2) 3-Methoxy-N-methyl-N-[2-(morpholin-4-yl)phenyl]-4-(3-nitro-2-trifluoroacetylaminobenzoyl)aminobenzamide
NMR (CDCl₃, δ) : 2.18-2.63 (2H, m), 2.74-2.93 (2H, m), 3.48 (3H, s), 3.54-3.82 (7H, m), 6.87-7.40 (7H, m), 7.91-8.24 (3H, m)

3) 3-Methoxy-N-methyl-N-[2-(4-methyl-1-piperazinyl)phenyl]-4-(3-nitro-2-trifluoroacetylaminobenzoyl)aminobenzamide
NMR (CDCl₃, δ) : 2.30-3.00 (11H, m), 3.46 (3H, s), 3.59 (3H, s), 6.85-6.96 (2H, m), 7.04 (1H, d, J=8Hz), 7.11-7.24 (2H, m), 7.25-7.39 (2H, m), 7.94 (1H, d, J=8Hz), 8.15 (1H, d, J=8Hz), 8.24 (1H, d, J=8Hz)

4) 3-Methoxy-N-methyl-4-(3-nitro-2-trifluoroacetylaminobenzoyl)amino-N-(2-piperidinophenyl)benzamide
NMR (CDCl₃, δ) : 1.42-1.74 (6H, m), 2.36-2.65 (2H, m), 2.70-2.88 (2H, m), 3.42-3.76 (6H, m), 6.39-8.24 (10H, m)

Example 42

The following compounds were obtained according to a similar manner to that of Example 7.

1) N-(2-Acetoxy-4-methylphenyl)-4-(2,3-diaminophenyl)-carbonylamino-3-methoxy-N-methylbenzamide
NMR (CDCl₃, δ) : 2.30 (6H, s), 3.35 (3H, s), 3.70 (3H, s), 6.67 (1H, t, J=8Hz), 6.81-7.05 (6H, m), 8.29 (1H, d, J=8Hz), 8.45 (1H, s)

2) 4-(2,3-Diaminophenyl)carbonylamino-3-methoxy-N-(2-methoxycarbonyl-4-methylphenyl)-N-methylbenzamide
NMR (CDCl₃, δ) : 2.31 (3H, s), 3.40 (3H, s), 3.71 (3H,

188

s), 3.85 (3H, s), 6.62 (1H, t, J=8Hz), 6.80 (1H, d, J=8Hz), 6.88 (1H, d, J=8Hz), 6.93 (1H, s), 7.02 (1H, d, J=8Hz), 7.10 (1H, d, J=8Hz), 7.59 (1H, s), 8.19 (1H, d, J=8Hz), 8.42 (1H, s)

5

- 3) 4-(2,3-Diaminophenyl)carbonylamino-3-methoxy-N-methyl-N-[2-(4-phthalimidobut-1-yloxy)-4-methylphenyl]benzamide

Example 43

10 The following compounds were obtained according to a similar manner to that of Example 13.

- 1) N-(2-Acetoxy-4-methylphenyl)-4-(1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methylbenzamide

15 NMR (CDCl₃, δ) : 2.27 (3H, s), 2.30 (3H, s), 3.35 (3H, s), 3.74 (3H, s), 6.86 (1H, s), 6.90-7.03 (3H, m), 7.11 (1H, d, J=8Hz), 7.39 (1H, t, J=8Hz), 7.76 (1H, d, J=8Hz), 7.95 (1H, d, J=8Hz), 8.18 (1H, s), 8.49 (1H, d, J=8Hz)

20

- 2) 4-(1H-Benzimidazol-4-yl)carbonylamino-3-methoxy-N-(2-methoxycarbonyl-4-methylphenyl)-N-methylbenzamide

25 NMR (CDCl₃, δ) : 2.30 (3H, s), 3.41 (3H, s), 3.70 (3H, s), 3.82 (3H, s), 6.82-6.90 (2H, m), 7.13 (1H, d, J=8Hz), 7.24 (1H, d, J=8Hz), 7.33 (1H, t, J=8Hz), 7.57 (1H, s), 7.72 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 8.10 (1H, s), 8.41 (1H, d, J=8Hz)

Example 44

30 The following compounds were obtained according to a similar manner to that of Example 16.

- 1) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-(4-phthalimidobut-1-yloxy)-4-methylphenyl]benzamide

35

189

NMR (CDCl₃, δ) : 1.77-1.92 (4H, m), 2.22 (3H, s), 2.64 (3H, s), 3.31 (3H, s), 3.69-3.80 (5H, m), 3.89 (1H, m), 3.97 (1H, m), 6.53-6.61 (3H, m), 6.85 (1H, d, J=8Hz), 6.90-6.97 (2H, m), 7.25 (1H, t, J=8Hz), 7.59-7.70 (3H, m), 7.78-7.90 (3H, m), 8.40 (1H, d, J=8Hz)

2) 3-Methoxy-N-(2-methoxycarbonyl-4-methylphenyl)-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylaminobenzamide
NMR (CDCl₃, δ) : 2.30 (3H, s), 2.60 (3H, s), 3.41 (3H, s), 3.70 (3H, s), 3.83 (3H, s), 6.81-6.89 (2H, m), 7.13 (1H, d, J=8Hz), 7.22-7.30 (2H, m), 7.51-7.58 (2H, m), 7.90 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

Example 45

The following compounds were obtained according to a similar manner to that of Example 18.

1) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(3-phthalimido-propyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide
NMR (CDCl₃, δ) : 1.47-1.60 (2H, m), 1.68-1.77 (2H, m), 1.78-1.90 (2H, m), 2.26 (3H, s), 2.28 (3H, s), 2.30-2.40 (8H, m), 3.00 (2H, t, J=7Hz), 3.32 (3H, s), 3.48 (2H, t, J=7Hz), 3.63 (2H, t, J=7Hz), 3.80-4.00 (7H, m), 6.55-6.62 (2H, m), 6.87 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.02 (1H, s), 7.32 (1H, t, J=8Hz), 7.60 (1H, d, J=8Hz), 7.66-7.87 (5H, m), 8.10 (1H, d, J=8Hz), 8.49 (1H, d, J=8Hz)

30

2) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-phenyl-1H-benzimidazol-4-yl)carbonylaminobenzamide
NMR (DMSO-d₆, δ) : 1.40-1.50 (2H, m), 1.50-1.63 (2H, m), 1.70-1.90 (2H, m), 2.13 (3H, s), 2.17-2.23 (4H,

35

190

m), 2.22 (3H, s), 2.31 (2H, t, J=7Hz), 3.20 (3H, s), 3.39 (4H, br s), 3.87 (3H, s), 3.85-4.00 (2H, m), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.94 (1H, d, J=8Hz), 7.00 (1H, s), 7.04 (1H, d, J=8Hz), 7.40 (1H, t, J=8Hz), 7.57-7.69 (3H, m), 7.80 (1H, d, J=8Hz), 7.97 (1H, d, J=8Hz), 8.32-8.37 (3H, m), 8.50 (1H, d, J=8Hz)

Example 46

The following compounds were obtained according to a similar manner to that of Example 23.

1) 4-[(2-Ethoxycarbonylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.27 (3H, t, J=8Hz), 1.48-1.59 (2H, m), 1.67-1.89 (4H, m), 2.29 (3H, s), 2.31 (3H, s), 2.36-2.45 (6H, m), 3.32 (3H, s), 3.48-3.53 (2H, m), 3.61-3.72 (3H, m), 3.77 (3H, s), 3.84-4.00 (2H, m), 4.09-4.23 (3H, m), 4.41 (1H, dd, J=7, 9Hz), 4.55-4.60 (1H, br s), 6.59 (1H, d, J=8Hz), 6.62 (1H, s), 6.80-6.88 (2H, m), 6.92 (1H, d, J=8Hz), 6.99-7.05 (2H, m), 7.15 (1H, t, J=8Hz), 8.23 (1H, d, J=8Hz), 8.40 (1H, s)

2) 4-[(Indolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.49-1.59 (2H, m), 1.67-1.88 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.32 (3H, s), 3.36 (2H, t, J=9Hz), 3.47-3.52 (2H, m), 3.56-3.66 (4H, m), 3.78 (3H, s), 3.84-3.99 (2H, m), 6.59 (1H, d, J=8Hz), 6.63 (1H, s), 6.73 (1H, d, J=8Hz), 6.86 (1H, d, J=8Hz), 6.90 (1H, d, J=8Hz), 6.97-7.03 (2H, m), 7.09 (1H, t, J=8Hz), 8.27 (1H,

191

d, J=8Hz), 8.39 (1H, s)

- 3) 4-[(2-Hydroxymethylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

5 NMR (CDCl₃, δ) : 1.47-1.57 (2H, m), 1.65-1.86 (4H, m),
2.28 (3H, s), 2.32 (3H, s), 2.34-2.45 (6H, m),
3.06-3.17 (1H, m), 3.32 (3H, s), 3.41-3.52 (3H, m),
3.57 (1H, dd, J=8, 13Hz), 3.60-3.67 (2H, m), 3.70
10 (1H, dd, J=5, 13Hz), 3.77 (3H, s), 3.83-3.98 (2H,
m), 4.02-4.10 (1H, m), 6.59 (1H, d, J=8Hz), 6.62
(1H, s), 6.74 (1H, d, J=8Hz), 6.87 (1H, d, J=8Hz),
6.92 (1H, d, J=8Hz), 6.98 (1H, d, J=8Hz), 7.01 (1H,
s), 7.10 (1H, t, J=8Hz), 8.25 (1H, d, J=8Hz), 8.39
15 (1H, s)

- 4) 4-[(Indolin-6-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

20 NMR (CDCl₃, δ) : 1.48-1.58 (2H, m), 1.64-1.87 (4H, m),
2.28 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.07
(2H, t, J=8Hz), 3.32 (3H, s), 3.46-3.51 (2H, m),
3.60 (2H, t, J=8Hz), 3.61-3.68 (2H, m), 3.77 (3H,
s), 3.83-3.98 (2H, m), 6.58 (1H, d, J=8Hz), 6.62
25 (1H, s), 6.84 (1H, d, J=8Hz), 6.91 (1H, d, J=8Hz),
6.99-7.16 (3H, m), 8.27 (1H, d, J=8Hz), 8.44 (1H,
s)

- 5) 4-[2-[[2-(Dimethylamino)ethyl]amino]-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

30 NMR (DMSO-d₆, δ) : 1.35-1.50 (2H, m), 1.50-1.66 (2H,
m), 1.66-1.85 (2H, m), 2.14 (3H, s), 2.17-2.39
35 (15H, m), 2.45-2.60 (2H, m), 3.21 (3H, s), 3.27-

192

3.53 (6H, m), 3.80-4.01 (5H, m), 6.46-6.54 (1H, m),
6.60-6.71 (1H, m), 6.72-6.85 (2H, m), 6.90 (1H, d,
J=8Hz), 6.95-7.04 (3H, m), 7.81-7.94 (2H, m)

5 Example 47

The following compounds were obtained according to a
similar manner to that of Example 25.

1) 4-[2-Carbamoyl-1H-benzimidazol-4-yl]carbonylamino-3-
10 methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.44-1.66 (2H, m), 1.66-1.80 (2H, m),
1.80-1.93 (2H, m), 2.26 (3H, s), 2.30 (3H, s),
2.32-2.46 (6H, m), 3.35 (3H, s), 3.45-3.54 (2H, m),
15 3.60-3.71 (2H, m), 3.79-3.92 (4H, m), 3.92-4.03
(1H, m), 6.32 (1H, br peak), 6.56-6.69 (2H, m),
6.90 (1H, d, J=8Hz), 6.94-7.04 (1H, m), 7.10 (1H,
s), 7.48-7.61 (2H, m), 7.73 (1H, d, J=8Hz), 8.28
(1H, d, J=8Hz), 8.47-8.57 (1H, m)

20

2) 4-[2-(N,N-Dimethylcarbamoyl)-1H-benzimidazol-4-yl]-
carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide

25 NMR (CDCl₃, δ) : 1.45-1.64 (2H, m), 1.64-1.78 (2H, m),
1.78-1.92 (2H, m), 2.26 (3H, s), 2.29 (3H, s),
2.31-2.43 (6H, m), 3.25 (3H, s), 3.33 (3H, s),
3.43-3.53 (2H, m), 3.59-3.68 (2H, m), 3.70-4.03
(8H, m), 6.54-6.69 (2H, m), 6.86 (1H, d, J=8Hz),
30 6.93 (1H, d, J=8Hz), 7.09 (1H, s), 7.49 (1H, br
peak), 7.71 (1H, br peak), 8.25 (1H, br peak), 8.34
(1H, d, J=8Hz)

Example 48

35

The following compounds were obtained according to a

similar manner to that of Example 26.

1) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-
[2-(5-ethoxycarbonylpent-1-yloxy)-4-methoxyphenyl]-3-
methoxy-N-methylbenzamide

2) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonyl-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.36-1.50 (2H, m), 1.50-1.64 (2H,
m), 1.70-1.82 (2H, m), 2.15 (3H, s), 2.17-2.40 (9H,
m), 3.21 (3H, s), 3.37-3.47 (4H, m), 3.81-4.04 (7H,
m), 6.64 (1H, d, J=8Hz), 6.81 (1H, s), 6.99-7.14
(4H, m), 7.14-7.24 (1H, m), 7.90 (1H, br peak),
8.09 (1H, br peak)

3) 4-[2-(2-Aminoethyl)-1H-benzimidazol-4-yl]carbonyl-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.36-1.51 (2H, m), 1.51-1.64 (2H,
m), 1.70-1.83 (2H, m), 2.14 (3H, s), 2.17-2.38 (9H,
m), 2.94 (2H, t, J=5Hz), 3.09 (2H, t, J=5Hz), 3.21
(3H, s), 3.24-3.49 (4H, m), 3.84-4.04 (5H, m), 6.64
(1H, d, J=8Hz), 6.81 (1H, s), 7.02 (1H, d, J=8Hz),
7.06-7.20 (4H, m), 7.85-7.94 (1H, m), 8.00-8.10
(1H, m)

Example 49

The following compounds were obtained according to a
similar manner to that of Example 28.

1) 4-(2-Amino-3-nitrobenzoyl)amino-N-[2-[4,4-dimethyl(2,5-
oxazolinyl)]phenyl]-3-methoxy-N-methylbenzamide

NMR (CDCl₃, δ) : 1.35 (6H, s), 3.39 (3H, s), 3.67 (3H,
s), 4.09 (2H, s), 6.68 (1H, dd, J=8, 8Hz), 6.99

194

(1H, s), 7.05 (1H, d, J=8Hz), 7.14 (1H, d, J=8Hz),
7.27 (1H, dd, J=8, 8Hz), 7.37 (1H, dd, J=8, 8Hz),
7.70 (1H, d, J=8Hz), 7.79 (1H, d, J=8Hz), 8.04-8.23
(3H, m), 8.25-8.36 (2H, m)

5

- 2) 4-(2-Amino-3-nitrobenzoyl)amino-3-methoxy-N-methyl-N-[2-(morpholin-4-yl)phenyl]benzamide

NMR (CDCl₃, δ) : 2.45-2.63 (2H, m), 2.80-2.98 (2H, m),
3.49 (3H, s), 3.63-3.86 (7H, m), 6.69 (1H, dd, J=8,
10 8Hz), 6.92 (1H, d, J=8Hz), 7.02 (1H, m), 7.05-7.16
(2H, m), 7.17-7.30 (2H, m), 7.72 (1H, d, J=8Hz),
8.10-8.22 (3H, m), 8.28-8.40 (2H, m)

- 3) 4-(2-Amino-3-nitrobenzoyl)amino-3-methoxy-N-methyl-N-(2-piperidinophenyl)benzamide

NMR (CDCl₃, δ) : 1.43-1.72 (6H, m), 2.42-2.56 (2H, m),
2.73-2.87 (2H, m), 3.50 (3H, s), 3.71 (3H, s), 6.68
(1H, dd, J=8, 8Hz), 6.90 (1H, d, J=8Hz), 6.97-7.07
(3H, m), 7.12-7.22 (2H, m), 7.73 (1H, d, J=8Hz),
20 8.11-8.22 (3H, m), 8.28-8.39 (2H, m)

- 4) 4-(2-Amino-3-nitrobenzoyl)amino-3-methoxy-N-methyl-N-[2-(4-methyl-1-piperazinyl)phenyl]benzamide

NMR (CDCl₃, δ) : 2.40 (3H, s), 2.45-2.73 (6H, m), 2.89-
25 3.04 (2H, m), 3.50 (3H, s), 3.69 (3H, s), 6.68 (1H,
dd, J=8, 8Hz), 6.89-7.01 (2H, m), 7.02-7.12 (2H,
m), 7.15-7.29 (2H, m), 7.72 (1H, d, J=8Hz), 8.09-
8.24 (3H, m), 8.28-8.38 (2H, m)

30 Example 50

The following compound was obtained according to a similar manner to that of Example 29.

4-(2-Carboxyphenylcarbamoyl)-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-

35

phenyl]benzamide

5 NMR (DMSO-d₆, δ) : 1.38-1.65 (4H, m), 1.70-1.85 (2H, m), 2.21 (3H, s), 2.30-2.60 (6H, m), 2.68-2.91 (3H, m), 3.17 (3H, s), 3.20 (3H, s), 3.66 (3H, s), 3.83-4.03 (3H, m), 6.10 (1H, d, J=8Hz), 6.82-7.02 (3H, m), 7.43-7.52 (2H, m), 7.63-7.70 (2H, m), 7.91-8.01 (2H, m), 8.67 (1H, d, J=8Hz)

Example 51

10 The following compound was obtained by using 4-(1H-benzimidazol-4-yl)carbonylamino-N-(2-methoxycarbonyl-4-methylphenyl)-3-methoxy-N-methylbenzamide as a starting compound according to a similar manner to that of Example 29.

15 4-(1H-Benzimidazol-4-yl)carbonylamino-N-(2-carboxy-4-methylphenyl)-3-methoxy-N-methylbenzamide

20 NMR (DMSO-d₆, δ) : 2.29 (3H, s), 3.28 (3H, s), 3.70 (3H, s), 6.80-6.89 (2H, m), 7.28-7.43 (3H, m), 7.51 (1H, s), 7.80 (1H, d, J=8Hz), 7.94 (1H, d, J=8Hz), 8.31 (1H, m), 8.53 (1H, s)

Example 52

The following compound was obtained according to a similar manner to that of Example 51.

25

N-(2-Carboxy-4-methylphenyl)-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylaminobenzamide

30 NMR (DMSO-d₆, δ) : 2.29 (3H, s), 2.65 (3H, s), 3.29 (3H, s), 3.73 (3H, s), 6.80-6.89 (2H, m), 7.28-7.38 (3H, m), 7.51 (1H, s), 7.70 (1H, d, J=8Hz), 7.89 (1H, d, J=8Hz), 8.30 (1H, br)

Example 53

35 The following compounds were obtained according to a similar manner to that of Example 30.

- 1) 4-[(2-Carbamoylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide

5 NMR (CDCl₃, δ) : 1.47-1.56 (2H, m), 1.63-1.87 (4H, m),
2.28 (3H, s), 2.29 (3H, s), 2.31-2.42 (6H, m), 3.33
(3H, s), 3.47-3.51 (2H, m), 3.58-3.65 (2H, m), 3.74
(3H, s), 3.85-4.00 (2H, m), 6.59-6.66 (2H, m), 6.91
(1H, d, J=8Hz), 6.98-7.03 (2H, m), 7.35 (1H, t,
J=8Hz), 7.52 (1H, d, J=8Hz), 7.57-7.63 (2H, m),
10 8.32 (1H, d, J=8Hz), 8.60 (1H, s), 9.88 (1H, s)

- 2) 4-[[2-(N-Methylcarbamoyl)indol-4-yl]carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

15 NMR (CDCl₃, δ) : 1.45-1.56 (2H, m), 1.63-1.86 (4H, m),
2.27 (3H, s), 2.29 (3H, s), 2.30-2.42 (6H, m), 3.04
and 3.06 (Total 3H, s), 3.34 (3H, s), 3.47-3.52
(2H, m), 3.57-3.63 (2H, m), 3.75 (3H, s), 3.83-4.00
(2H, m), 6.58-6.64 (2H, m), 6.90 (1H, d, J=8Hz),
20 6.98-7.03 (2H, m), 7.32 (1H, t, J=8Hz), 7.46 (1H,
s), 7.51 (1H, d, J=8Hz), 7.60 (1H, d, J=8Hz), 8.02
(1H, s), 8.33 (1H, d, J=8Hz), 8.59 (1H, s), 9.76
(1H, s)

- 25 3) 4-[2-(N,N-Dimethylcarbamoyl)phenylcarbamoyl]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

30 NMR (CDCl₃, δ) : 1.49-1.59 (2H, m), 1.67-1.89 (4H, m),
2.27 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 2.93
(3H, s), 3.12 (3H, s), 3.35 (3H, s), 3.47-3.52 (2H,
m), 3.60-3.67 (2H, m), 3.84-3.98 (2H, m), 3.92 (3H,
s), 6.58 (1H, d, J=8Hz), 6.61 (1H, s), 6.83 (1H, d,
J=8Hz), 6.98 (1H, d, J=8Hz), 7.01 (1H, s), 7.10
(1H, t, J=8Hz), 7.20 (1H, d, J=8Hz), 7.40 (1H, t,
35 J=8Hz), 8.01 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)

Example 54

The following compounds were obtained according to a similar manner to that of Example 32.

- 5 1) 4-[(1-tert-Butoxycarbonyl-3-hydroxymethylindol-4-yl)-
carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide

10 NMR (CDCl₃, δ) : 1.49-1.61 (2H, m), 1.66 (9H, s), 1.67-
1.89 (4H, m), 2.29 (3H, s), 2.30 (3H, s), 2.32-2.42
(6H, m), 3.34 (3H, s), 3.46-3.51 (2H, m), 3.60-3.67
(2H, m), 3.76 (3H, s), 3.88-4.01 (2H, m), 4.61 (2H,
s), 6.61 (1H, d, J=8Hz), 6.67 (1H, s), 6.87 (1H, d,
J=8Hz), 6.93 (1H, d, J=8Hz), 7.07 (1H, s), 7.37
15 (1H, t, J=8Hz), 7.44 (1H, d, J=8Hz), 7.68 (1H, s),
8.29 (1H, d, J=8Hz), 8.39-8.47 (2H, m)

- 20 2) 4-(2-Hydroxymethyl-1H-benzimidazol-4-yl)carbonyl-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

25 NMR (DMSO-d₆, δ) : 1.40-1.50 (2H, m), 1.50-1.63 (2H, m),
1.69-1.82 (2H, m), 2.14 (3H, s), 2.17-2.37 (9H, m),
3.21 (3H, s), 3.37-3.48 (4H, m), 3.83-4.03 (5H, m),
4.72 (2H, d, J=5Hz), 5.74 (1H, br peak), 6.64 (1H,
d, J=8Hz), 6.80 (1H, s), 6.99-7.23 (5H, m), 7.90
30 (1H, d, J=8Hz), 8.09 (1H, d, J=8Hz)

Example 55

30 The following compounds were obtained according to
similar manners to those of Examples 4 and 28.

- 1) 4-(2-Amino-3-nitrophenyl)carbonylamino-3-methoxy-N-(2-
methoxycarbonyl-4-methylphenyl)-N-methylbenzamide

35 NMR (CDCl₃, δ) : 2.23 (3H, s), 3.39 (3H, s), 3.75 (3H,
s), 3.89 (3H, s), 6.67 (1H, t, J=8Hz), 6.83 (1H, d,

J=8Hz), 6.97 (1H, s), 7.12 (1H, d, J=8Hz), 7.27 (1H, d, J=8Hz), 7.59 (1H, s), 7.70 (1H, d, J=8Hz), 8.04-8.18 (3H, m), 8.12 (1H, d, J=8Hz)

- 5 2) 4-(2-Amino-3-nitrophenyl)carbonylamino-3-methoxy-N-[2-(4-methoxyphenylmethoxy)-4-methylphenyl]-N-methylbenzamide

NMR (CDCl₃, δ) : 2.28 (3H, s), 3.37 (3H, s), 3.66 (3H, s), 3.81 (3H, s), 4.87 (1H, d, J=12Hz), 5.01 (1H, d, J=12Hz), 6.60-6.73 (3H, m), 6.85-6.99 (4H, m), 7.23-7.31 (3H, m), 7.71 (1H, d, J=8Hz), 8.10-8.19 (3H, m), 8.29-8.34 (2H, m)

10

- 15 3) 4-(2-Amino-3-nitrophenyl)carbonylamino-3-methoxy-N-methyl-N-[2-(5-tert-butoxycarbonylamino-pent-1-yl)oxy-4-methylphenyl]benzamide

NMR (CDCl₃, δ) : 1.41 (9H, s), 1.45-1.60 (4H, m), 1.75-1.84 (2H, m), 2.28 (3H, s), 3.09-3.18 (2H, m), 3.31 (3H, s), 3.78 (3H, s), 3.80-3.97 (2H, m), 4.67 (1H, br), 6.58-6.63 (2H, m), 6.69 (1H, t, J=8Hz), 6.89 (1H, d, J=8Hz), 6.96 (1H, d, J=8Hz), 7.03 (1H, s), 7.71 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz), 8.29-8.37 (3H, m)

20

25 Example 56

The following compounds were obtained according to similar manners to those of Examples 7 and 16.

- 30 1) N-(2-Amino-4-methylphenyl)-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-benzamide

NMR (CDCl₃, δ) : 2.18 (3H, s), 2.76 (3H, s), 3.31 (3H, s), 3.80 (3H, s), 6.37 (1H, d, J=8Hz), 6.53 (1H, s), 6.66 (1H, d, J=8Hz), 7.00-7.08 (2H, m), 7.26 (1H, t, J=8Hz), 7.60 (1H, d, J=8Hz), 7.86 (1H, br), 8.44 (1H, d, J=8Hz)

35

- 2) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-(5-tert-butoxycarbonylamino-pent-1-yl)oxy-4-methyl]phenylbenzamide

NMR (CDCl₃, δ) : 1.42 (9H, s), 1.42-1.60 (4H, m), 1.72-1.85 (2H, m), 2.28 (3H, s), 2.67 (3H, s), 3.08-3.17 (2H, m), 3.36 (3H, s), 3.60-3.97 (2H, m), 3.78 (3H, s), 4.80 (1H, br), 6.57-6.63 (2H, m), 6.80-7.08 (3H, m), 7.30 (1H, m), 7.59 (1H, m), 7.91 (1H, br), 8.45 (1H, m)

- 3) N-[2-[4,4-Dimethyl(2,5-oxazolinyl)]phenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylamino-benzamide

NMR (CDCl₃, δ) : 1.33 (3H, s), 1.35 (3H, s), 1.60 (3H, s), 3.41 (3H, s), 3.68 (3H, s), 4.04-4.14 (2H, m), 6.95 (1H, s), 7.07 (1H, d, J=8Hz), 7.14 (1H, d, J=8Hz), 7.18-7.39 (4H, m), 7.59 (1H, d, J=8Hz), 7.80 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.47 (1H, d, J=8Hz)

- 4) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-(morpholin-4-yl)phenyl]benzamide

NMR (DMSO-d₆, δ) : 2.25-2.44 (2H, m), 2.62 (3H, s), 2.72-2.90 (2H, m), 3.20-3.80 (10H, m), 6.87-6.99 (2H, m), 7.05 (1H, d, J=8Hz), 7.09-7.26 (2H, m), 7.30 (1H, dd, J=8, 8Hz), 7.42 (1H, d, J=8Hz), 7.67 (1H, d, J=8Hz), 7.87 (1H, d, J=8Hz), 8.41 (1H, m)

- 5) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-N-(2-piperidinophenyl)benzamide

NMR (CDCl₃, δ) : 1.41-1.72 (6H, m), 2.36-2.53 (2H, m), 2.66 (3H, s), 2.70-2.87 (2H, m), 3.51 (3H, s), 3.71 (3H, s), 6.88 (1H, d, J=8Hz), 6.93-7.33 (7H, m), 7.62 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.42 (1H, d, J=8Hz)

- 6) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-
carbonylamino-N-[2-(4-methyl-1-piperazinyl)phenyl]-
benzamide

5 NMR (CDCl₃, δ) : 2.40 (3H, br s), 2.46-2.70 (9H, m),
2.86-3.01 (2H, m), 3.51 (3H, s), 3.70 (3H, s),
6.82-6.98 (2H, m), 7.02-7.21 (3H, m), 7.22-7.36
(2H, m), 7.43-7.54 (1H, m), 8.11 (1H, d, J=8Hz),
8.58 (1H, d, J=8Hz)

10 Example 57

To a solution of 2-(4-pyridyl)-1H-benzimidazole-4-
carboxylic acid (155 mg) in dichloromethane (2 ml) was added
oxalyl chloride (0.056 ml) and N,N-dimethylformamide (2
drops) and stirred at ambient temperature for 2 hours. The
15 reaction mixture was concentrated in vacuo to give 2-(4-
pyridyl)-1H-benzimidazole-4-carbonyl chloride. To a solution
of 4-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
(76.8 mg) and triethylamine (65 mg) in dichloromethane (5 ml)
20 was added 2-(4-pyridyl)-1H-benzimidazole-4-carbonyl chloride
in dichloromethane (2 ml) under ice bath cooling and stirred
at ambient temperature for 6 hours. After the reaction
mixture was concentrated in vacuo, the residue was diluted
with chloroform and saturated sodium hydrogen carbonate
25 aqueous solution. The organic layer was separated and washed
with brine. The solution was dried over magnesium sulfate
and concentrated in vacuo. The residue was purified by
silica gel (Chromatorex, Fuji Silysia Chemical Ltd.) column
chromatography (methanol:chloroform = 1:49). To the purified
30 product was added water and 1N hydrochloric acid (0.51 ml).
The solution was lyophilized to give 3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-
phenyl]-4-[2-(4-pyridyl)-1H-benzimidazol-4-yl]carbonylamino-
benzamide trihydrochloride (120 mg).

35 NMR (DMSO-d₆, δ) : 1.42-1.52 (2H, m), 1.52-1.62 (2H,

201

m), 1.73-1.83 (2H, m), 2.22 (3H, s), 2.41 (2H, t, J=7Hz), 2.73 (3H, s), 2.82-3.07 (4H, m), 3.21 (3H, s), 3.32-3.52 (3H, m), 3.88 (3H, s), 3.90-4.13 (2H, m), 4.40-4.50 (1H, m), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.98 (1H, d, J=8Hz), 7.03 (1H, s), 7.07 (1H, d, J=8Hz), 7.55 (1H, t, J=8Hz), 7.94 (1H, d, J=8Hz), 8.07 (1H, d, J=8Hz), 8.48 (1H, d, J=8Hz), 8.75 (2H, br s)

10 Example 58

The following compounds were obtained according to a similar manner to that of Example 57.

- 1) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(3-pyridyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.65 (2H, m), 1.70-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7Hz), 2.73 (3H, s), 2.85-3.12 (4H, m), 3.21 (3H, s), 3.35-3.53 (3H, m), 3.88 (3H, s), 3.92-4.13 (2H, m), 4.40-4.45 (1H, m), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.96 (1H, d, J=8Hz), 7.01 (1H, s), 7.06 (1H, d, J=8Hz), 7.48 (1H, t, J=8Hz), 7.89 (1H, d, J=8Hz), 8.03 (1H, d, J=8Hz), 8.13 (1H, t, J=8Hz), 8.48 (1H, d, J=8Hz), 9.00 (1H, br s), 9.12 (1H, d, J=8Hz), 9.67 (1H, br s)

- 2) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(2-pyridyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.63 (2H, m), 1.68-1.82 (2H, m), 2.22 (3H, s), 2.37 (2H, t, J=7Hz), 2.72 (3H, s), 2.82-3.10 (4H, m), 3.20 (3H,

s), 3.33-3.56 (3H, m), 3.85 (3H, s), 3.90-4.10 (2H, m), 4.40-4.45 (1H, m), 6.62 (1H, t, J=8Hz), 6.80-7.06 (4H, m), 7.43 (1H, t, J=8Hz), 7.63 (1H, t, J=7Hz), 7.79 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 8.17 (1H, t, J=8Hz), 8.48 (1H, t, J=8Hz), 8.82 (1H, d, J=5Hz)

3) 4-(2H-1,4-Benzoxazin-3-oxo-8-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.51 (2H, m), 1.51-1.63 (2H, m), 1.70-1.80 (2H, m), 2.22 (3H, s), 2.37 (2H, t, J=7Hz), 2.73 (3H, s), 2.80-3.04 (4H, m), 3.17 (3H, s), 3.36-3.50 (3H, m), 3.73 (3H, s), 3.80-4.15 (2H, m), 4.40-4.47 (1H, m), 4.89 (2H, s), 6.64 (1H, d, J=8Hz), 6.80 (1H, s), 6.88-6.93 (2H, m), 7.03 (1H, d, J=8Hz), 7.10-7.13 (2H, m), 7.59-7.62 (1H, m), 8.22 (1H, d, J=8Hz)

Example 59

The following compound was obtained by using N-(2-phthalimido-4-methylphenyl)-2-amino-3-methoxy-N-methylbenzamide as a starting compound according to a similar manner to that of Example 4.

N-(2-Amino-4-methylphenyl)-4-(2-amino-3-nitrophenyl)carbonylamino-3-methoxy-N-methylbenzamide

NMR (CDCl₃, δ) : 2.20 (3H, s), 3.30 (3H, s), 3.79 (3H, s), 3.89 (2H, br s), 6.39 (1H, d, J=8Hz), 6.52 (1H, s), 6.63-6.71 (2H, m), 7.05 (1H, d, J=8Hz), 7.10 (1H, s), 7.70 (1H, d, J=8Hz), 8.12 (2H, br s), 8.20 (1H, d, J=8Hz), 8.31 (1H, d, J=8Hz), 8.34 (1H, br)

Example 60

A mixture of N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-

methyphenyl]-3-methoxy-N-methyl-4-(3-nitro-2-phthalimidomethylcarbonylaminophenyl)carbonylaminobenzamide (3.96 g), iron powder (1.42 g) and acetic acid (3.05 g) in ethanol (50 ml) was refluxed for 4 hours and the solvent was removed under reduce pressure. The residue was stirred in a mixture of chloroform (100 ml) and saturated aqueous sodium hydrogen carbonate (100 ml) for 30 minutes and the solution was filtered through a bed of celite. The organic phase was separated and washed with brine. The solution was dried over magnesium sulfate and the solvent was evaporated in vacuo. The residue was solidified with diethyl ether to give N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methyphenyl]-3-methoxy-N-methyl-4-(2-phthalimidomethyl-1H-benzimidazol-4-yl)-carbonylaminobenzamide (3.64 g).

NMR (CDCl₃, δ) : 1.24 (3H, t, J=7.5Hz), 1.46-1.57 (2H, m), 1.63-1.75 (2H, m), 1.75-1.88 (2H, m), 2.25 (3H, s), 2.34 (2H, t, J=7.5Hz), 3.31 (3H, s), 3.74-4.00 (2H, m), 4.02 (3H, s), 4.12 (2H, q, J=7.5Hz), 5.21 (2H, s), 6.53-6.63 (2H, m), 6.86 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 7.03 (1H, s), 7.35 (1H, t, J=8Hz), 7.53 (1H, d, J=8Hz), 7.68-7.78 (2H, m), 7.84-7.93 (2H, m), 8.14 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz)

25 Example 61

To a solution of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide (240 mg) in acetonitrile (1 ml) was added cyanoacetic acid (662 mg). The solution was heated at 100°C for 8 hours. After cooling, aqueous sodium hydrogen carbonate was added to the mixture and extracted with ethyl acetate. The extract was washed with brine and dried over sodium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography (7% methanol in chloroform) and preparative

thin-layer chromatography (ethyl acetate:methanol = 1:1) to give 4-[[2-cyanomethyl-1H-benzimidazol-4-yl]carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (80 mg).

5 NMR (CDCl₃, δ) : 1.42-1.94 (6H, m), 2.24 (3H, s), 2.29 (3H, s), 2.32-2.48 (6H, m), 3.36 (3H, s), 3.43-3.55 (2H, m), 3.55-4.21 (9H, m), 6.50-6.68 (2H, m), 6.78 (1H, br), 6.81-7.02 (2H, m), 7.20-7.31 (1H, m), 7.36-7.48 (1H, m), 8.08 (1H, d, J=8Hz), 8.47 (1H, d, J=8Hz)

10

Example 62

A solution of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (100 mg) and mercaptoacetic acid (448 mg) was heated at 80°C for 5 hours. The reaction mixture was diluted with chloroform and washed with aqueous sodium hydrogen carbonate. The extract was dried over sodium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography (10% methanol in chloroform) to give 3-methoxy-4-(2-mercaptomethyl-1H-benzimidazol-4-yl)carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (60 mg).

15

20

25 NMR (CDCl₃, δ) : 1.42-1.90 (6H, m), 2.19-2.47 (12H, m), 3.34 (3H, s), 3.49 (2H, m), 3.57-4.07 (9H, m), 6.51-6.68 (2H, m), 6.81-7.05 (3H, m), 7.31 (1H, dd, J=8, 8Hz), 7.51 (1H, d, J=8Hz), 8.12 (1H, d, J=8Hz), 8.50 (1H, m)

30

Example 63

A solution of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (150 mg), butyrolactone (628 mg) and p-toluenesulfonic acid (139 mg) was heated at 100°C for 4

35

hours. The reaction mixture was diluted with chloroform and washed with aqueous sodium hydrogen carbonate. The extract was dried over sodium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography (10% methanol in chloroform) and then preparative thin-layer chromatography (chloroform/methanol = 20/3) to give 4-[2-(3-hydroxypropyl)-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (70 mg).

NMR (CDCl₃, δ) : 1.44-1.59 (2H, m), 1.63-1.88 (4H, m), 2.01-2.14 (2H, m), 2.25 (3H, s), 2.27 (3H, s), 2.30-2.43 (6H, m), 2.90-3.03 (2H, m), 3.34 (3H, s), 3.42-3.52 (2H, m), 3.56-4.02 (9H, m), 6.52-6.66 (2H, m), 6.78-7.03 (3H, m), 7.24 (1H, dd, J=8, 8Hz), 7.46 (1H, m), 8.07 (1H, m), 8.52 (1H, m)

Example 64

To a mixture of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (140 mg) and sodium carbonate (14 mg) in ethyl acetate (1.5 ml) was added dropwise a solution of 1,1-dichloro-1,1-diphenoxymethane (67 mg) in ethyl acetate (1 ml) in water bath and the mixture was stirred at same temperature for 5 hours. The reaction mixture was evaporated in vacuo and dissolved in chloroform. The organic layer was washed with water, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform:methanol = 10:1) to give 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-phenoxy-1H-benzimidazol-4-yl)carbonylaminobenzamide (18 mg).

NMR (DMSO-d₆, δ) : 1.33-1.48 (2H, m), 1.48-1.62 (2H, m), 1.62-1.78 (2H, m), 2.14 (3H, s), 2.17-2.35 (9H, m), 3.09 (3H, s), 3.17 (3H, s), 3.36-3.45 (4H, m),

3.76-3.87 (1H, m), 3.87-3.99 (1H, m), 6.64 (1H, d, J=8Hz), 6.75 (1H, s), 6.81 (1H, s), 6.88 (1H, d, J=8Hz), 7.00 (1H, d, J=8Hz), 7.25-7.38 (2H, m), 7.45-7.52 (4H, m), 7.60 (1H, d, J=8Hz), 7.88 (1H, d, J=8Hz), 8.25 (1H, d, J=8Hz)

Example 65

A mixture of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (150 mg) and diphenyl N-sulfamoylcarbonimide (85 mg) in dichloromethane (8 ml) was refluxed for 24 hours under nitrogen. The reaction mixture was poured into water and extracted with chloroform. The organic layer was washed with water, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform:methanol = 10:1/chloroform:methanol = 6:1) to give 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-sulfamoylamino-1H-benzimidazol-4-yl)-carbonylaminobenzamide (38 mg).

NMR (DMSO-d₆, δ) : 1.37-1.50 (2H, m), 1.50-1.64 (2H, m), 1.64-1.82 (2H, m), 2.23 (6H, s), 2.27-2.43 (6H, m), 3.19 (3H, s), 3.40-3.51 (4H, m), 3.70 (3H, s), 3.80-4.03 (2H, m), 6.30 (2H, br peak), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.86-6.95 (2H, m), 7.03 (1H, d, J=8Hz), 7.07 (1H, t, J=8Hz), 7.32 (1H, d, J=8Hz), 7.70 (1H, d, J=8Hz), 7.91 (1H, br peak), 10.48 (1H, br peak), 11.49 (1H, br peak)

Example 66

A suspension of 4-(2,3-diaminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (150 mg) and diphenyl N-cyanocarbonimide (64 mg) in 2-propanol (2 ml) was refluxed for 3 hours under nitrogen. The reaction mixture was

evaporated in vacuo and dissolved in chloroform. The organic solution was washed with water, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by preparative thin-layer chromatography (chloroform:ethyl acetate:methanol = 8:1:1/chloroform:methanol = 10:1) to give 4-(2-cyanoamino-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide (45 mg).

NMR (DMSO-d₆, δ) : 1.38-1.51 (2H, m), 1.51-1.66 (2H, m), 1.66-1.81 (2H, m), 2.23 (3H, s), 2.35 (2H, t, J=7Hz), 2.60 (2H, br s), 2.80-2.99 (4H, m), 3.20 (3H, s), 3.50-3.68 (2H, m), 3.74-3.90 (3H, m), 3.90-4.02 (1H, m), 6.65 (1H, d, J=8Hz), 6.80-6.95 (4H, m), 7.03 (1H, d, J=8Hz), 7.07 (1H, d, J=8Hz), 7.55 (1H, d, J=8Hz), 8.29-8.40 (1H, m)

Example 67

A mixture of 4-(2,3-diaminophenyl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg), glyoxal (47 mg) and sodium hydrogen sulfite (169 mg) in ethanol (15 ml) was refluxed for 5 hours. The solution was diluted with chloroform (30 ml) and the solution was washed with water and brine. The organic solution was dried over magnesium sulfate and the solvent was evaporated in vacuo to give an oil. The crude product was purified by silica gel column (1% methanol in chloroform) to give 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(quinoxalin-5-yl)carbonylaminobenzamide (131 mg).

NMR (CDCl₃, δ) : 1.49-1.60 (2H, m), 1.63-1.77 (2H, m), 1.77-1.90 (2H, m), 2.27 (3H, s), 2.32 (3H, s), 2.33-2.46 (6H, m), 3.31 (3H, s), 3.45-3.53 (2H, m), 3.60-3.69 (2H, m), 3.74-4.00 (2H, m), 3.81 (3H, s), 6.54-6.66 (2H, m), 6.87 (1H, d, J=8Hz), 6.98 (1H, d, J=8Hz), 7.06 (1H, s), 8.18-8.27 (3H, m), 8.52

(1H, s), 8.75 (1H, s), 8.93 (2H, s)

Example 68

To a solution of 4-[N-[1-[(tert-butyl)oxycarbonyl]-
5 benzimidazol-4-yl]carbamoyl]-N-[2-[4,4-dimethyl(2,5-
oxazolinyl)]phenyl]-3-methoxy-N-methylbenzamide (400 mg) in
dichloromethane (10 ml) was added trifluoroacetic acid (4
ml). The mixture was stirred at ambient temperature for 3
hours. The reaction mixture was concentrated in vacuo and
10 the residue was diluted with a mixture of chloroform and
saturated aqueous sodium bicarbonate solution. The organic
layer was separated and washed with water and brine. The
solution was dried over magnesium sulfate and the solvent was
evaporated in vacuo. The residue was purified by column
15 chromatography (chloroform:methanol = 100:3) to give 4-[N-
(1H-benzimidazol-4-yl)carbamoyl]-N-[2-[4,4-dimethyl(2,5-
oxazolinyl)]phenyl]-3-methoxy-N-methylbenzamide (290 mg).

NMR (CDCl₃, δ) : 1.38 (3H, s), 1.39 (3H, s), 3.40 (3H,
s), 3.84 (3H, s), 4.01-4.17 (2H, m), 7.05-7.45 (7H,
20 m), 7.62 (1H, d, J=8Hz), 7.79 (1H, d, J=8Hz), 7.96
(1H, s), 8.07 (1H, d, J=8Hz)

Example 69

The following compounds were obtained according to a
25 similar manner to that of Example 68.

- 1) 4-[N-(1H-Benzimidazol-4-yl)carbamoyl]-3-methoxy-N-
methyl-N-[2-(morpholin-4-yl)phenyl]benzamide

NMR (CDCl₃, δ) : 2.30-2.48 (2H, m), 2.77-2.94 (2H, m),
30 3.52 (3H, s), 3.60-3.94 (7H, m), 6.75-7.37 (9H, m),
7.59-8.43 (3H, m)

- 2) 4-[N-(1H-Benzimidazol-4-yl)carbamoyl]-3-methoxy-N-
methyl-N-[2-(1-pyrrolyl)phenyl]benzamide

35 NMR (CDCl₃, δ) : 3.50 (3H, s), 3.91 (3H, s), 6.21-6.30

(2H, m), 6.38-6.46 (2H, m), 6.56-6.68 (2H, m),
7.06-7.53 (7H, m), 7.87-8.07 (2H, m)

- 3) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)carbamoyl]-N-[2-(4-methyl-1-piperazinyl)phenyl]-benzamide

NMR (CDCl₃, δ) : 2.39 (3H, s), 2.41-2.68 (9H, m), 2.86-3.01 (2H, m), 3.52 (3H, s), 3.84 (3H, s), 6.89 (1H, d, J=8Hz), 6.99 (1H, s), 7.07-7.36 (7H, m), 8.10 (1H, d, J=8Hz)

- 4) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)carbamoyl]-N-[2-(2,5-oxazolyl)phenyl]benzamide

NMR (CDCl₃, δ) : 2.60 (3H, s), 3.48 (3H, s), 3.80 (3H, s), 6.78-6.87 (2H, m), 7.13 (1H, dd, J=8, 8Hz), 7.28-7.47 (7H, m), 7.78 (1H, s), 7.88 (1H, m), 7.95 (1H, d, J=8Hz)

Example 70

The solution of 4-[(1-tert-butoxycarbonyl-2-phthalimido-methylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (312 mg) in ethanol (5.0 ml) and 1N sodium hydroxide aqueous solution (1.76 ml) was stirred at ambient temperature for 6 hours. The resulting solution was diluted with water and extracted with chloroform. The organic layer was dried over magnesium sulfate and concentrated in vacuo to afford 4-[(2-aminomethylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (228 mg).

NMR (DMSO-d₆, δ) : 1.38-1.62 (4H, m), 1.68-1.80 (2H, m), 2.15 (3H, s), 2.18-2.37 (6H, m), 2.23 (3H, s), 3.19 (3H, s), 3.26-3.48 (3H, m), 3.69 (3H, s), 3.80-4.01 (3H, m), 4.18 (2H, s), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.90-6.98 (2H, m), 7.14 (1H,

210

d, J=8Hz), 7.20 (1H, t, J=9Hz), 7.56 (1H, d, J=9Hz), 7.62 (1H, d, J=9Hz), 7.95 (1H, d, J=8Hz), 9.06 (1H, s)

5 Example 71

The following compounds were obtained according to a similar manner to that of Example 70.

- 1) 4-[(2-Methylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.46-1.60 (2H, m), 1.66-1.88 (4H, m), 2.28 (6H, s), 2.32-2.42 (6H, m), 2.48 (3H, s), 3.33 (3H, s), 3.44-3.51 (2H, m), 3.58-3.67 (2H, m), 3.78 (3H, s), 3.85-4.00 (2H, m), 6.59 (1H, d, J=8Hz), 6.63 (1H, s), 6.70 (1H, s), 6.87 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 7.04 (1H, s), 7.17 (1H, t, J=8Hz), 7.42 (1H, d, J=8Hz), 7.57 (1H, d, J=8Hz), 8.30-8.40 (2H, m), 8.71 (1H, s)

- 2) 4-[(3-Hydroxymethylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.49-1.89 (6H, m), 2.29 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.34 (3H, s), 3.46-3.51 (2H, m), 3.60-3.66 (2H, m), 3.77 (3H, s), 3.89-4.00 (2H, m), 4.68 (2H, s), 6.62 (1H, d, J=8Hz), 6.67 (1H, s), 6.87 (1H, d, J=8Hz), 6.92 (1H, d, J=8Hz), 7.07 (1H, s), 7.22 (1H, t, J=8Hz), 7.29 (1H, s), 7.36 (1H, d, J=8Hz), 7.52 (1H, d, J=8Hz), 8.32 (1H, d, J=8Hz), 8.48-8.52 (2H, br s)

- 3) 4-[(Indol-6-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.47-1.58 (2H, m), 1.65-1.87 (4H, m),
2.28 (3H, s), 2.29 (3H, s), 2.31-2.41 (6H, m), 3.33
(3H, s), 3.45-3.51 (2H, m), 3.59-3.67 (2H, m), 3.77
(3H, s), 3.83-3.99 (2H, m), 6.58-6.64 (3H, m), 6.88
5 (1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.03 (1H, s),
7.37 (1H, t, J=3Hz), 7.50 (1H, d, J=9Hz), 7.68 (1H,
d, J=9Hz), 7.98 (1H, s), 8.32 (1H, d, J=8Hz), 8.60
(1H, s), 8.83-8.88 (1H, br s)

10 Example 72

The following compound was obtained by using 3-methoxy-
N-methyl-4-[2-[[2-((N-tert-butoxycarbonyl)methylamino)ethyl]-
amino-1-tert-butoxycarbonyl-1H-benzimidazol-4-yl]carbamoyle-N-
[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
15 yloxy]phenyl]benzamide as a starting compound according to a
similar manner to that of Example 23.

3-Methoxy-N-methyl-4-[2-[[2-(methylamino)ethyl]amino]-
1H-benzimidazol-4-yl]carbamoyle-N-[4-methyl-2-[5-(4-
20 methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
NMR (CDCl₃, δ) : 1.36-1.50 (2H, m), 1.50-1.64 (2H, m),
1.64-1.82 (2H, m), 2.15 (3H, s), 2.17-2.36 (9H, m),
2.40 (3H, s), 2.88 (2H, t, J=5Hz), 3.22 (3H, s),
3.25-3.55 (7H, m), 3.81-4.02 (5H, m), 6.64 (1H, d,
25 J=8Hz), 6.70-6.85 (3H, m), 6.90 (1H, d, J=8Hz),
7.00 (1H, d, J=8Hz), 7.03-7.13 (2H, m), 7.77-7.90
(2H, m)

Example 73

30 The following compound was obtained according to a
similar manner to that of Example 72.

4-[2-[(2-Aminoethyl)methylamino]-1H-benzimidazol-4-yl]-
carbamoyle-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
35 methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

212

NMR (DMSO-d₆, δ) : 1.37-1.50 (2H, m), 1.50-1.63 (2H, m), 1.68-1.82 (2H, m), 2.14 (3H, s), 2.16-2.38 (9H, m), 2.93 (2H, t, J=5Hz), 3.14 (3H, s), 3.21 (3H, s), 3.36-3.47 (4H, m), 3.55 (2H, t, J=5Hz), 3.81-4.02 (2H, m), 6.63 (1H, d, J=8Hz), 6.76-6.87 (2H, m), 6.91 (1H, d, J=8Hz), 7.00 (1H, d, J=8Hz), 7.04-7.12 (2H, m), 7.88 (1H, d, J=8Hz), 7.95 (1H, d, J=8Hz)

10 Example 74

The following compound was obtained by using 4-[2-tert-butoxycarbonylamino-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide as a starting compound according to a similar manner to that of Example 23.

4-[2-Amino-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

20 NMR (DMSO-d₆, δ) : 1.37-1.50 (2H, m), 1.50-1.65 (2H, m), 1.65-1.82 (2H, m), 2.14 (3H, s), 2.17-2.38 (9H, m), 3.23 (3H, s), 3.36-3.48 (4H, m), 3.67-4.05 (5H, m), 6.20 (2H, br peak), 6.65 (1H, d, J=8Hz), 6.73-6.93 (3H, m), 6.96-7.14 (3H, m), 7.84-7.92 (2H, m)

25

Example 75

The following compounds were obtained according to a similar manner to that of Preparation 13.

30 1) 4-(2-Aminomethyl-1-methyl-1H-benzimidazol-4-yl)carbonyl-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
NMR (CDCl₃, δ) : 1.44-1.75 (4H, m), 1.75-1.90 (2H, m), 2.25 (3H, s), 2.28 (3H, s), 2.30-2.41 (6H, m), 3.32 (3H, s), 3.43-3.52 (2H, m), 3.57-3.66 (2H, m),

35

213

3.74-3.90 (7H, m), 3.95 (1H, br peak), 4.20 (2H, s), 6.53-6.63 (2H, m), 6.87 (1H, d, J=8Hz), 6.98 (1H, d, J=8Hz), 7.03 (1H, s), 7.37 (1H, t, J=8Hz), 7.46 (1H, d, J=8Hz), 8.15 (1H, d, J=8Hz), 8.55 (1H, d, J=8Hz)

2) 4-(2-Aminomethyl-3-methyl-3H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.48-1.62 (2H, m), 1.62-1.93 (4H, m), 2.30 (3H, s), 2.32-2.41 (5H, m), 2.41-2.55 (4H, m), 3.33 (3H, s), 3.51-3.60 (2H, m), 3.60-3.77 (5H, m), 3.81 (3H, s), 3.87-4.03 (2H, m), 4.13 (2H, br peak), 6.61 (1H, d, J=8Hz), 6.65 (1H, s), 6.88 (1H, d, J=8Hz), 6.97 (1H, d, J=8Hz), 7.03 (1H, s), 7.22-7.30 (1H, m), 7.42 (1H, d, J=8Hz), 7.86 (1H, br peak), 8.30 (1H, d, J=8Hz), 8.35 (1H, s)

Example 76

A solution of 3-methoxy-N-methyl-N-(4-methyl-2-benzyloxyphenyl)-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylamino benzamide (260 mg) in 90% trifluoroacetic acid (2 ml) was stirred at ambient temperature for 2 hours and the solvent was evaporated in vacuo. The residue was stirred with chloroform (10 ml) and saturated aqueous sodium hydrogencarbonate (10 ml) and the organic phase was separated. The solution was washed with brine and dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by NH type silica gel column (chloroform) to give 4-(2-aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-(4-methyl-2-benzyloxyphenyl)benzamide (130 mg).

NMR (CDCl₃, δ) : 2.22 (3H, s), 3.39 (3H, s), 3.57 (3H, s), 4.15 (2H, s), 4.88 (1H, d, J=12Hz), 5.04 (1H, d, J=12Hz), 6.60-6.70 (2H, m), 6.85-7.01 (3H, m),

214

7.32 (1H, m), 7.38 (5H, s), 7.46 (1H, br s), 8.08 (1H, br s), 8.47 (1H, br s)

Example 77

5 The following compounds were obtained according to a similar manner to that of Example 76.

- 1) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-(2,4-dimethylphenyl)-3-methoxy-N-methylbenzamide

10 NMR (DMSO-d₆, δ) : 2.11 (3H, s), 2.19 (3H, s), 3.21 (3H, s), 3.72 (3H, s), 4.08 (2H, s), 6.88-7.05 (4H, m), 7.14 (1H, d, J=8Hz), 7.31 (1H, t, J=8Hz), 7.71 (1H, d, J=8Hz), 7.89 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

15

- 2) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-(2-methoxy-4-methylphenyl)-N-methylbenzamide

20 NMR (CDCl₃, δ) : 2.25 (3H, s), 3.34 (3H, s), 3.67 (3H, s), 3.70 (3H, s), 4.15 (2H, s), 6.55-6.63 (2H, m), 6.82-7.00 (3H, m), 7.26 (1H, t, J=8Hz), 7.48 (1H, br), 8.08 (1H, br), 8.52 (1H, br)

20

- 3) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonylphenylmethoxy]phenyl]benzamide

25

30 NMR (CDCl₃, δ) : 2.26 (3H, s), 2.30 (3H, s), 2.31-2.50 (4H, m), 3.40 (3H, s), 3.40-3.51 (2H, m), 3.62 (3H, s), 3.68-3.83 (2H, m), 4.17 (2H, s), 4.88 (1H, d, J=12Hz), 5.07 (1H, d, J=12Hz), 6.62 (1H, s), 6.70 (1H, d, J=8Hz), 6.87-6.94 (2H, m), 7.03 (1H, d, J=8Hz), 7.20-7.44 (5H, m), 7.52 (1H, m), 7.99 (1H, br), 8.46 (1H, m)

30

- 4) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[3-(4-methylpiperazin-1-

35

yl)carbonylprop-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 2.05-2.16 (2H, m), 2.23 (3H, s), 2.30-
2.41 (4H, m), 2.52 (2H, t, J=7.5Hz), 3.33 (3H, s),
3.43-3.50 (2H, m), 3.59-3.65 (2H, m), 3.75 (3H, s),
3.86-4.06 (2H, m), 4.21 (2H, s), 6.58-6.67 (2H, m),
6.90-7.02 (3H, m), 7.28 (1H, t, J=8Hz), 7.56 (1H,
br), 8.06 (1H, br), 8.49 (1H, br)

5) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonylbut-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.76-1.93 (4H, m), 2.25 (3H, s), 2.28 (3H, s), 2.31-2.47 (4H, m), 3.33 (3H, s), 3.49 (1H, m), 3.60-3.74 (5H, m), 3.86 (1H, m), 3.96 (1H, m), 4.17 (2H, s), 6.59-6.65 (2H, m), 6.86-6.95 (2H, m), 7.00 (1H, d, J=8Hz), 7.21 (1H, t, J=8Hz), 7.39 (1H, m), 8.02 (1H, m), 8.50 (1H, m)

6) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-(5-piperazin-1-ylcarbonylpent-1-yloxy)phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.63 (2H, m), 1.70-1.81 (2H, m), 2.22 (3H, s), 2.37 (2H, t, J=7.5Hz), 2.98-3.13 (4H, m), 3.20 (3H, s), 3.61-3.71 (4H, m), 3.77 (3H, s), 3.88 (1H, m), 3.97 (1H, m), 4.42 (2H, s), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.92-6.99 (2H, m), 7.05 (1H, d, J=8Hz), 7.41 (1H, t, J=8Hz), 7.83 (1H, d, J=8Hz), 7.97 (1H, d, J=8Hz), 8.35 (1H, br)

7) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-(5-carbamoylpent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methylbenzamide

NMR (CDCl₃, δ) : 1.46-1.57 (2H, m), 1.66-1.82 (4H, m), 2.20-2.30 (2H, m), 2.23 (3H, s), 3.32 (3H, s), 3.61

216

(3H, s), 3.76 (1H, m), 3.91 (1H, m), 4.12 (2H, s),
5.93 (1H, br), 6.32 (1H, br), 6.54-6.63 (2H, m),
6.73 (1H, s), 6.93 (1H, d, J=8Hz), 7.01 (1H, d,
J=8Hz), 7.18 (1H, t, J=8Hz), 7.43 (1H, m), 7.90
(1H, m), 8.46 (1H, d, J=8Hz)

8) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-
[2-(5-dimethylcarbamoylpent-1-yloxy)-4-methylphenyl]-3-
methoxy-N-methylbenzamide

10 NMR (DMSO-d₆, δ) : 1.48-1.60 (2H, m), 1.66-1.77 (2H,
m), 1.77-1.90 (2H, m), 2.28 (3H, s), 2.38 (2H, t,
J=7.5Hz), 2.93 (3H, s), 3.00 (3H, s), 3.35 (3H, s),
3.72 (3H, s), 3.85 (1H, m), 3.96 (1H, m), 4.17 (2H,
s), 6.55-6.63 (2H, m), 6.86 (1H, d, J=8Hz), 6.91
15 (1H, s), 6.97 (1H, d, J=8Hz), 7.24 (1H, t, J=8Hz),
7.49 (1H, br), 8.05 (1H, br), 8.50 (1H, br)

9) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-
[2-[5-(2,2-dimethylhydrazino)carbonylpent-1-yloxy]-4-
methylphenyl]-3-methoxy-N-methylcarbonylaminobenzamide

20 NMR (CDCl₃, δ) : 1.42-1.62 (2H, m), 1.62-1.90 (4H, m),
2.12 (2H, t, J=7.5Hz), 2.27 (3H, s), 2.50 (3H, s),
2.58 (3H, s), 3.34 (3H, s), 3.71 (3H, s), 3.77-4.00
(2H, m), 4.20 (2H, s), 6.27 (1H, br), 6.52-6.67
25 (2H, m), 6.83-7.11 (3H, m), 7.24 (1H, m), 7.50 (1H,
br), 8.07 (1H, br), 8.53 (1H, br)

10) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-oxopiperidin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

30 NMR (CDCl₃, δ) : 1.41-1.60 (2H, m), 1.60-1.89 (4H, m),
2.21 (3H, s), 2.30-2.50 (6H, m), 3.30 (3H, s),
3.41-3.98 (6H, m), 4.22 (2H, s), 6.51-6.62 (2H, m),
6.78-6.99 (3H, m), 7.24 (1H, m), 7.55 (1H, br),
35 7.96 (1H, br), 8.45 (1H, br)

11) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-[5-(4-hydroxypiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-3-methoxy-N-methylbenzamide

NMR (CDCl₃, δ) : 1.35-1.54 (4H, m), 1.54-1.72 (2H, m),
1.72-1.91 (4H, m), 2.24 (3H, s), 2.28-2.41 (2H, m),
2.95-3.21 (2H, m), 3.30 (3H, s), 3.46-3.98 (4H, m),
3.70 (3H, s), 4.09 (1H, m), 4.20 (2H, s), 6.52-6.63
(2H, m), 6.78-6.97 (3H, m), 7.13 (1H, m), 7.41 (1H,
br), 7.88 (1H, br), 8.38 (1H, d, J=8Hz)

12) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-[5-[N-(2-dimethylaminoeth-1-yl)-N-methylamino-carbonyl]pent-1-yloxy]-4-methylphenyl]-3-methoxy-N-methylbenzamide

NMR (CDCl₃, δ) : 1.31-1.52 (2H, m), 1.59-1.81 (4H, m),
2.23 (3H, s), 2.30-2.41 (2H, m), 2.73-2.88 (2H, m),
2.85 (6H, sx2), 3.05 (3H, s), 3.29 (1H, s), 3.55-
3.94 (4H, m), 4.46 (2H, m), 6.52-6.63 (2H, m), 6.67
(1H, s), 6.82-7.10 (3H, m), 7.31 (1H, m), 7.75 (1H,
m), 8.33 (1H, m)

Example 78

The following compound was obtained according to similar manners to those of Examples 38 and 76.

4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-(6-hydroxyhex-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methylbenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.31-1.50 (4H, m), 1.67-1.76 (2H,
m), 2.21 (3H, s), 3.18 (3H, s), 3.40 (2H, t,
J=7.5Hz), 3.74 (3H, s), 3.74-3.85 (2H, m), 3.85
(1H, m), 3.97 (1H, m), 4.45 (1H, m), 4.79 (2H, m),
6.64 (1H, d, J=8Hz), 6.81 (1H, s), 7.90 (1H, d,
J=8Hz), 6.97 (1H, s), 7.03 (1H, d, J=8Hz), 7.39
(1H, t, J=8Hz), 7.77 (1H, d, J=8Hz), 7.94 (1H, d,

J=8Hz), 8.34 (1H, d, J=8Hz), 8.71 (1H, br)

Example 79

To the solution of 3-methoxy-N-methyl-N-[4-methyl-2-[5-
5 (4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(3-p
hthalimidopropyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide
(160 mg) in ethanol (5 ml) was added hydrazine hydrate (49 mg)
and stirred at ambient temperature for 6 hours. After the
reaction mixture was concentrated in vacuo, the residue was
10 purified by preparative thin-layer chromatography
(methanol:chloroform:28% ammonia solution = 3:5:1). To the
purified product was added water and 1N hydrochloric acid
(0.51 ml). The solution was lyophilized to give 4-[2-(3-
aminopropyl)-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-
15 methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-
1-yloxy]phenyl]benzamide trihydrochloride (70 mg).

NMR (DMSO-d₆, δ) : 1.42-1.50 (2H, m), 1.50-1.63 (2H,
m), 1.70-1.82 (2H, m), 2.12-2.22 (2H, m), 2.22 (3H,
s), 2.40 (2H, t, J=7Hz), 2.73 (3H, s), 2.80-3.17
20 (6H, m), 3.20 (3H, s), 3.33-3.57 (4H, m), 3.72 (3H,
s), 3.87-4.10 (3H, m), 4.40-4.47 (1H, m), 6.63 (1H,
d, J=8Hz), 6.82 (1H, s), 6.90-6.93 (2H, m), 7.04
(1H, d, J=8Hz), 7.43 (1H, t, J=8Hz), 7.82 (1H, d,
J=8Hz), 7.97-8.13 (2H, m), 8.21 (2H, br s)

25

Example 80

The solution of 4-[(1-tert-butoxycarbonyl-2-tert-
butyldiphenylsilyloxymethylindol-4-yl)carbonyl]amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
30 yl)carbonylpent-1-yloxy]phenyl]benzamide (615 mg) in 1N
sodium hydroxide aqueous solution (3.1 ml) and methanol (10
ml) was stirred at ambient temperature overnight. The
resulting solution was neutralized with 1N hydrochloric acid
and extracted with ethyl acetate (20 ml). The organic layer
35 was washed with brine, dried over magnesium sulfate and

concentrated in vacuo. The residue was purified by silica gel column chromatography (eluent; 8-10% methanol in chloroform) to give 4-[(2-hydroxymethylindol-4-yl)carbonyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (362 mg).

NMR (CDCl₃, δ) : 1.44-1.57 (2H, m), 1.63-1.85 (4H, m), 2.27 (3H, s), 2.29 (3H, s), 2.30-2.40 (6H, m), 3.33 (3H, s), 3.43-3.50 (2H, m), 3.57-3.63 (2H, m), 3.70 (3H, s), 3.82-3.99 (2H, m), 4.80 (2H, s), 6.59-6.67 (2H, m), 6.77 (1H, s), 6.89 (1H, d, J=8Hz), 6.95 (1H, d, J=8Hz), 6.98 (1H, s), 7.18 (1H, t, J=8Hz), 7.42 (1H, d, J=8Hz), 7.57 (1H, d, J=8Hz), 8.32 (1H, d, J=8Hz), 8.64 (1H, s), 8.98-9.03 (1H, br s)

Example 81

The following compound was obtained by using 4-[(1-tert-butoxycarbonyl-2-benzyloxymethylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide as a starting compound according to a similar manner to that of Preparation 10.

4-[(1-tert-Butoxycarbonyl-2-hydroxymethylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.40-1.61 (4H, m), 1.50 (9H, s), 1.69-1.79 (2H, s), 2.23 (3H, s), 2.35-2.42 (2H, m), 2.69 (3H, s), 2.98-3.22 (9H, m), 3.28-3.54 (5H, m), 3.64 (3H, s), 3.84-4.00 (2H, m), 4.10-4.18 (1H, br), 4.31-4.40 (1H, m), 4.89 (1H, t, J=5Hz), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.88-6.92 (2H, m), 7.03 (1H, d, J=8Hz), 7.20-7.33 (2H, m), 7.76 (1H, d, J=8Hz), 9.13 (1H, s)

Example 82

To an ice bath cooled solution of 4-(2-aminomethyl-1H-

benzimidazol-4-yl)carbonylamino-N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methylbenzamide (2.95 g) in dichloromethane (30 ml) were added triethylamine (496 mg) and di-tert-butyl dicarbonate (1.07 g) and the mixture was stirred at ambient temperature for 3 hours. The solution was washed successively with water, 10% hydrochloric acid, saturated aqueous sodium hydrogencarbonate and brine and the organic phase was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by silica gel column (1% methanol in chloroform) to give N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonylamino)methyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (2.65 g).

NMR (CDCl₃, δ) : 1.24 (3H, t, J=7.5Hz), 1.41-1.55 (2H, m), 1.48 (9H, s), 1.63-1.84 (4H, m), 2.23 (3H, s), 2.31 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.78 (3H, s), 3.78-3.98 (2H, m), 4.12 (2H, q, J=7.5Hz), 4.58 (2H, m), 5.68 (1H, br t, J=7Hz), 6.53-6.63 (2H, m), 6.83-7.04 (3H, m), 7.30 (1H, t, J=8Hz), 7.50 (1H, d, J=8Hz), 8.11 (1H, d, J=8Hz), 8.49 (1H, d, J=8Hz)

Example 83

The solution of 4-[(2-ethoxycarbonylindolin-4-yl)-carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (175 mg) in ammonia-methanol solution (8.0 ml) was stood overnight at ambient temperature. The solvent was evaporated in vacuo and the residue was purified by silica gel column chromatography (eluent; 8-10% methanol in chloroform) to give 4-[(2-carbamoylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (160 mg).

NMR (CDCl₃, δ) : 1.48-1.58 (2H, m), 1.64-1.87 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.41 (6H, m), 3.32 (3H, s), 3.46-3.51 (2H, m), 3.59-3.65 (2H, m), 3.77

221

(3H, s), 3.84-3.98 (3H, m), 4.38-4.49 (2H, m),
5.46-5.50 (1H, br s), 6.59 (1H, d, J=8Hz), 6.62
(1H, s), 6.69-6.74 (1H, br), 6.85 (1H, d,
J=8Hz), 6.91 (1H, d, J=8Hz), 7.01 (1H, s), 7.09 (1H,
5 d, J=8Hz), 7.18 (1H, t, J=8Hz), 8.22 (1H, d,
J=8Hz), 8.37 (1H, s)

Example 84

To a solution of 4-[(2-hydroxymethylindol-4-yl)-
10 carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
(295 mg) in dichloromethane (8.0 ml) was added manganese(IV)
oxide (196 mg) and the mixture was stirred at ambient
15 temperature for 3 hours. The resulting mixture was filtered
through a bed of celite and the filtrate was concentrated in
vacuo. The residue was triturated with diethyl ether-n-
hexane (1:3) to give 4-[(2-formylindol-4-yl)carbonyl]amino-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide (255 mg).

20 NMR (CDCl₃, δ) : 1.49-1.61 (2H, m), 1.68-1.89 (4H, m),
2.28 (3H, s), 2.29 (3H, s), 2.31-2.43 (6H, m), 3.36
(3H, s), 3.47-3.52 (2H, m), 3.60-3.68 (2H, m), 3.78
(3H, s), 3.86-4.00 (2H, m), 6.60 (1H, d, J=8Hz),
6.65 (1H, s), 6.88 (1H, d, J=8Hz), 6.98 (1H, d,
25 J=8Hz), 7.07 (1H, s), 7.46 (1H, t, J=8Hz), 7.58
(1H, d, J=8Hz), 7.62 (1H, d, J=8Hz), 7.84 (1H, s),
8.35 (1H, d, J=8Hz), 8.62 (1H, s), 9.44-9.50 (1H,
br s), 9.89 (1H, s)

Example 85

A solution of 4-(2-aminomethyl-1H-benzimidazol-4-
yl)carbonylamino-3-methoxy-N-methyl-N-(2-benzyloxy-4-
methylphenyl)benzamide (120 mg) in methanol (15 ml) was
stirred under atmospheric pressure of hydrogen in the presence
35 of palladium hydroxide (20 mg) at ambient temperature

overnight. After removal of the catalyst by filtration, the filtrate was evaporated in vacuo to give 4-(2-aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-(2-hydroxy-4-methylphenyl)benzamide (85 mg).

5 NMR (DMSO-d₆, δ) : 2.13 (3H, s), 3.19 (3H, s), 3.77
(3H, s), 4.63 (2H, m), 6.47 (1H, d, J=8Hz), 6.69
(1H, s), 6.86 (1H, d, J=8Hz), 6.97 (1H, d, J=8Hz),
7.03 (1H, s), 7.42 (1H, t, J=8Hz), 7.83 (1H, d,
J=8Hz), 7.98 (1H, d, J=8Hz), 8.30 (1H, br), 8.78
0 (2H, br)

Example 86

The following compound was obtained by using 4-(2-amino-3-nitrophenyl)carbonylamino-N-[2-(4-methoxybenzyloxy)-4-methylphenyl]-3-methoxy-N-methylbenzamide as a starting compound according to a similar manner to that of Example 85.

4-(2-Amino-3-nitrophenyl)carbonylamino-N-(2-hydroxy-4-methylphenyl)-3-methoxy-N-methylbenzamide

20 NMR (CDCl₃, δ) : 2.23 (3H, s), 3.37 (3H, s), 3.67 (3H, s), 6.48-7.02 (7H, m), 7.63 (1H, d, J=8Hz), 8.03-8.13 (3H, m), 8.21-8.30 (2H, m)

Example 87

25 The following compound was obtained by using N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonylamino)methyl-1H-benzimidazol-4-yl]carbonylaminobenzamide as a starting compound according to a similar manner to that of Example 29.

30

N-[2-(5-Carboxypent-1-yloxy)-4-methylphenyl]-3-methoxy-
N-methyl-4-[2-(tert-butoxycarbonylamino)methyl-1H-
benzimidazol-4-yl]carbonylaminobenzamide

35 NMR (CDCl₃, δ) : 1.50 (9H, s), 1.50-1.63 (2H, m), 1.69-1.85 (4H, m), 2.24 (3H, s), 2.36-2.50 (2H, m), 3.33

223

(3H, s), 3.63 (3H, s), 3.80 (1H, m), 3.95 (1H, m),
4.54 (2H, m), 6.09 (1H, br), 6.52-6.63 (2H, m),
6.78-7.01 (3H, m), 7.22 (1H, t, J=8Hz), 7.48 (1H,
m), 7.90 (1H, m), 8.39 (1H, m)

5

Example 88

A mixture of N-[2-(5-carboxypent-1-yloxy)-4-methyl-phenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonylamino)-methyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (200 mg),
10 4-(tert-butoxycarbonyl)piperazine (66.3 mg), 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride (68.3 mg) and 1-hydroxybenzotriazol (48.1 mg) in N,N-dimethylformamide (5 ml) was stirred at ambient temperature overnight and the mixture was diluted with ethyl acetate. The solution was
15 washed successively with saturated aqueous sodium hydrogen carbonate solution, water and brine and dried over magnesium sulfate. The solvent was evaporated and the residue was purified by silica gel column (chloroform) to give 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-tert-butoxycarbonylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(tert-butoxycarbonyl)-aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (250
20 mg).

NMR (CDCl₃, δ) : 1.45-1.56 (2H, m), 1.45 (9H, s), 1.50 (9H, s), 1.62-1.87 (4H, m), 2.26 (3H, s), 2.35 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.35-3.47 (6H, m), 3.51-3.60 (2H, m), 3.77-3.99 (2H, m), 3.82 (3H, s), 4.37 (2H, m), 5.74 (1H, br), 6.55-6.62 (2H, m), 6.84-7.03 (3H, m), 7.31 (1H, t, J=8Hz), 7.51 (1H, d, J=8Hz), 8.12 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz)

30

Example 89

The following compounds were obtained according to a similar manner to that of Example 88.

35 1) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-oxopiperidin-1-

yl)carbonylpent-1-yloxy]phenyl]-4-[2-(tert-butoxy-carbonyl)aminomethyl-1H-benzimidazol-4-yl]-carbonylaminobenzamide

5 NMR (CDCl₃, δ) : 1.47-1.60 (2H, m), 1.47 (9H, s), 1.70-1.88 (4H, m), 2.23 (3H, s), 2.37-2.57 (2H, m), 3.33 (3H, s), 3.69-3.99 (4H, m), 3.80 (3H, s), 4.60 (2H, d, J=7Hz), 5.75 (1H, br), 6.55-6.66 (2H, m), 6.85-7.01 (3H, m), 7.31 (1H, t, J=8Hz), 7.59 (1H, d, J=8Hz), 8.00 (1H, d, J=8Hz), 8.46 (1H, d, J=8Hz)

10

2) N-[2-(5-Carbamoylpent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonylamino)methyl-1H-benzimidazol-4-yl]carbonylaminobenzamide

15 NMR (CDCl₃, δ) : 1.50 (9H, s), 1.60-1.86 (6H, m), 2.21-2.30 (2H, m), 2.26 (3H, s), 3.33 (3H, s), 3.70 (3H, s), 3.72-3.97 (2H, m), 4.56 (2H, m), 6.57-6.68 (2H, m), 6.93-7.05 (3H, m), 7.33 (1H, m), 7.54 (1H, m), 8.01 (1H, s), 8.11 (1H, d, J=8Hz), 8.50 (1H, m)

20 3) N-[2-(5-Dimethylcarbamoylpent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonylamino)methyl-1H-benzimidazol-4-yl]carbonylaminobenzamide

25 NMR (CDCl₃, δ) : 1.45-1.57 (2H, m), 1.49 (9H, s), 1.64-1.85 (4H, m), 2.27 (3H, s), 2.35 (2H, t, J=7.5Hz), 2.92 (3H, s), 3.00 (3H, s), 3.34 (3H, s), 3.79 (3H, s), 3.84 (1H, m), 3.94 (1H, m), 4.58 (2H, d, J=7Hz), 5.72 (1H, br), 6.55-6.64 (2H, m), 6.90 (1H, d, J=8Hz), 6.92-7.01 (2H, m), 7.28 (1H, t, J=8Hz), 7.48 (1H, d, J=8Hz), 8.02 (1H, s), 8.10 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz)

30

4) N-[2-[5-(2,2-Dimethylhydrazino)carbonylpent-1-yloxy]-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonylamino)methyl-1H-benzimidazol-4-yl]-carbonylaminobenzamide

35

225

NMR (CDCl₃, δ) : 1.41-1.58 (2H, m), 1.50 (9H, s), 1.63-1.89 (4H, m), 2.13 (2H, t, J=7.5Hz), 2.27 (3H, s), 2.50 (3H, s), 2.58 (3H, s), 3.35 (3H, s), 3.72-4.01 (2H, m), 3.79 (3H, s), 4.58 (2H, m), 5.81 (1H, br), 6.52-6.67 (2H, m), 6.84-7.05 (3H, m), 7.30 (1H, t, J=8Hz), 7.49 (1H, m), 8.11 (1H, m), 8.51 (1H, m)

5) N-[2-[5-[N-(2-Dimethylaminoeth-1-yl)-N-methylamino-carbonyl]pent-1-yloxy]-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylamino benzamide

NMR (CDCl₃, δ) : 1.41-1.56 (2H, m), 1.48 (9H, s), 1.61-1.74 (2H, m), 1.74-1.85 (2H, m), 2.24 (3H, s), 2.32 (6H, s), 2.32-2.44 (2H, m), 2.53 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.52 (2H, t, J=7.5Hz), 3.78 (3H, s), 3.86 (1H, m), 3.96 (1H, m), 4.58 (2H, m), 5.92 (1H, br), 6.53-6.61 (2H, m), 6.87 (1H, d, J=8Hz), 6.91-7.02 (2H, m), 7.29 (1H, t, J=8Hz), 7.50 (1H, m), 8.11 (1H, d, J=8Hz), 8.49 (1H, m)

Example 90

The following compound was obtained by using 4-(1H-benzimidazol-4-yl)carbonylamino-N-(2-carboxy-4-methylphenyl)-3-methoxy-N-methylbenzamide as a starting compound according to a similar manner to that of Example 88.

4-(1H-Benzimidazol-4-yl)carbonylamino-N-(2-carbamoyl-4-methylphenyl)-3-methoxy-N-methylbenzamide

NMR (CDCl₃, δ) : 2.29 (3H, s), 3.45 (3H, s), 3.71 (3H, s), 6.93 (1H, d, J=8Hz), 7.00-7.10 (2H, m), 7.14-7.43 (3H, m), 7.67 (1H, d, J=8Hz), 7.93 (1H, d, J=8Hz), 8.09 (1H, s), 8.31 (1H, d, J=8Hz)

Example 91

The following compound was obtained according to a

similar manner to that of Example 90.

N-(2-Dimethylcarbamoyl-4-methylphenyl)-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-
5 benzamide

NMR (CDCl₃, δ) : 2.31 (3H, s), 2.56 (3H, s), 2.99 (6H, s), 3.42 (3H, s), 3.76 (3H, s), 6.89-7.32 (6H, m), 7.44 (1H, d, J=8Hz), 8.08 (1H, d, J=8Hz), 8.59 (1H, br)

Example 92

To a solution of 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-oxopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]-
15 carbonylaminobenzamide (150 mg) in methanol (5 ml) was added sodium borohydride (7.52 mg) under an ice bath cooling and the mixture was stirred at the same temperature for 1 hour. The mixture was diluted with chloroform and the solution was washed with water and brine. The organic phase was dried
20 over magnesium sulfate and the solvent was evaporated in vacuo to give N-[2-[5-(4-hydroxypiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (150 mg).

NMR (CDCl₃, δ) : 1.48 (9H, s), 1.48-1.57 (2H, m), 1.64-1.94 (8H, m), 2.27 (3H, s), 2.30-2.41 (2H, m), 3.03-3.23 (2H, m), 3.33 (3H, s), 3.79 (3H, s), 3.80-3.97 (4H, m), 4.12 (1H, m), 4.61 (2H, d, J=7Hz), 5.86 (1H, br), 6.54-6.63 (2H, m), 6.84-7.00 (3H, m), 7.32 (1H, t, J=8Hz), 7.59 (1H, d, J=8Hz),
30 7.98 (1H, br), 8.43 (1H, d, J=8Hz)

Example 93

A mixture of 4-(2-amino-3-nitrophenyl)carbonylamino-N-(2-hydroxy-4-methylphenyl)-3-methoxy-N-methylbenzamide (400

mg), acetic anhydride (90.7 mg) and triethylamine (89.9 mg) in dichloromethane (20 ml) was stirred in an ice bath for 4 hours. The mixture was diluted with chloroform and the solution was washed with water and brine. The organic phase
5 was dried over magnesium sulfate and the solvent was evaporated in vacuo to give N-(2-acetoxy-4-methylphenyl)-4-(2-amino-3-nitrophenyl)carbonylamino-3-methoxy-N-methylbenzamide (425 mg).

10 NMR (CDCl₃, δ) : 2.21 (3H, s), 2.30 (3H, s), 3.35 (3H, s), 3.72 (3H, s), 6.68 (1H, t, J=8Hz), 6.88 (1H, s), 6.92-7.01 (2H, m), 7.10 (1H, d, J=8Hz), 7.70 (1H, d, J=8Hz), 8.14 (2H, br), 8.23 (1H, d, J=8Hz), 8.22 (1H, d, J=8Hz), 8.36 (1H, br)

15 Example 94

A mixture of 4-(2-amino-3-nitrophenyl)carbonylamino-N-(2-hydroxy-4-methylphenyl)-3-methoxy-N-methylbenzamide (520 mg), N-(4-bromobutyl)phthalimide (326 mg) and potassium carbonate (160 mg) in N,N-dimethylformamide (10 ml) was
20 heated at 60°C for 8 hours. The mixture was diluted with ethyl acetate and the solution was washed with water and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo. The crude oil was purified by silica gel column (1% methanol in chloroform) to
25 give 4-(2-amino-3-nitrophenyl)carbonylamino-3-methoxy-N-methyl-N-[2-(4-phthalimidobut-1-yloxy)-4-methylphenyl]-benzamide (670 mg).

30 NMR (CDCl₃, δ) : 1.78-1.96 (4H, m), 2.27 (3H, s), 3.31 (3H, s), 3.68-3.80 (5H, m), 3.92 (1H, m), 4.00 (1H, m), 6.57-6.72 (3H, m), 6.81-7.08 (3H, m), 7.66-7.73 (2H, m), 7.81-7.88 (2H, m), 8.09-8.21 (2H, m), 8.32 (1H, m)

Example 95

35 The following compound was obtained by using N-[2-(4-

phthalimidobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylaminobenzamide as a starting compound according to a similar manner to that of Example 26.

5

N-[2-(4-Aminobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylaminobenzamide

10 NMR (CDCl₃, δ) : 1.53-1.70 (2H, m), 1.75-1.86 (2H, m),
2.23 (3H, s), 2.57 (3H, s), 2.77 (2H, t, J=7.5Hz),
3.34 (3H, s), 3.66 (3H, s), 3.80 (1H, m), 3.92 (1H, m),
6.54-6.61 (2H, m), 6.81-6.92 (2H, m), 6.98 (1H, d, J=8Hz),
7.21 (1H, t, J=8Hz), 7.51 (1H, br), 7.94 (1H, br), 8.45 (1H, br)

15

Example 96

The following compound was obtained according to a similar manner to that of Example 95.

20 N-[2-(6-Aminohex-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide

25 NMR (CDCl₃, δ) : 1.33-1.50 (4H, m), 1.48 (9H, s), 1.52-1.63 (2H, m), 1.68-1.80 (2H, m), 2.25 (3H, s), 2.79 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.72 (3H, s), 3.78 (1H, m), 3.90 (1H, m), 4.52 (2H, m), 6.02 (1H, br), 6.51-6.62 (2H, m), 6.86-7.00 (3H, m), 7.20 (1H, t, J=8Hz), 7.48 (1H, d, J=8Hz), 7.91 (1H, d, J=8Hz), 8.32 (1H, d, J=8Hz)

30

Example 97

35 The following compound was obtained by using N-[2-(5-tert-butoxycarbonylaminopent-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylaminobenzamide as a starting compound according to a

similar manner to that of Example 23.

N-[2-(5-Aminopent-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-
5 benzamide

NMR (CDCl₃, δ) : 1.40-1.53 (2H, m), 1.53-1.64 (2H, m),
1.71-1.80 (2H, m), 2.25 (3H, s), 2.59 (3H, s), 2.78
(2H, t, J=7.5Hz), 3.35 (3H, s), 3.68 (3H, s), 3.76
(1H, m), 3.92 (1H, m), 6.51-6.62 (2H, m), 6.83 (1H,
10 s), 6.91 (1H, d, J=8Hz), 7.00 (1H, d, J=8Hz), 7.22
(1H, t, J=8Hz), 7.52 (1H, d, J=8Hz), 7.90 (1H, br),
8.41 (1H, d, J=8Hz)

Example 98

15 The following compound was obtained according to a
similar manner to that of Example 97.

N-[2-(4-Aminobut-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylbenzamide

20 MASS (m/z) : 516

Example 99

To a solution of N-[2-(4-aminobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-
25 4-yl)carbonylaminobenzamide (120 mg) in dichloromethane (10 ml) was added acetic anhydride (23.8 mg) and the mixture was stirred at ambient temperature for 1 hour. The solution was washed with water and brine and dried over magnesium sulfate and the solvent was evaporated in vacuo. The residue was
30 purified by silica gel column (2% methanol in chloroform) and the product was solidified with diethyl ether to give N-[2-(4-acetylaminobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylaminobenzamide (87 mg).

35 NMR (CDCl₃, δ) : 1.58-1.86 (2H, m), 2.08 (3H, s), 2.26

230

(3H, s), 2.69 (3H, s), 3.19-3.41 (2H, m), 3.35 (3H, s), 3.73-3.87 (2H, m), 3.76 (3H, s), 6.23 (1H, br), 6.51-6.78 (3H, m), 6.90-7.10 (3H, m), 7.27 (1H, m), 7.59 (1H, d, J=8Hz), 7.91 (1H, d, J=8Hz), 8.45 (1H, d, J=8Hz)

5

Example 100

A mixture of N-[2-(4-aminobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylaminobenzamide (150 mg), 37% formaldehyde solution (43.7 mg) and sodium cyanoborohydride (18.3 mg) in methanol (10 ml) was stirred at ambient temperature for 6 hours. The mixture was diluted with chloroform and the solution was washed with saturated aqueous sodium hydrogen carbonate and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo. The crude product was purified by silica gel column (5% methanol in chloroform) to give N-[2-(4-dimethylaminobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylaminobenzamide (142 mg).

15

20

NMR (CDCl₃, δ) : 1.67-1.87 (4H, m), 2.25 (3H, s), 2.35 (6H, s), 2.46-2.61 (2H, m), 2.61 (3H, s), 3.32 (3H, s), 3.73 (3H, s), 3.81 (1H, m), 3.94 (1H, m), 6.53-6.64 (2H, m), 6.81-7.01 (3H, m), 7.24 (1H, t, J=8Hz), 7.46 (1H, br), 8.03 (1H, br), 8.50 (1H, br)

25

Example 101

The following compound was obtained according to similar manners to those of Preparation 25 and Example 38.

30

N-(2-Amino-4-methylbenzyl)-3-methoxy-N-methyl-4-(2-methylbenzimidazol-4-yl)carbamoylbenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 2.60 (3H, s), 2.79 (3H, s), 4.00 (3H, s), 4.03 (3H, s), 7.45-7.62 (3H, m), 7.69-7.81 (3H, m), 7.96 (1H, s), 8.03-8.11 (2H, m)

35

Example 102

A mixture of N-(2-carboxy-4-methylphenyl)-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-benzamide (350 mg), diphenylphosphorylazide (224 mg) and triethylamine (82.5 mg) in dioxane (10 ml) was heated at 80°C for 6 hours. After evaporation, the residual oil was subjected to silica gel column and the column was eluted with 5% methanol in chloroform. Object fractions were collected and evaporated in vacuo and the residue was solidified from chloroform to give N-(2-amino-4-methylphenyl)-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylamino-benzamide (280 mg).

NMR (DMSO-d₆, δ) : 2.10 (3H, s), 2.64 (3H, s), 3.11 (3H, s), 4.80 (3H, s), 5.41 (2H, s), 6.13 (1H, d, J=8Hz), 6.48-6.55 (2H, m), 7.02 (1H, d, J=8Hz), 7.09 (1H, s), 7.30 (1H, t, J=8Hz), 7.57 (1H, d, J=8Hz), 7.86 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

Example 103

The following compound was obtained according to a similar manner to that of Example 102.

N-(2-Amino-4-methylphenyl)-4-(1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methylbenzamide

NMR (DMSO-d₆, δ) : 2.08 (3H, s), 3.11 (3H, s), 3.76 (3H, s), 5.41 (2H, s), 6.14 (1H, d, J=8Hz), 6.47-6.53 (2H, m), 7.02 (1H, d, J=8Hz), 7.09 (1H, s), 7.40 (1H, t, J=8Hz), 7.81 (1H, d, J=8Hz), 7.97 (1H, d, J=8Hz), 8.35 (1H, d, J=8Hz), 8.52 (1H, s)

Example 104

A mixture of N-[2-(5-carboxypent-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)-aminomethyl-1H-benzimidazol-4-yl]carbonylamino-benzamide (850 mg), N-hydroxysuccinimide (145 mg) and N,N'-dicyclohexyl-

carbodiimide (260 mg) in tetrahydrofuran (20 ml) was stirred at ambient temperature overnight and the resulting insoluble urea was filtered off. To the filtrate was added lithium borohydride (55 mg) and the mixture was stirred at ambient temperature for 5 hours. The mixture was diluted with chloroform and the solution was washed successively with 1N hydrochloric acid, saturated aqueous sodium hydrogen carbonate and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo. The crude product was purified by silica gel column (5% methanol in chloroform) to give N-[2-(6-hydroxyhex-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (340 mg).

NMR (CDCl₃, δ) : 1.40-1.51 (2H, m), 1.48 (9H, s), 1.57-1.65 (2H, m), 1.71-1.82 (2H, m), 2.25 (3H, s), 3.35 (3H, s), 3.61-3.69 (2H, m), 3.73 (3H, s), 3.80 (1H, m), 3.91 (1H, m), 4.59 (2H, d, J=7Hz), 5.92 (1H, br t, J=7Hz), 6.56-6.63 (2H, m), 6.89-7.01 (3H, m), 7.29 (1H, t, J=8Hz), 7.55 (1H, d, J=8Hz), 7.96 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

Example 105

A mixture of N-[2-(6-hydroxyhex-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (210 mg), mesyl chloride (36.5 mg) and triethylamine (32.2 mg) in dichloromethane (10 ml) was stirred in an ice bath for 2 hours. The mixture was diluted with chloroform and the solution was washed successively with 1N hydrochloric acid, saturated aqueous sodium hydrogen carbonate and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo to give N-[2-(6-methanesulfonyloxyhex-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)aminomethyl-1H-benzimidazol-4-

233

yl]carbonylaminobenzamide (235 mg).

NMR (CDCl₃, δ) : 1.40-1.56 (4H, m), 1.49 (9H, s), 1.71-1.82 (2H, m), 2.28 (3H, s), 3.00 (3H, s), 3.36 (3H, s), 3.81 (3H, s), 3.81 (1H, m), 3.91 (1H, m), 4.22 (2H, t, J=7.5Hz), 4.58 (2H, d, J=7Hz), 5.61 (1H, br), 6.56-6.63 (2H, m), 6.90 (1H, d, J=8Hz), 6.95 (1H, d, J=8Hz), 7.00 (1H, s), 7.30 (1H, t, J=8Hz), 7.55 (1H, br), 8.10 (1H, br), 8.48 (1H, br)

10 Example 106

A mixture of N-[2-(6-methanesulfonyloxyhex-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-[2-(tert-butoxycarbonyl)-aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (235 mg) and potassium phthalimide (88.5 mg) in dimethyl sulfoxide (10 ml) was heated at 50°C for 6 hours. The mixture was diluted with ethyl acetate and the solution was washed with water and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo. The crude product was purified by silica gel column (1% methanol in chloroform) to give 3-methoxy-N-methyl-N-[2-(6-phthalimido-hex-1-yl)oxy-4-methyl]phenyl-4-[2-(tert-butoxycarbonyl)-aminomethyl-1H-benzimidazol-4-yl]carbonylaminobenzamide (224 mg).

NMR (CDCl₃, δ) : 1.36-1.54 (4H, m), 1.48 (9H, s), 1.64-1.87 (4H, m), 2.26 (3H, s), 2.61 (3H, s), 3.32 (3H, s), 3.69 (2H, t, J=7.5Hz), 3.79 (3H, s), 3.79 (1H, m), 3.90 (1H, m), 4.60 (1H, d, J=7Hz), 5.72 (1H, br), 6.54-6.63 (2H, m), 6.81-7.01 (3H, m), 7.31 (1H, t, J=8Hz), 7.60 (1H, d, J=8Hz), 7.70 (1H, m), 7.76 (1H, m), 7.80-7.88 (2H, m), 7.99 (1H, br), 8.41 (1H, d, J=8Hz)

Example 107

A solution of 4-[2-chloro-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-

methylnpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (70 mg) and N-methylnpiperazine (106 mg) was heated at 80°C for 2.5 hours. The excess of N-methylnpiperazine was evaporated in vacuo and the residue was purified by preparative thin-layer chromatography (chloroform:methanol = 10:1) to give 3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylnpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-methylnpiperazin-1-yl)-1H-benzimidazol-4-yl]carbonylaminobenzamide (66 mg).

NMR (CDCl₃, δ) : 1.43-1.60 (2H, m), 1.62-1.93 (4H, m),
2.23 (3H, s), 2.27 (3H, s), 2.30-2.43 (9H, m),
2.47-2.60 (4H, m), 3.34 (3H, s), 3.42-3.52 (2H, m),
3.54-3.72 (9H, m), 3.74-4.00 (2H, m), 6.50-6.67
(2H, m), 6.78-7.09 (4H, m), 7.21 (1H, m), 7.92 (1H,
m), 8.58 (1H, m)

Example 108

The following compounds were obtained according to a similar manner to that of Example 107.

- 1) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylnpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(morpholin-4-yl)-1H-benzimidazol-4-yl]carbonylaminobenzamide

NMR (CDCl₃, δ) : 1.45-1.60 (2H, m), 1.62-1.95 (4H, m),
2.25 (3H, s), 2.30 (3H, s), 2.32-2.44 (6H, m), 3.33
(3H, s), 3.41-3.52 (5H, m), 3.53-3.68 (6H, m),
3.75-4.00 (6H, m), 6.50-6.65 (2H, m), 6.73 (1H, s),
6.84 (1H, m), 6.91 (1H, m), 7.03 (1H, m), 7.23 (1H,
m), 7.93 (1H, m), 8.59 (1H, m)

- 2) 4-[(2-Dimethylamino-1H-benzimidazol-4-yl)carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylnpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.42-1.59 (2H, m), 1.61-1.96 (4H, m),
2.24 (3H, s), 2.29 (3H, s), 2.31-2.44 (6H, m), 3.12
(6H, s), 3.32 (3H, s), 3.38-3.53 (5H, m), 3.57-3.69

(2H, m), 3.72-4.00 (2H, m), 6.50-6.64 (2H, m), 6.70
(1H, s), 6.78-7.08 (3H, m), 7.16-7.29 (1H, m), 7.91
(1H, m), 8.60 (1H, m), 9.28 (1H, s)

- 5 3) 4-[2-[4-(Dimethylamino)piperidino]-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

10 NMR (CDCl₃, δ) : 1.29-2.00 (10H, m), 2.19-2.48 (18H, m), 2.67 (1H, m), 2.97-3.16 (2H, m), 3.33 (3H, s), 3.42-4.01 (9H, m), 4.18-4.46 (2H, m), 6.50-6.66 (2H, m), 6.78-7.08 (4H, m), 7.21 (1H, d, J=8Hz), 7.92 (1H, d, J=8Hz), 8.01 (1H, s), 8.59 (1H, m), 9.59 (1H, m)

15

- 4) 4-[[2-(Dimethylamino)amino-1H-benzimidazol-4-yl]-carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

20 NMR (CDCl₃, δ) : 1.21-1.36 (2H, m), 1.46-1.70 (4H, m), 2.12-2.38 (12H, m), 3.17 (3H, s), 3.25-3.54 (7H, m), 3.56-3.92 (8H, m), 6.39 (2H, s), 6.50 (1H, s), 6.60 (1H, d, J=8Hz), 6.69 (1H, d, J=8Hz), 6.72 (1H, s), 6.97 (1H, d, J=8Hz), 7.20 (1H, dd, J=8, 8Hz),
25 7.72 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 8.60 (1H, d, J=8Hz)

- 5) 4-[2-[(2-Aminoethyl)methylamino]-1H-benzimidazol-4-yl]-carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

30 NMR (CDCl₃, δ) : 1.43-1.59 (2H, m), 1.61-1.90 (4H, m), 2.26 (3H, s), 2.28 (3H, s), 2.30-2.44 (6H, m), 2.98-3.07 (2H, m), 3.29 (3H, s), 3.33 (3H, s),
35 3.41-3.52 (4H, m), 3.56-3.66 (2H, m), 3.70 (3H, s),

236

3.77-4.00 (2H, m), 6.52-6.66 (2H, m), 6.80-7.06
(4H, m), 7.22 (1H, m), 7.89 (1H, m), 8.56 (1H, m)

6) 4-[[2-[[2-(Dimethylamino)ethyl]amino]-1H-benzimidazol-4-
yl]carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-
(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide

NMR (CDCl₃, δ) : 1.43-1.58 (2H, m), 1.62-1.88 (4H, m),
2.20-2.42 (18H, m), 2.52-2.62 (2H, m), 3.23 (3H,
s), 3.42-3.56 (4H, m), 3.58-3.67 (2H, m), 3.72 (3H,
s), 3.77-4.00 (2H, m), 5.69 (1H, m), 6.53-6.65 (2H,
m), 6.87 (1H, d, J=8Hz), 6.91-7.00 (2H, m), 7.03
(1H, dd, J=8, 8Hz), 7.24 (1H, d, J=8Hz), 7.82 (1H,
d, J=8Hz), 8.50 (1H, d, J=8Hz)

Example 109

The following compound was obtained by using 3-methoxy-
N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]-4-[2-(1-benzyloxycarbonyl-4-
piperidyl)-1H-benzimidazol-4-yl]carbonylaminobenzamide as a
starting compound according to a similar manner to that of
Example 24.

3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-piperidyl)-1H-
benzimidazol-4-yl]carbonylaminobenzamide

NMR (DMSO-d₆, δ) : 1.37-1.50 (2H, m), 1.50-1.63 (2H,
m), 1.68-1.81 (2H, m), 1.85-2.04 (2H, m), 2.13 (3H,
s), 2.15-2.37 (11H, m), 2.83-2.98 (2H, m), 3.11-
3.49 (10H, m), 3.79 (3H, s), 3.83-4.05 (2H, m),
6.65 (1H, d, J=8Hz), 6.84 (1H, s), 6.92 (1H, d,
J=8Hz), 6.96 (1H, s), 7.03 (1H, d, J=8Hz), 7.33
(1H, t, J=8Hz), 7.70 (1H, d, J=8Hz), 7.91 (1H, d,
J=8Hz), 8.43 (1H, d, J=8Hz)

Example 110

4-(2-Formyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (158 mg),
5 hydroxylamine hydrochloride (25 mg), sodium acetate (30 mg) and ethanol (60% solution in water, 1.5 ml) were combined and the mixture was stirred at 60°C for 3 hours. After cooled to ambient temperature, the reaction mixture was concentrated in vacuo. The residue was dissolved in chloroform, washed with
10 water and brine and dried over magnesium sulfate and evaporated in vacuo. The residue was purified by basic preparative thin-layer chromatography (chloroform:methanol = 15:1) to give 4-(2-syn-hydroxyiminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
15 (38 mg) and 4-(2-anti-hydroxyiminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (89 mg).

20 Syn isomer :

NMR (DMSO-d₆, δ) : 1.37-1.49 (2H, m), 1.49-1.63 (2H, m), 1.67-1.80 (2H, m), 2.13 (3H, s), 2.15-2.35 (9H, m), 3.20 (3H, s), 3.35-3.44 (4H, m), 3.76-3.90 (4H, m), 3.90-4.01 (1H, m), 6.65 (1H, d, J=8Hz), 6.82
25 (1H, s), 6.90-6.98 (2H, m), 7.05 (1H, d, J=8Hz), 7.45 (1H, t, J=8Hz), 7.82 (1H, s), 7.87 (1H, d, J=8Hz), 8.00 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

Anti isomer :

30 NMR (DMSO-d₆, δ) : 1.37-1.49 (2H, m), 1.49-1.62 (2H, m), 1.67-1.81 (2H, m), 2.12 (3H, s), 2.16-2.35 (9H, m), 3.19 (3H, s), 3.35-3.45 (4H, m), 3.73-3.90 (4H, m), 3.90-4.02 (1H, m), 6.64 (1H, d, J=8Hz), 6.81 (1H, s), 6.89-6.98 (2H, m), 7.04 (1H, d, J=8Hz),
35 7.43 (1H, t, J=8Hz), 7.72 (1H, d, J=8Hz), 7.96 (1H,

d, J=8Hz), 8.23 (1H, s), 8.40 (1H, d, J=8Hz)

Example 111

To a solution of 4-[[2-cyanomethyl-1H-benzimidazol-4-yl]carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (224 mg) in ethanol (4 ml) and water (2 ml) was added hydroxylamine hydrochloride (93.5 mg) and sodium hydrogen carbonate (113 mg). The solution was heated at 90°C for 2 hours. After being concentrated in vacuo, aqueous sodium hydrogen carbonate was added to the residue and extracted with ethyl acetate. The extract was washed with brine and dried over sodium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography (10% methanol in chloroform) to give 4-[[2-[(2-amino-2-(hydroxyimino)ethyl)-1H-benzimidazol-4-yl]carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg).

NMR (CDCl₃, δ) : 1.37-1.58 (2H, m), 1.60-1.87 (4H, m), 2.22-2.49 (12H, m), 3.34 (3H, s), 3.39-3.51 (2H, m), 3.52-4.00 (9H, m), 5.49 (2H, br s), 6.51-6.66 (2H, m), 6.72 (1H, s), 6.81-7.01 (2H, m), 7.17 (1H, dd, J=8, 8Hz), 7.41 (1H, d, J=8Hz), 8.02 (1H, d, J=8Hz), 8.45 (1H, d, J=8Hz)

Example 112

To a solution of 4-[(2-formylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg) in pyridine (4.0 ml) was added hydroxylamine hydrochloride (21.3 mg) and the solution was stirred at ambient temperature for 1 hour. The resulting solution was concentrated in vacuo and the residue was diluted with chloroform. The organic layer was washed successively with water and brine, dried over magnesium sulfate and concentrated in vacuo to give 4-[(2-

hydroxyiminomethylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (203 mg).

NMR (CDCl₃, δ) : 1.48-1.60 (2H, m), 1.67-1.88 (4H, m),
5 2.27 (3H, s), 2.29 (3H, s), 2.32-2.47 (6H, m), 3.34
(3H, s), 3.48-3.56 (2H, m), 3.61-3.68 (2H, m), 3.70
(3H, s), 3.82-3.99 (2H, m), 6.57-6.64 (2H, m), 6.88
(1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.00 (1H, s),
7.12 (1H, s), 7.22 (1H, t, J=8Hz), 7.44 (1H, d,
10 J=8Hz), 7.50 (1H, d, J=8Hz), 8.14 (1H, s), 8.32
(1H, d, J=8Hz), 8.58-8.67 (2H, m), 9.32-9.38 (1H,
br s)

Example 113

15 To a solution of 2-methoxy-4-[N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
carbamoyl]benzoic acid (200 mg) in N,N-dimethylformamide (3
ml) at 0°C were added 1-ethyl-3-(3-dimethylaminopropyl)-
carbodiimide hydrochloride (97 mg), N-hydroxybenzotriazole
20 (79 mg) and 4-amino-2-[(4-methylpiperazin-1-yl)methyl]-1H-
benzimidazole (105 mg) and the mixture was stirred at ambient
temperature for 15 hours. The reaction mixture was poured
into saturated sodium bicarbonate aqueous solution and
extracted with chloroform. The organic layer was washed with
25 saturated sodium bicarbonate aqueous solution and brine,
dried over magnesium sulfate and evaporated in vacuo. The
residue was purified by preparative thin-layer chromatography
(chloroform:methanol = 10:1) to give 3-methoxy-N-methyl-N-[4-
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-
30 phenyl]-4-[2-[(4-methylpiperazin-1-yl)methyl]-1H-
benzimidazol-4-yl]carbamoylbenzamide (104 mg).

NMR (CDCl₃, δ) : 1.45-1.63 (2H, m), 1.63-1.79 (2H, m),
1.79-1.92 (2H, m), 2.28 (3H, s), 2.30-2.60 (12H,
m), 2.60-3.00 (8H, m), 3.35 (3H, s), 3.46-3.58 (2H,
35 m), 3.58-3.74 (2H, m), 3.82-4.06 (7H, m), 6.55-6.69

(2H, m), 6.88 (1H, d, J=8Hz), 7.00-7.13 (2H, m),
7.13-7.41 (2H, m), 8.06 (1H, d, J=8Hz), 8.32 (1H,
br peak)

5 Example 114

 The following compounds were obtained according to a
similar manner to that of Example 113.

- 1) 4-(1H-Benzimidazol-4-yl)carbamoyl-3-methoxy-N-methyl-N-
10 [4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
 yloxy]phenyl]benzamide
 NMR (CDCl₃, δ) : 1.49-1.60 (2H, m), 1.67-1.90 (4H, m),
 2.29 (3H, s), 2.36 (2H, t, J=8Hz), 2.38 (3H, s),
 2.42-2.53 (4H, m), 3.36 (3H, s), 3.52-3.58 (2H, m),
15 3.60 (3H, s), 3.65-3.73 (2H, m), 3.87-4.00 (2H, m),
 4.30-4.39 (2H, m), 6.59-6.68 (3H, m), 6.88 (1H, d,
 J=8Hz), 7.01-7.07 (2H, m), 7.16 (1H, t, J=8Hz),
 7.28 (1H, d, J=8Hz), 7.38 (1H, d, J=8Hz), 7.68 (1H,
 s)
20
- 2) 4-[(Naphthalen-1-yl)carbamoyl]-3-methoxy-N-methyl-N-[4-
 methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
 yloxy]phenyl]benzamide
 NMR (CDCl₃, δ) : 1.50-1.90 (6H, m), 2.28 (3H, s), 2.30
25 (3H, s), 2.32-2.41 (6H, m), 3.37 (3H, s), 3.46-3.51
 (2H, m), 3.59-3.67 (2H, m), 3.87-3.99 (2H, m), 4.02
 (3H, s), 6.58-6.64 (2H, m), 6.88 (1H, d, J=8Hz),
 7.02 (1H, d, J=8Hz), 7.16 (1H, s), 7.48-7.58 (3H,
 m), 7.68 (1H, d, J=8Hz), 7.86-7.95 (2H, m), 8.10
30 (1H, d, J=8Hz), 8.28 (1H, d, J=8Hz)
- 3) 3-(2-Carbamoylphenylcarbamoyl)-3-methoxy-N-methyl-N-[4-
 methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
 yloxy]phenyl]benzamide
35 NMR (CDCl₃, δ) : 1.48-1.59 (2H, m), 1.67-1.89 (4H, m),

2.27 (3H, s), 2.30 (3H, s), 2.32-2.43 (6H, m), 3.34 (3H, s), 3.47-3.52 (2H, m), 3.60-3.67 (2H, m), 3.82-3.98 (2H, m), 3.89 (3H, s), 5.84-5.95 (1H, br), 6.56-6.62 (2H, m), 6.87 (1H, d, J=8Hz), 6.95-7.08 (3H, m), 7.44 (1H, d, J=9Hz), 7.50 (1H, d, J=9Hz), 7.93 (1H, d, J=8Hz), 8.63 (1H, d, J=8Hz)

4) 4-(2-Methoxycarbonylphenylcarbamoyl)-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.49-1.58 (2H, m), 1.67-1.88 (4H, m), 2.27 (3H, s), 2.37 (2H, t, J=8Hz), 2.55 (3H, s), 2.77-2.88 (4H, m), 3.33 (3H, s), 3.62-3.68 (2H, m), 3.74-3.80 (2H, m), 3.87-3.98 (2H, m), 3.90 (3H, s), 3.92 (3H, s), 6.57-6.60 (2H, m), 6.86 (1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.02 (1H, s), 7.29-7.37 (1H, m), 7.60 (1H, d, J=8Hz), 7.82 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.01 (1H, d, J=8Hz), 8.82 (1H, d, J=8Hz)

20

5) 4-(2-Sulfamoylphenylcarbamoyl)-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.47-1.54 (2H, m), 1.64-1.84 (4H, m), 2.28 (3H, s), 2.34 (2H, t, J=8Hz), 2.50 (3H, s), 2.68-2.79 (4H, m), 3.28-3.42 (2H, br), 3.32 (3H, s), 3.58-3.64 (2H, m), 3.72-3.78 (2H, m), 3.82-3.97 (2H, m), 3.88 (3H, s), 6.57-6.61 (2H, m), 6.87 (1H, d, J=8Hz), 6.98 (1H, d, J=8Hz), 7.04 (1H, s), 7.29-7.39 (2H, m), 7.62 (1H, dd, J=2, 8Hz), 7.83 (2H, d, J=8Hz)

25

30

6) 4-[(Indol-4-yl)carbamoyl]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

35

- NMR (CDCl₃, δ) : 1.49-1.76 (4H, m), 1.80-1.90 (2H, m),
2.28 (3H, s), 2.30 (3H, s), 2.32-2.43 (6H, m), 3.38
(3H, s), 3.47-3.52 (2H, m), 3.60-3.68 (2H, m),
3.87-3.97 (2H, m), 4.00 (3H, s), 6.48 (1H, s),
5 6.58-6.64 (2H, m), 6.88 (1H, d, J=8Hz), 7.01 (1H,
d, J=8Hz), 7.11 (1H, s), 7.18-7.22 (2H, m), 7.23-
7.32 (1H, br), 8.06-8.11 (2H, m), 8.37-8.41 (1H, br
s)
- 10 7) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-
yl)carbamoyl-N-(4-methyl-2-nitrophenyl)benzamide
NMR (CDCl₃, δ) : 2.39 (3H, s), 2.59 (3H, s), 3.44 (3H,
s), 4.01 (3H, s), 6.87 (1H, d, J=8Hz), 7.07 (1H,
s), 7.11-7.23 (4H, m), 7.34 (1H, m), 7.62 (1H, s),
15 8.01 (1H, d, J=8Hz)
- 8) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-
yl)carbamoyl-N-[2-(4-tert-butoxycarbonylaminobut-1-
yl)oxy-4-methyl]phenylbenzamide
20 NMR (CDCl₃, δ) : 1.46 (9H, s), 1.61-1.74 (2H, m), 1.74-
1.88 (2H, m), 2.25 (3H, s), 2.65 (3H, s), 3.13-3.22
(2H, m), 3.34 (3H, s), 3.82-3.97 (2H, m), 3.93 (3H,
s), 4.67 (1H, br), 6.58-6.63 (2H, m), 6.90 (1H, d,
J=8Hz), 7.00-7.10 (2H, m), 7.17 (1H, t, J=8Hz),
25 7.30 (1H, m), 7.41 (1H, d, J=8Hz), 8.03 (1H, d,
J=8Hz)
- 9) 4-[N-[1-[(tert-Butyl)oxycarbonyl]benzamidazol-4-yl]-
carbamoyl]-N-[2-[4,4-dimethyl(2,5-oxazolinyl)]phenyl]-3-
30 methoxy-N-methylbenzamide
NMR (CDCl₃, δ) : 1.37 (3H, s), 1.38 (3H, s), 1.70 (9H,
s), 3.40 (3H, s), 3.90 (3H, s), 4.02-4.16 (2H, m),
7.02-7.41 (7H, m), 7.61 (1H, d, J=8Hz), 7.78 (1H,
d, J=8Hz), 8.08 (1H, dd, J=8, 8Hz), 8.44 (1H, d,
35 J=8Hz)

- 10) 4-[N-[2-[(Dimethylamino)methyl]-1H-benzimidazol-4-yl]-
carbamoyl]-N-[2-[4,4-dimethyl(2,5-oxazolinyl)]phenyl]-3-
methoxy-N-methylbenzamide
NMR (CDCl₃, δ) : 1.38 (3H, s), 1.39 (3H, s), 2.39 (6H,
5 s), 3.39 (3H, s), 3.81 (2H, s), 3.89 (3H, s), 4.07-
4.17 (2H, m), 7.04-7.39 (8H, m), 7.80 (1H, d,
J=8Hz), 8.10 (1H, d, J=8Hz)
- 11) 4-[N-[1-[(tert-Butyl)oxycarbonyl]benzimidazol-4-yl]-
10 carbamoyl]-3-methoxy-N-methyl-N-[2-(morpholin-4-yl)-
phenyl]benzamide
NMR (CDCl₃, δ) : 1.72 (9H, s), 2.32-2.46 (2H, m),
2.80-2.92 (2H, m), 3.53 (3H, s), 3.63-3.84 (4H, m),
3.91 (3H, s), 6.88 (1H, d, J=8Hz), 7.05 (1H, s),
15 7.09-7.25 (3H, m), 7.32 (1H, m), 7.38 (1H, dd, J=8,
8Hz), 7.64 (1H, d, J=8Hz), 8.10 (1H, d, J=8Hz),
8.37 (1H, s), 8.47 (1H, d, J=8Hz)
- 12) 4-[N-[1-[(tert-Butyl)oxycarbonyl]benzimidazol-4-yl]-
20 carbamoyl]-3-methoxy-N-methyl-N-[2-(1-pyrrolyl)phenyl]-
benzamide
NMR (CDCl₃, δ) : 1.72 (9H, s), 3.50 (3H, s), 3.96 (3H,
s), 6.22-6.31 (2H, m), 6.40-6.49 (2H, m), 6.54-6.69
(2H, m), 7.06-8.05 (8H, m), 8.38 (1H, s), 8.46 (1H,
25 d, J=8Hz)
- 13) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-
yl)carbamoyl]-N-(2-piperidinophenyl)benzamide
NMR (CDCl₃, δ) : 1.41-1.78 (6H, m), 2.26-2.41 (2H, m),
30 2.60 (3H, s), 2.70-2.86 (2H, m), 3.53 (3H, s),
3.72-3.93 (3H, m), 6.66-7.57 (9H, m), 8.00-8.39
(1H, m)
- 14) 4-[N-[1-[(tert-Butyl)oxycarbonyl]-2-methylbenzimidazol-
35 4-yl]carbamoyl]-3-methoxy-N-methyl-N-[2-(4-methyl-1-

piperazinyl)phenyl]benzamide

NMR (CDCl₃, δ) : 1.70 (9H, s), 2.38 (3H, s), 2.41-2.65 (6H, m), 2.81 (3H, s), 2.86-3.01 (2H, m), 3.52 (3H, s), 3.89 (3H, s), 6.89 (1H, d, J=8Hz), 6.98 (1H, s), 7.06-7.34 (5H, m), 7.57 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

15) 4-[N-[1-[(tert-Butyl)oxycarbonyl]-2-methylbenzimidazol-4-yl]carbamoyl]-3-methoxy-N-methyl-N-[2-(2,5-oxazolyl)-phenyl]benzamide

NMR (CDCl₃, δ) : 1.71 (9H, s), 2.81 (3H, s), 3.49 (3H, s), 3.90 (3H, s), 6.79-6.87 (2H, m), 7.17-7.46 (6H, m), 7.56 (1H, d, J=8Hz), 7.78 (1H, s), 7.88 (1H, m), 8.00 (1H, d, J=8Hz), 8.37 (1H, d, J=8Hz)

16) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)carbamoyl]-N-[2-(2,5-oxazolinyl)phenyl]benzamide

NMR (CDCl₃, δ) : 2.60 (3H, br s), 3.44 (3H, s), 3.79-3.96 (3H, m), 4.02-4.16 (2H, m), 4.29-4.49 (2H, m), 6.72 (1H, m), 6.98-7.59 (8H, m), 7.78 (1H, d, J=8Hz), 8.06 (1H, d, J=8Hz)

17) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)carbamoyl]-N-[2-(3H,4H,5H-2,6-oxazinyl)phenyl]-benzamide

NMR (CDCl₃, δ) : 1.93-2.09 (2H, m), 2.52-2.65 (3H, m), 3.43 (3H, s), 3.52-3.65 (2H, m), 3.80 and 3.88 (Total 3H, s), 4.27-4.42 (2H, m), 6.73 (1H, d, J=8Hz), 6.96-8.36 (10H, m)

18) N-[2-(1-Aza-3-oxaspiro[4.4]non-1-en-2-yl)phenyl]-3-methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)-carbamoyl]benzamide

NMR (CDCl₃, δ) : 1.62-1.78 (4H, m), 1.83-2.06 (4H, m), 2.60 (3H, s), 3.40 (3H, s), 3.86 (3H, br s), 4.18-

4.29 (2H, m), 6.72 (1H, m), 7.02-7.20 (4H, m),
7.22-7.40 (2H, m), 7.50 (1H, m), 7.77 (1H, m), 8.07
(1H, d, J=8Hz)

- 5 19) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-phthalimido-
methyl-1H-benzimidazol-4-yl)carbamoylbenzamide
NMR (CDCl₃, δ) : 1.47-1.67 (2H, m), 1.67-1.80 (2H, m),
1.80-1.94 (2H, m), 2.23-2.33 (6H, m), 2.33-2.45
10 (6H, m), 3.38 (3H, s), 3.45-3.55 (2H, m), 3.59-3.71
(2H, m), 3.81-4.09 (5H, m), 5.15 (2H, s), 6.54-6.68
(2H, m), 6.76-6.95 (1H, m), 6.98-7.17 (3H, m),
7.67-7.83 (2H, m), 7.83-7.95 (2H, m), 8.03-8.17
(1H, m), 8.33 (1H, d, J=8Hz), 9.75-9.83 (1H, m)
- 15 20) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(2-phthalimido-
ethyl)-1H-benzimidazol-4-yl]carbamoylbenzamide
NMR (CDCl₃, δ) : 1.46-1.63 (2H, m), 1.63-1.78 (2H, m),
20 1.78-1.91 (2H, m), 2.20-2.49 (9H, m), 2.49-2.71
(3H, m), 3.28-3.43 (5H, m), 3.43-4.04 (9H, m), 4.23
(2H, t, J=7.5Hz), 6.56-6.69 (2H, m), 6.81-7.11 (3H,
m), 7.18 (1H, br peak), 7.24-7.33 (1H, m), 7.48
(1H, br peak), 7.60-7.73 (2H, m), 7.78-7.87 (2H,
25 m), 8.03 (1H, br peak)
- 30 21) 4-(2-tert-Butyldiphenylsiloxymethyl-1H-benzimidazol-4-
yl)carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide
NMR (CDCl₃, δ) : 1.16 (9H, s), 1.44-1.64 (2H, m),
1.64-1.79 (2H, m), 1.79-1.94 (2H, m), 2.20-2.31
(6H, m), 2.31-2.43 (6H, m), 3.33 (3H x 1/2, s),
3.38 (3H x 1/2, s), 3.44-3.54 (2H, m), 3.57-3.69
35 (2H, m), 3.77-4.01 (5H, m), 4.99-5.06 (2H, m),

6.54-6.68 (2H, m), 6.74-6.94 (2H, m), 6.94-7.32
(3H, m), 7.32-7.59 (6H, m), 7.65-7.77 (4H, m),
8.00-8.14 (1H, m), 8.35 (1H x 1/2, d, J=8Hz), 9.34
(1H x 1/2, s)

5

- 22) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-[(tert-butoxy)-
carbonylamino]-1H-benzimidazol-4-yl]carbamoylbenzamide

10 NMR (DMSO-d₆, δ) : 1.36-1.50 (2H, m), 1.50-1.61 (2H,
m), 1.66 (9H, s), 1.70-1.80 (2H, m), 2.14 (3H, s),
2.17-2.37 (9H, m), 3.21 (3H, s), 3.38-3.46 (4H, m),
3.79-4.04 (5H, m), 6.65 (1H, d, J=8Hz), 6.80 (1H,
s), 6.96 (1H, t, J=8Hz), 7.02 (1H, d, J=8Hz), 7.05-
7.14 (2H, m), 7.30 (1H, d, J=8Hz), 7.34 (2H, br
15 peak), 7.90 (1H, d, J=9Hz), 8.11 (1H, d, J=8Hz)

- 23) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-[(methyl-
sulfonyl)amino]-1H-benzimidazol-4-yl]carbamoylbenzamide

20 NMR (DMSO-d₆, δ) : 1.36-1.50 (2H, m), 1.50-1.64 (2H,
m), 1.64-1.82 (2H, m), 2.15 (3H, s), 2.18-2.36 (9H,
m), 3.20 (3H, s), 3.37-3.46 (4H, m), 3.49 (3H, s),
3.77-4.03 (5H, m), 6.64 (1H, d, J=8Hz), 6.81 (1H,
s), 6.96-7.14 (6H, m), 7.23 (1H, d, J=8Hz), 7.90
25 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz)

- 24) 3-Methoxy-4-[2-methoxymethyl-1H-benzimidazol-4-yl]-
carbamoyl-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]benzamide

30 NMR (DMSO-d₆, δ) : 1.38-1.50 (2H, m), 1.50-1.64 (2H,
m), 1.64-1.83 (2H, m), 2.13 (3H, s), 2.16-2.38 (9H,
m), 3.20 (3H, s), 3.35-3.47 (7H, m), 3.82-4.01 (5H,
m), 4.68 (2H, s), 6.65 (1H, d, J=8Hz), 6.81 (1H,
s), 7.00-7.23 (5H, m), 7.91 (1H, d, J=8Hz), 8.11
35 (1H, d, J=8Hz)

- 25) 3-Methoxy-N-methyl-4-[2-methyl-1H-benzimidazol-4-yl]-
carbamoyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.38-1.51 (2H, m), 1.51-1.66 (2H,
5 m), 1.66-1.83 (2H, m), 2.15 (3H, s), 2.18-2.38 (9H,
m), 2.52 (3H, s), 3.22 (3H, s), 3.36-3.48 (4H, m),
3.80-4.05 (5H, m), 6.64 (1H, d, J=8Hz), 6.81 (1H,
s), 6.98-7.18 (5H, m), 7.90 (1H, d, J=8Hz), 8.06
(1H, d, J=8Hz)

10

- 26) 4-[1,2-Dimethyl-1H-benzimidazol-4-yl]carbamoyl-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.38-1.51 (2H, m), 1.51-1.65 (2H,
15 m), 1.70-1.82 (2H, m), 2.15 (3H, s), 2.18-2.39 (9H,
m), 2.56 (3H, s), 3.21 (3H, s), 3.37-3.50 (4H, m),
3.73 (3H, s), 3.81-4.05 (5H, m), 6.63 (1H, d,
J=8Hz), 6.81 (1H, s), 6.99-7.25 (5H, m), 7.90 (1H,
d, J=8Hz), 8.10 (1H, d, J=8Hz)

20

- 27) 4-[2-Ethyl-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-
methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.33-1.50 (5H, m), 1.50-1.66 (2H,
25 m), 1.66-1.83 (2H, m), 2.14 (3H, s), 2.16-2.28 (7H,
m), 2.28-2.38 (2H, m), 2.87 (2H, q, J=7.5Hz), 3.22
(3H, s), 3.37-3.46 (4H, m), 3.81-4.02 (5H, m), 6.64
(1H, d, J=8Hz), 6.81 (1H, s), 7.00-7.17 (5H, m),
7.93 (1H, d, J=8Hz), 8.09 (1H, d, J=8Hz)

30

- 28) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(n-propyl)-1H-
benzimidazol-4-yl]carbamoylbenzamide

NMR (DMSO-d₆, δ) : 1.00 (3H, t, J=7.5Hz), 1.37-1.50
35 (2H, m), 1.50-1.65 (2H, m), 1.70-1.94 (4H, m), 2.15

(3H, s), 2.18-2.38 (9H, m), 2.83 (2H, t, J=7.5Hz),
3.22 (3H, s), 3.36-3.45 (4H, m), 3.81-4.05 (5H, m),
6.63 (1H, d, J=8Hz), 6.80 (1H, s), 6.99-7.18 (5H,
m), 7.92 (1H, d, J=8Hz), 8.08 (1H, d, J=8Hz)

5

- 29) 4-[2-Isopropyl-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-
N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.35-1.50 (8H, m), 1.50-1.63 (2H,
10 m), 1.70-1.81 (2H, m), 2.13 (3H, s), 2.17-2.37 (9H,
m), 3.10-3.25 (4H, m), 3.36-3.46 (4H, m), 3.80-4.03
(5H, m), 6.64 (1H, d, J=8Hz), 6.80 (1H, s), 7.00-
7.18 (5H, m), 7.93 (1H, d, J=8Hz), 8.09 (1H, d,
J=8Hz)

15

- 30) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-trifluoromethyl-
1H-benzimidazol-4-yl]carbamoylbenzamide

NMR (DMSO-d₆, δ) : 1.35-1.50 (2H, m), 1.50-1.64 (2H,
20 m), 1.69-1.82 (2H, m), 2.14 (3H, s), 2.19-2.38 (9H,
m), 3.21 (3H, s), 3.37-3.49 (4H, m), 3.82-4.05 (5H,
m), 6.63 (1H, d, J=8Hz), 6.81 (1H, s), 6.99-7.13
(3H, m), 7.32-7.42 (2H, m), 7.84-7.96 (1H, m), 8.20
(1H, br peak)

25

- 31) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(3-pyridyl)-1H-
benzimidazol-4-yl]carbamoylbenzamide

NMR (DMSO-d₆, δ) : 1.39-1.50 (2H, m), 1.50-1.66 (2H,
30 m), 1.66-1.84 (2H, m), 2.12 (3H, s), 2.16-2.29 (7H,
m), 2.31 (2H, t, J=5Hz), 3.23 (3H, s), 3.36-3.50
(4H, m), 3.82-4.02 (2H, m), 4.07 (3H, s), 6.65 (1H,
d, J=8Hz), 6.82 (1H, s), 7.03 (1H, d, J=8Hz), 7.07-
7.16 (2H, m), 7.23 (1H, t, J=8Hz), 7.30 (1H, d,
35 J=8Hz), 7.64 (1H, dd, J=5, 8Hz), 7.95 (1H, d,

J=8Hz), 8.20 (1H, d, J=8Hz), 8.50 (1H, d, J=8Hz),
8.70 (1H, d, J=5Hz), 9.37 (1H, s)

32) 4-[2-(N,N-Dimethylcarbamoyl)-1H-benzimidazol-4-yl]-
5 carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide

NMR (CDCl₃, δ) : 1.44-1.66 (2H, m), 1.66-1.79 (2H,
m), 1.79-1.92 (2H, m), 2.26 (3H, s), 2.30 (3H, s),
10 2.32-2.45 (6H, m), 3.32 (3H, s), 3.36 (3H, s),
3.43-3.54 (2H, m), 3.59-3.69 (2H, m), 3.82-4.04
(8H, m), 6.54-6.67 (2H, m), 6.82-6.95 (2H, m),
6.95-7.06 (1H, m), 7.11 (1H, s), 7.19-7.41 (2H, m),
8.08 (1H, d, J=8Hz), 8.48 (1H, d, J=8Hz)

15

33) 3-Methoxy-4-[2-methoxy-1H-benzimidazol-4-yl]carbamoyl-N-
methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.35-1.50 (2H, m), 1.50-1.64 (2H,
20 m), 1.68-1.83 (2H, m), 2.13 (3H, s), 2.17-2.36 (9H,
m), 3.21 (3H, s), 3.36-3.45 (4H, m), 3.83-4.02 (5H,
m), 4.13 (3H, s), 6.64 (1H, d, J=8Hz), 6.81 (1H,
s), 6.97-7.13 (5H, m), 7.93 (1H, br peak), 8.06
(1H, br peak)

25

34) 4-[2-(N,N-Dimethylaminomethyl)-1H-benzimidazol-4-yl]-
carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methyl-
piperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (DMSO-d₆, δ) : 1.35-1.50 (2H, m), 1.50-1.63 (2H,
30 m), 1.68-1.82 (2H, m), 2.15 (3H, s), 2.20-2.38
(15H, m), 3.20 (3H, s), 3.36-3.46 (4H, m), 3.70
(3H, s), 3.80-4.03 (5H, m), 6.63 (1H, d, J=8Hz),
6.80 (1H, s), 6.99-7.19 (5H, m), 7.91 (1H, d,
J=8Hz), 8.08 (1H, d, J=8Hz)

35

- 35) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(1-imidazolyl)-methyl-1H-benzimidazol-4-yl]carbamoylbenzamide

5 NMR (CDCl₃, δ) : 1.45-1.62 (2H, m), 1.62-1.78 (2H, m),
1.78-1.90 (2H, m), 2.26 (3H, s), 2.35 (2H, t,
J=7.5Hz), 2.44 (3H, s), 2.54-2.72 (6H, m), 3.33
(3H, s), 3.57-3.69 (2H, m), 3.69-3.82 (2H, m),
3.82-4.03 (5H, m), 5.44 (2H, s), 6.53-6.63 (2H, m),
6.88 (1H, d, J=8Hz), 6.95-7.40 (5H, m), 7.74 (1H,
10 br peak), 8.80 (1H, d, J=8Hz)

- 36) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(morpholin-4-ylmethyl)-1H-benzimidazol-4-yl]carbamoylbenzamide

15 NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.65 (2H,
m), 1.69-1.82 (2H, m), 2.21 (3H, s), 2.31-2.53
(13H, m), 3.23 (3H, s), 3.27-3.36 (4H, m), 3.58-
3.67 (4H, m), 3.78 (2H, s), 3.82-4.01 (5H, m), 6.65
(1H, d, J=8Hz), 6.81 (1H, s), 7.00-7.23 (5H, m),
20 7.93 (1H, d, J=8Hz), 8.09 (1H, d, J=8Hz)

- 37) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(pyrrolidin-1-ylmethyl)-1H-benzimidazol-4-yl]carbamoylbenzamide

25 NMR (CDCl₃, δ) : 1.45-1.60 (2H, m), 1.65-1.78 (2H,
m), 1.78-1.91 (2H, m), 1.91-2.10 (4H, m), 2.27 (3H,
s), 2.30 (3H, s), 2.32-2.45 (6H, m), 2.95 (4H, br
peak), 3.35 (3H, s), 3.45-3.53 (2H, m), 3.58-3.68
(2H, m), 3.79-4.01 (5H, m), 4.21 (2H, br s), 6.52-
30 6.65 (2H, m), 6.86 (1H, d, J=8Hz), 7.00 (1H, d,
J=8Hz), 7.09 (1H, s), 7.18-7.29 (3H, m), 8.04 (1H,
d, J=8Hz)

- 38) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(piperidino-
- 35

methyl)-1H-benzimidazol-4-yl]carbamoylbenzamide

NMR (CDCl₃, δ) : 1.33-1.64 (10H, m), 1.69-1.83 (2H, m), 2.14 (3H, s), 2.16-2.36 (9H, m), 2.40-2.49 (4H, m), 3.21 (3H, s), 3.36-3.49 (4H, m), 3.71 (2H, s), 3.80-4.04 (5H, m), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.99-7.21 (5H, m), 7.91 (1H, d, J=8Hz), 8.09 (1H, d, J=8Hz)

- 39) 4-[2-[2-(Dimethylamino)ethyl]-1H-benzimidazol-4-yl]-carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.45-1.95 (8H, m), 2.26 (3H, s), 2.30 (3H, s), 2.32-2.45 (12H, m), 2.79 (2H, t, J=5Hz), 3.11 (2H, t, J=5Hz), 3.36 (3H, s), 3.45-3.55 (2H, m), 3.60-3.67 (2H, m), 3.80-4.02 (5H, m), 6.55-6.64 (2H, m), 6.88 (1H, d, J=8Hz), 7.00-7.10 (2H, m), 7.13-7.26 (2H, m), 7.93 (1H, br peak), 8.08 (1H, d, J=8Hz)

- 40) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-[2-(4-methylpiperazin-1-yl)ethyl]-1H-benzimidazol-4-yl]-carbamoylbenzamide

NMR (DMSO-d₆, δ) : 1.37-1.50 (2H, m), 1.50-1.64 (2H, m), 1.68-1.82 (2H, m), 2.13 (6H, s), 2.17-2.54 (17H, m), 2.81 (2H, t-like), 3.01 (2H, t-like), 3.21 (3H, s), 3.38-3.48 (4H, m), 3.81-4.02 (5H, m), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.89-7.21 (5H, m), 7.91 (1H, br peak), 8.07 (1H, br peak)

- 41) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-methylpiperazin-1-yl)-1H-benzimidazol-4-yl]carbamoylbenzamide

NMR (DMSO-d₆, δ) : 1.38-1.50 (2H, m), 1.50-1.64 (2H, m), 1.64-1.82 (2H, m), 2.15 (3H, s), 2.19-2.39

(12H, m), 2.39-2.53 (4H, m), 3.20 (3H, s), 3.36-3.47 (4H, m), 3.47-3.61 (4H, m), 3.82-4.03 (5H, m), 6.63 (1H, d, J=8Hz), 6.80 (1H, s), 6.83-6.98 (2H, m), 7.00 (1H, d, J=8Hz), 7.03-7.13 (2H, m), 7.91 (1H, d, J=8Hz), 8.00 (1H, d, J=8Hz)

42) 4-[2-Dimethylamino-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

10 NMR (DMSO-d₆, δ) : 1.38-1.50 (2H, m), 1.50-1.65 (2H, m), 1.65-1.82 (2H, m), 2.14 (3H, s), 2.18-2.38 (9H, m), 3.11 (6H, s), 3.21 (3H, s), 3.38-3.48 (4H, m), 3.82-4.01 (5H, m), 6.63 (1H, d, J=8Hz), 6.78-6.87 (2H, m), 6.91 (1H, d, J=8Hz), 7.01 (1H, d, J=8Hz),
15 7.03-7.13 (2H, m), 7.91 (1H, d, J=8Hz), 7.99 (1H, d, J=8Hz)

43) 4-[1-(tert-Butoxycarbonyl)-2-[[2-[N-(tert-butoxycarbonyl)-N-methylamino]ethyl]amino]-1H-benzimidazol-4-yl]-carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

20 NMR (CDCl₃, δ) : 1.42 (9H, br peak), 1.50-1.64 (2H, m), 1.64-1.78 (11H, m), 1.78-1.93 (2H, m), 2.25 (3H, s), 2.30 (3H, s), 2.32-2.43 (6H, m), 2.93 (3H, s),
25 3.35 (3H, s), 3.44-3.53 (2H, m), 3.58-3.68 (4H, m), 3.73 (2H, br peak), 3.80-4.01 (5H, m), 6.53-6.63 (2H, m), 6.85 (1H, d, J=8Hz), 6.96-7.09 (3H, m), 7.30 (1H, d, J=8Hz), 7.44 (1H, br peak), 8.06 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

30 44) 4-[1-(tert-Butoxycarbonyl)-2-[[2-((tert-butoxy)carbonylamino)ethyl]methylamino]-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

35 NMR (DMSO-d₆, δ) : 1.21 (9H, s), 1.38-1.50 (2H, m),

- 1.50-1.68 (11H, m), 1.68-1.81 (2H, m), 2.14 (3H, s), 2.18-2.38 (9H, m), 3.05 (3H, s), 3.15-3.28 (5H, m), 3.36-3.46 (4H, m), 3.46-3.56 (2H, m), 3.81-4.01 (5H, m), 6.63 (1H, d, J=8Hz), 6.70-6.82 (2H, m), 7.00-7.13 (4H, m), 7.34 (1H, d, J=8Hz), 7.92 (1H, d, J=8Hz), 8.20 (1H, d, J=8Hz)
- 5
- 45) 4-[2-(1-Imidazolyl)-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
- 10
- NMR (DMSO-d₆, δ) : 1.36-1.51 (2H, m), 1.51-1.64 (2H, m), 1.69-1.84 (2H, m), 2.13 (3H, s), 2.17-2.38 (9H, m), 3.23 (3H, s), 3.38-3.49 (4H, m), 3.82-4.08 (5H, m), 6.64 (1H, d, J=8Hz), 6.81 (1H, s), 7.03 (1H, d, J=8Hz), 7.08-7.14 (2H, m), 7.17-7.32 (3H, m), 7.92 (2H, br peak), 8.18 (1H, br peak), 8.49 (1H, s)
- 15
- 46) 4-[1-(tert-Butoxycarbonyl)-2-[[2-(dimethylamino)ethyl]-amino]-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide
- 20
- NMR (CDCl₃, δ) : 1.45-1.88 (13H, m), 1.88-1.91 (2H, m), 2.22-2.43 (18H, m), 2.64 (2H, t, J=5Hz), 3.35 (3H, s), 3.45-3.53 (2H, m), 3.60-3.73 (4H, m), 3.80-4.02 (5H, m), 6.54-6.64 (2H, m), 6.86 (1H, d, J=8Hz), 6.95-7.10 (3H, m), 7.34 (1H, d, J=8Hz), 7.44 (1H, br peak), 8.07 (1H, d, J=8Hz), 8.36 (1H, d, J=8Hz)
- 25
- 47) 4-[2-[[2-(Dimethylamino)ethyl]methylamino]-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
- 30
- NMR (DMSO-d₆, δ) : 1.35-1.50 (2H, m), 1.50-1.63 (2H, m), 1.68-1.82 (2H, m), 2.13 (3H, s), 2.17-2.37 (15H, m), 2.45-2.53 (2H, m), 3.11 (3H, s), 3.21
- 35

(3H, s), 3.37-3.48 (4H, m), 3.62 (2H, t, J=5Hz),
3.79-4.02 (5H, m), 6.64 (1H, d, J=8Hz), 6.76-6.86
(2H, m), 6.90 (1H, d, J=8Hz), 6.99-7.13 (3H, m),
7.91 (1H, d, J=8Hz), 7.99 (1H, d, J=8Hz)

5

- 48) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(1,2,4-triazol-1-yl)-1H-benzimidazol-4-yl]carbamoylbenzamide

NMR (DMSO-d₆, δ) : 1.38-1.50 (2H, m), 1.50-1.65 (2H, m), 1.69-1.82 (2H, m), 2.13 (3H, s), 2.17-2.37 (9H, m), 3.21 (3H, s), 3.36-3.45 (4H, m), 3.82-4.05 (5H, m), 6.64 (1H, d, J=8Hz), 6.81 (1H, s), 7.03 (1H, d, J=8Hz), 7.07-7.15 (2H, m), 7.18-7.31 (2H, m), 7.89 (1H, br peak), 8.17 (1H, br peak), 8.46 (1H, s), 9.40 (1H, s)

15

- 49) 4-[2-[(2-Methoxyethyl)amino]-1H-benzimidazol-4-yl]-carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl₃, δ) : 1.42-1.76 (4H, m), 1.76-1.91 (2H, m), 2.26 (3H, s), 2.30 (3H, s), 2.31-2.45 (6H, m), 3.34 (4H, s), 3.40 (3H, s), 3.45-3.55 (2H, m), 3.55-3.69 (6H, m), 3.79-4.02 (5H, m), 5.17 (1H, br peak), 6.56-6.65 (2H, m), 6.81 (1H, d, J=8Hz), 6.95-7.10 (3H, m), 7.10-7.35 (2H, m), 8.03 (1H, d, J=8Hz)

25

Example 115

The following compounds were obtained according to a similar manner to that of Example 38.

30

- 1) 4-(Imidazo[1,5-a]pyridine-1-carbonyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.63 (4H, m), 1.70-1.82 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=8Hz), 2.75 (3H,

35

s), 2.90-3.08 (3H, m), 3.19 (3H, s), 3.33-3.50 (3H, m), 3.75 (3H, s), 3.88-4.10 (3H, m), 4.40-4.48 (1H, m), 6.66 (1H, d, J=8Hz), 6.80-6.87 (1H, m), 6.90-7.07 (3H, m), 7.25 (1H, t, J=8Hz), 8.08 (1H, d, J=9Hz), 8.27 (1H, d, J=8Hz), 8.52-8.58 (2H, m), 9.52 (1H, s)

2) 4-[(2-Ethoxycarbonylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
NMR (DMSO-d₆, δ) : 1.21 (3H, t, J=8Hz), 1.41-1.63 (4H, m), 1.70-1.80 (2H, m), 2.23 (3H, s), 2.74 and 2.75 (Total 3H, s), 2.86-3.07 (3H, m), 3.19 (3H, s), 3.32-3.61 (6H, m), 3.65 (3H, s), 3.83-4.05 (3H, m), 4.11 (2H, q, J=8Hz), 4.37-4.49 (2H, m), 6.66 (1H, d, J=8Hz), 6.71 (1H, d, J=8Hz), 6.82 (1H, s), 6.88-6.93 (2H, m), 6.96 (1H, d, J=8Hz), 7.01-7.09 (2H, m), 7.77 (1H, d, J=8Hz), 8.99 (1H, s)

3) 4-[(2-Carbamoylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.70-1.80 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=8Hz), 2.74 (3H, s), 2.90-3.05 (3H, m), 3.19 (3H, s), 3.33-3.59 (6H, m), 3.65 (3H, s), 3.92-4.45 (6H, m), 6.67 (1H, d, J=8Hz), 6.80-6.94 (3H, m), 7.01-7.13 (3H, m), 7.19 (1H, s), 7.48 (1H, s), 7.76 (1H, d, J=8Hz), 9.03 (1H, s)

4) 4-[(2-Carbamoylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide hydrochloride
NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.71-1.81 (2H, m), 2.26 (3H, s), 2.40 (2H, t, J=8Hz), 2.74 (3H,

s), 2.88-3.05 (3H, m), 3.20 (3H, s), 3.28-3.48 (4H, m), 3.68 (3H, s), 3.83-4.02 (3H, m), 6.68 (1H, d, J=8Hz), 6.83 (1H, s), 6.91-6.98 (2H, m), 7.07 (1H, d, J=8Hz), 7.29 (1H, t, J=8Hz), 7.39-7.45 (1H, br s), 7.56 (1H, d, J=8Hz), 7.57 (1H, s), 7.61 (1H, d, J=8Hz), 7.91 (1H, d, J=8Hz), 8.06-8.11 (1H, br s), 9.12 (1H, s)

5) 4-[[2-(N-Methylcarbamoyl)indol-4-yl]carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride
NMR (CDCl₃, δ) : 1.42-1.62 (4H, m) 1.71-1.80 (2H, m), 2.24 (3H, s), 2.40 (2H, t, J=8Hz), 2.74 (3H, s), 2.79 and 2.81 (Total 3H, s), 2.88-3.02 (3H, m), 3.19 (3H, s), 3.27-3.42 (4H, m), 3.67 (3H, s), 3.86-4.10 (3H, m), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.91-6.98 (2H, m), 7.06 (1H, d, J=8Hz), 7.28 (1H, t, J=8Hz), 7.52 (1H, s), 7.54 (1H, d, J=8Hz), 7.62 (1H, d, J=8Hz), 7.90 (1H, d, J=8Hz), 8.57-8.62 (1H, m), 9.11 (1H, s)

6) 4-[(Indolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
NMR (DMSO-d₆, δ) : 1.40-1.62 (4H, m), 1.70-1.80 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=8Hz), 2.74 (3H, s), 2.87-3.06 (3H, m), 3.19 (3H, s), 3.30-3.51 (5H, m), 3.63 (3H, s), 3.67 (2H, t, J=8Hz), 3.81-4.10 (4H, m), 4.39-4.48 (1H, m), 6.64 (1H, d, J=8Hz), 6.82 (1H, s), 6.88-6.93 (2H, m), 7.04 (1H, d, J=8Hz), 7.40-7.47 (2H, m), 7.61-7.72 (2H, m), 9.40 (1H, s)

7) 4-[(2-Hydroxymethylindolin-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-

yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.62 (4H, m), 1.69-1.80 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=8Hz), 2.73 and 2.74 (Total 3H, s), 2.90-3.12 (4H, m), 3.19 (3H, s), 3.30-3.60 (6H, m), 3.64 (3H, s), 3.82-4.12 (5H, m), 4.39-4.49 (1H, m), 6.65 (1H, d, J=9Hz), 6.82 (1H, s), 6.88-6.93 (2H, m), 7.03 (1H, d, J=9Hz), 7.15 (1H, d, J=7Hz), 7.30 (1H, t, J=9Hz), 7.40 (1H, d, J=7Hz), 7.72 (1H, d, J=9Hz), 9.24 (1H, s)

10

- 8) 4-[(2-Aminomethylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.70-1.81 (2H, m), 2.24 (3H, s), 2.39 (2H, t, J=8Hz), 2.70 (3H, s), 2.98-3.12 (3H, m), 3.19 (3H, s), 3.28-3.46 (4H, m), 3.69 (3H, s), 3.84-4.02 (3H, m), 4.23 (2H, s), 6.67 (1H, d, J=8Hz), 6.84 (1H, s), 6.90-6.98 (3H, m), 7.05 (1H, d, J=8Hz), 7.22 (1H, t, J=9Hz), 7.56 (1H, d, J=9Hz), 7.62 (1H, d, J=9Hz), 7.93 (1H, d, J=8Hz), 8.68-8.77 (2H, br), 9.08 (1H, s)

15

20

- 9) 4-[(2-Methylindol-4-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.70-1.80 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=8Hz), 2.43 (3H, s), 2.76 (3H, s), 2.88-3.02 (3H, m), 3.19 (3H, s), 3.28-3.43 (4H, m), 3.70 (3H, s), 3.86-4.07 (3H, m), 6.53 (1H, s), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.90-6.97 (2H, m), 7.06 (1H, d, J=9Hz), 7.11 (1H, d, J=9Hz), 7.49 (2H, d, J=9Hz), 8.02 (1H, d, J=8Hz), 8.98 (1H, s)

25

30

- 10) 4-[(Indolin-6-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-

35

methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

5 NMR (DMSO-d₆, δ) : 1.40-1.63 (4H, m), 1.69-1.80 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=8Hz), 2.73 (3H, s), 2.90-3.07 (2H, m), 3.19 (3H, s), 3.20 (2H, t, J=8Hz), 3.32-3.50 (3H, m), 3.63 (3H, s), 3.70 (2H, t, J=8Hz), 3.82-4.12 (4H, m), 4.38-4.48 (1H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.88-6.95 (2H, m), 7.04 (1H, d, J=8Hz), 7.48 (1H, d, J=8Hz), 7.53 (1H, d, J=8Hz), 7.62 (1H, d, J=8Hz), 7.72 (1H, s), 7.78 (1H, d, J=8Hz), 9.46 (1H, s)

11) 4-[(Indol-6-yl)carbonyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride
15 NMR (DMSO-d₆, δ) : 1.40-1.63 (4H, m), 1.70-1.81 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=8Hz), 2.73 and 2.74 (Total 3H, s), 2.83-3.06 (3H, m), 3.18 (3H, s), 3.31-3.46 (3H, m), 3.67 (3H, s), 3.82-4.12 (3H, m), 4.39-4.49 (1H, m), 6.50 (1H, s), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.89-6.96 (2H, m), 7.05 (1H, d, J=8Hz), 7.51-7.58 (2H, m), 7.62 (1H, d, J=8Hz), 7.81 (1H, d, J=8Hz), 8.00 (1H, s), 9.18 (1H, s)

25 12) 4-(1H-Benzimidazol-4-yl)carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
ESI-MASS : 627.5 (M+H)

30 13) 4-[(Naphthalen-1-yl)carbamoyl]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride
NMR (DMSO-d₆, δ) : 1.42-1.66 (4H, m), 1.73-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=8Hz), 2.73 (3H, s), 2.86-3.08 (3H, m), 3.22 (3H, s), 3.31-3.48 (3H, m), 7.1-7.8 (8H, m), 8.1-8.4 (2H, m), 8.6-8.8 (2H, m), 9.1-9.3 (2H, m)

m), 3.87 (3H, s), 3.90-4.02 (3H, m), 4.38-4.50 (1H, m), 6.68 (1H, d, J=8Hz), 6.82 (1H, s), 6.99 (1H, d, J=8Hz), 7.08 (1H, s), 7.12 (1H, d, J=8Hz), 7.49-7.67 (4H, m), 7.77-7.86 (2H, m), 7.93-8.02 (2H, m)

5

- 14) 4-(2-Carbamoylphenylcarbamoyl)-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

10 NMR (DMSO-d₆, δ) : 1.40-1.63 (4H, m), 1.70-1.81 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=8Hz), 2.73 (3H, s), 2.88-3.05 (3H, m), 3.20 (3H, s), 3.30-3.50 (4H, m), 3.80 (3H, s), 3.83-4.00 (3H, m), 6.65 (1H, d, J=8Hz), 6.80 (1H, s), 6.95-7.00 (2H, m), 7.08-7.18 (2H, m), 7.48 (1H, t, J=8Hz), 7.64 (1H, s), 7.69 (1H, d, J=8Hz), 7.77 (1H, d, J=8Hz), 8.18 (1H, s), 8.53 (1H, d, J=8Hz)

15

- 15) 4-(2-Methoxycarbonylphenylcarbamoyl)-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

20

25 NMR (DMSO-d₆, δ) : 1.41-1.62 (4H, m), 1.71-1.80 (2H, m), 2.21 (3H, s), 2.40 (2H, t, J=8Hz), 2.74-2.78 (3H, br s), 2.88-3.02 (3H, m), 3.20 (3H, s), 3.32-3.52 (3H, m), 3.85 (3H, s), 3.88 (3H, s), 3.90-4.12 (3H, m), 4.40-4.48 (1H, m), 6.66 (1H, d, J=8Hz), 6.81 (1H, s), 6.98 (1H, d, J=8Hz), 7.03 (1H, s), 7.09 (1H, d, J=8Hz), 7.21 (1H, t, J=8Hz), 7.63 (1H, t, J=8Hz), 7.72 (1H, d, J=8Hz), 7.79 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 8.67 (1H, d, J=8Hz)

25

30

- 16) 4-[2-(N,N-Dimethylcarbamoyl)phenylcarbamoyl]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

35 NMR (DMSO-d₆, δ) : 1.41-1.63 (4H, m), 1.72-1.81 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=8Hz), 2.76 (3H,

s), 2.83-3.08 (3H, m), 2.89 (3H, s), 3.05 (3H, s),
3.20 (3H, s), 3.25-3.45 (4H, m), 3.83 (3H, s),
3.86-4.00 (3H, m), 6.65 (1H, d, J=8Hz), 6.81 (1H,
s), 6.97-7.02 (2H, m), 7.09 (1H, d, J=8Hz), 7.18
5 (1H, t, J=8Hz), 7.35 (1H, d, J=8Hz), 7.42 (1H, t,
J=8Hz), 7.82 (1H, d, J=8Hz), 8.27 (1H, d, J=8Hz)

17) 4-(2-Sulfamoylphenylcarbamoyl)-3-methoxy-N-methyl-N-[4-
methoxy-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
10 yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.62 (4H, m), 1.69-1.79 (2H,
m), 2.22 (3H, s), 2.38 (2H, t, J=8Hz), 2.77 (3H,
s), 2.86-3.04 (3H, m), 3.19 (3H, s), 3.35-3.58 (5H,
m), 3.61 (3H, s), 3.82-4.12 (3H, m), 4.39-4.49 (1H,
15 m), 6.63 (1H, d, J=8Hz), 6.80 (1H, s), 6.86-6.92
(2H, m), 7.06 (1H, d, J=8Hz), 7.39-7.47 (2H, m),
7.55 (1H, t, J=8Hz), 7.72 (1H, d, J=8Hz), 7.98 (1H,
d, J=8Hz)

20 18) 4-[(Indol-4-yl)carbamoyl]-3-methoxy-N-methyl-N-[4-
methoxy-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d₆, δ) : 1.42-1.63 (4H, m), 1.72-1.81 (2H,
m), 2.22 (3H, s), 2.40 (2H, t, J=8Hz), 2.75 (3H,
25 s), 2.90-3.03 (3H, m), 3.21 (3H, s), 3.30-3.44 (4H,
m), 3.88 (3H, s), 3.90-4.00 (3H, m), 6.50 (1H, s),
6.67 (1H, d, J=8Hz), 6.82 (1H, s), 6.99 (1H, d,
J=8Hz), 7.04-7.08 (2H, m), 7.12 (1H, d, J=8Hz),
7.18 (1H, d, J=8Hz), 7.34 (1H, t, J=3Hz), 7.69 (1H,
30 d, J=8Hz), 7.78 (1H, d, J=8Hz)

19) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-(quinolin-8-yl)-
carbonylaminobenzamide dihydrochloride

35 NMR (DMSO-d₆, δ) : 1.40-1.51 (2H, m), 1.53-1.63 (2H,

- m), 1.70-1.81 (2H, m), 2.23 (3H, s), 2.34-2.42 (2H, m), 2.50 (3H, s), 2.80-3.07 (4H, m), 3.20 (3H, s), 3.31-3.55 (4H, m), 3.83 (3H, s), 4.08 (1H, m), 4.45 (1H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.90-6.99 (2H, m), 7.05 (1H, d, J=8Hz), 7.71-7.85 (2H, m), 8.29 (1H, d, J=8Hz), 8.42 (1H, d, J=8Hz), 8.63 (1H, d, J=8Hz), 8.72 (1H, d, J=8Hz), 9.14 (1H, m)
- 5
- 20) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(1,2,3,4-tetrahydroquinolin-8-yl)carbonylaminobenzamide dihydrochloride
- 10
- NMR (DMSO-d₆, δ) : 1.39-1.50 (2H, m), 1.52-1.61 (2H, m), 1.68-1.83 (4H, m), 2.23 (3H, s), 2.38 (2H, t, J=7.5Hz), 2.49 (3H, s), 2.80-3.07 (4H, m), 3.18 (3H, s), 3.23-3.50 (6H, m), 3.60 (3H, s), 3.76-4.11 (3H, m), 4.42 (1H, m), 6.50 (1H, t, J=8Hz), 6.64 (1H, d, J=8Hz), 6.82 (1H, s), 6.86-6.92 (2H, m), 7.01-7.04 (2H, m), 7.43 (1H, d, J=8Hz), 7.61 (1H, d, J=8Hz), 9.08 (1H, s)
- 15
- 20
- 21) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(quinoxalin-5-yl)-carbonylaminobenzamide dihydrochloride
- NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.64 (2H, m), 1.70-1.81 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.73 (1H, s), 2.78-3.10 (3H, m), 3.20 (3H, s), 3.30-3.57 (3H, m), 3.65 (3H, s), 3.82-4.11 (3H, m), 4.41 (1H, m), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.88-6.95 (2H, m), 7.04 (2H, d, J=8Hz), 7.64 (1H, d, J=8Hz), 8.20 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz), 8.60 (1H, s), 9.04 (2H, s), 9.84 (1H, s)
- 25
- 30
- 22) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonylphenylmethoxy]phenyl]benzamide
- 35

trihydrochloride

- 23) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[3-(4-methylpiperazin-1-yl)carbonylprop-1-yloxy]phenyl]benzamide trihydrochloride
- 24) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylbut-1-yloxy]phenyl]benzamide trihydrochloride
- NMR (DMSO-d₆, δ) : 1.60-1.83 (4H, m), 2.22 (3H, s), 2.46 (2H, t, J=7.5Hz), 2.72 (3 x 1/2H, s), 2.74 (3 x 1/2H, s), 2.81-3.09 (3H, m), 3.20 (3H, s), 3.32-3.56 (3H, m), 3.70-4.12 (3H, m), 3.77 (3H, s), 4.40-4.50 (3H, m), 6.67 (1H, d, J=8Hz), 6.82 (1H, s), 6.94-6.99 (2H, m), 7.08 (1H, d, J=8Hz), 7.43 (1H, t, J=8Hz), 7.84 (1H, d, J=8Hz), 7.98 (1H, d, J=8Hz), 8.29 (1H, d, J=8Hz), 8.88-9.00 (3H, br)
- 25) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-[5-[N-(2-dimethylaminoeth-1-yl)-N-methylamino-carbonyl]-pent-1-yloxy]-4-methylphenyl]-3-methoxy-N-methylbenzamide trihydrochloride
- NMR (DMSO-d₆, δ) : 1.40-1.51 (2H, m), 1.51-1.63 (2H, m), 1.69-1.80 (2H, m), 2.22 (3H, s), 2.33 (2H, t, J=7.5Hz), 2.74 (3H, s), 2.76 (3H, s), 2.99 (3H, s), 3.17 (2H, m), 3.19 (3H, s), 3.62 (2H, t, J=7.5Hz), 3.77 (3H, s), 3.86 (1H, m), 3.97 (1H, m), 4.46 (2H, m), 6.66 (1H, d, J=8Hz), 6.84 (1H, s), 6.91-6.97 (2H, m), 7.02 (1H, d, J=8Hz), 7.43 (1H, t, J=8Hz), 7.83 (1H, d, J=8Hz), 7.97 (1H, d, J=8Hz), 8.29 (1H, d, J=8Hz), 8.94 (2H, br)
- 26) N-[2-(5-Carbamoylpent-1-yloxy)-4-methylphenyl]-4-(2-aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-

methoxy-N-methylbenzamide dihydrochloride

- 27) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbonylamino-N-[2-[5-(2,2-dimethylhydrazino)carbonylpent-1-yloxy]-4-methylphenyl]-3-methoxy-N-methylcarbonylaminobenzamide trihydrochloride

5
10
15
NMR (DMSO-d₆, δ) : 1.39-1.51 (2H, m), 1.58-1.69 (2H, m), 1.69-1.81 (2H, m), 2.21 (3H, s), 2.25 (2H, t, J=7.5Hz), 3.02 (6H, sx2), 3.19 (3H, s), 3.76 (3H, s), 3.89 (1H, m), 3.99 (1H, m), 4.45 (2H, m), 6.67 (1H, d, J=8Hz), 6.83 (1H, s), 6.91-6.96 (2H, m), 7.03 (1H, d, J=8Hz), 7.42 (1H, t, J=8Hz), 7.84 (1H, d, J=8Hz), 7.97 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz), 8.82-8.98 (3H, br)

- 28) N-[2-(4-Aminobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylaminobenzamide dihydrochloride

20
25
NMR (DMSO-d₆, δ) : 1.69-1.86 (2H, m), 2.21 (3H, s), 2.76 (3H, s), 2.81-2.92 (2H, m), 3.21 (3H, s), 3.69 (3H, s), 3.89 (1H, m), 4.01 (1H, m), 6.64 (1H, d, J=8Hz), 6.83 (1H, s), 6.90-6.98 (2H, m), 7.02 (1H, d, J=8Hz), 7.51 (1H, t, J=8Hz), 7.88 (1H, d, J=8Hz), 8.01-8.18 (4H, m)

- 29) N-[2-(4-Dimethylaminobut-1-yloxy)-4-methylphenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)carbonylaminobenzamide dihydrochloride

30
35
NMR (DMSO-d₆, δ) : 1.73-1.89 (4H, m), 2.24 (3H, s), 2.69 (3H, s), 2.70 (3H, s), 2.73 (3H, s), 3.07-3.16 (2H, m), 3.23 (3H, s), 3.67 (3H, s), 3.88 (1H, m), 3.99 (1H, m), 6.68 (1H, d, J=8Hz), 6.83 (1H, s), 6.93 (1H, s), 6.98 (1H, d, J=8Hz), 7.09 (1H, d, J=8Hz), 7.52 (1H, t, J=8Hz), 7.83 (1H, br), 7.90 (1H, d, J=8Hz), 8.06 (1H, d, J=8Hz)

- 30) N-[2-(4-Aminobut-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbamoylbenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.70-1.88 (4H, m), 2.23 (3H, s),
2.78 (3H, s), 2.82-2.92 (2H, m), 3.23 (3H, s), 3.73
(3H, s), 3.87-6.08 (2H, m), 6.67 (1H, d, J=8Hz),
6.84 (1H, s), 6.97-7.02 (2H, m), 7.09 (1H, d,
J=8Hz), 7.45 (1H, t, J=8Hz), 7.55 (1H, d, J=8Hz),
7.59-7.68 (2H, m), 8.08 (2H, br)

- 31) N-[2-(5-Aminopent-1-yl)oxy-4-methyl]phenyl-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-carbonylaminobenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.41-1.51 (2H, m), 1.58-1.69 (2H,
m), 1.69-1.81 (2H, m), 2.23 (3H, s), 2.71 (3H, s),
2.71-2.82 (2H, m), 3.20 (3H, s), 3.70 (3H, s), 3.88
(1H, m), 3.98 (1H, m), 6.67 (1H, d, J=8Hz), 6.82
(1H, s), 6.91 (1H, s), 6.93 (1H, d, J=8Hz), 7.07
(1H, d, J=8Hz), 7.49 (1H, m), 7.85 (1H, d, J=8Hz),
7.92-8.10 (3H, m)

- 32) N-[2-(6-Aminohex-1-yl)oxy-4-methyl]phenyl-4-(2-aminomethyl-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methylbenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.34-1.50 (4H, m), 1.52-1.65 (2H,
m), 1.70-1.79 (2H, m), 2.21 (3H, s), 2.71-2.80 (2H,
m), 3.17 (3H, s), 3.75 (3H, s), 3.80-4.20 (2H, m),
4.41 (2H, m), 6.67 (1H, d, J=8Hz), 7.82 (1H, s),
6.90-6.98 (2H, m), 7.06 (1H, d, J=8Hz), 7.40 (1H,
t, J=8Hz), 7.83 (1H, d, J=8Hz), 7.90-8.03 (3H, m),
8.30 (1H, m), 8.82-8.97 (2H, br)

- 33) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-methylpiperazin-1-yl)-1H-benzimidazol-4-yl]-

carbonylaminobenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.64 (4H, m), 1.65-1.82 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=8Hz), 2.68-2.84 (6H, m), 2.85-4.72 (24H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.87-7.00 (2H, m), 7.03 (1H, d, J=8Hz), 7.10 (1H, dd, J=8, 8Hz), 7.45 (1H, d, J=8Hz), 7.74 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

34) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(morpholin-4-yl)-1H-benzimidazol-4-yl]carbonylaminobenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.66 (4H, m), 1.68-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.69-2.78 (3H, m), 2.80-3.11 (2H, m), 3.20 (3H, s), 3.28-3.59 (3H, m), 3.60-4.18 (15H, m), 4.42 (1H, m), 6.64 (1H, d, J=8Hz), 6.82 (1H, s), 6.87-6.97 (2H, m), 7.02 (1H, d, J=8Hz), 7.11 (1H, dd, J=8, 8Hz), 7.44 (1H, d, J=8Hz), 7.72 (1H, d, J=8Hz), 8.26 (1H, m), 8.50 (1H, m)

35) 4-[(2-Dimethylamino-1H-benzimidazol-4-yl)carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.37-1.66 (4H, m), 1.68-1.82 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.72 and 2.74 (Total 3H, s), 2.80-3.11 (3H, m), 3.18 (3H, s), 3.21 (6H, s), 3.30-4.18 (9H, m), 4.44 (1H, m), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.87-6.97 (2H, m), 6.97-7.09 (2H, m), 7.35 (1H, d, J=8Hz), 7.68 (1H, d, J=8Hz), 8.37 (1H, m)

36) 4-[2-[4-(Dimethylamino)piperidino]-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-

methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.32-1.88 (10H, m), 1.99-4.60 (35H,
m), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.87-7.00
(2H, m), 7.01-7.12 (2H, m), 7.40 (1H, d, J=8Hz),
7.71 (1H, d, J=8Hz), 8.00 (1H, s), 8.41 (1H, m)

37) 4-[[2-(Dimethylamino)amino-1H-benzimidazol-4-yl]-
carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.37-1.66 (4H, m), 1.67-1.84 (2H,
m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.73 (3H,
s), 2.80-3.11 (3H, m), 3.19 (3H, s), 3.28-4.15
(15H, m), 4.43 (1H, m), 6.66 (1H, d, J=8Hz), 6.83
(1H, s), 6.88-7.12 (3H, m), 7.59 (1H, dd, J=8,
8Hz), 7.92 (1H, d, J=8Hz), 8.10 (1H, d, J=8Hz),
8.41 (1H, m)

38) 4-[[2-Cyanomethyl-1H-benzimidazol-4-yl]carbonylamino]-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.36-1.66 (4H, m), 1.67-1.83 (2H,
m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.72 and
2.73 (Total 3H, s), 2.80-3.11 (3H, m), 3.20 (3H,
s), 3.28-3.58 (3H, m), 3.73-4.15 (6H, m), 4.43 (1H,
m), 4.59 (2H, s), 6.66 (1H, d, J=8Hz), 6.83 (1H,
s), 6.88-7.00 (2H, m), 7.04 (1H, d, J=8Hz), 7.41
(1H, dd, J=8, 8Hz), 7.79 (1H, d, J=8Hz), 7.97 (1H,
d, J=8Hz), 8.39 (1H, d, J=8Hz)

39) 4-[[2-[(2-Amino-2-(hydroxyimino)ethyl)-1H-benzimidazol-
4-yl]carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-
(4-methylpiperazin-1-yl)carbonylpent-1-
yloxy]phenyl]benzamide trihydrochloride

- 5 NMR (DMSO-d₆, δ) : 1.37-1.65 (4H, m), 1.66-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7Hz), 2.72 and 2.73 (Total 3H, s), 2.78-3.15 (3H, m), 3.19 (3H, s), 3.24-3.59 (3H, m), 3.77 (3H, s), 3.81-4.17 (3H, m), 4.32 (2H, s), 4.43 (1H, m), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.88-6.99 (2H, m), 7.05 (1H, d, J=8Hz), 7.40 (1H, dd, J=8, 8Hz), 7.80 (1H, d, J=8Hz), 7.96 (1H, d, J=8Hz), 8.29 (1H, d, J=8Hz)
- 10 40) 4-[[2-[(2-Aminoethyl)methylamino]-1H-benzimidazol-4-yl]-carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide trihydrochloride
- 15 NMR (DMSO-d₆, δ) : 1.37-1.65 (4H, m), 1.66-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7Hz), 2.70 and 2.72 (Total 3H, s), 2.80-3.58 (14H, m), 3.70 (3H, s), 3.77-4.15 (5H, m), 4.43 (1H, m), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.87-6.99 (2H, m), 7.04 (1H, d, J=8Hz), 7.12 (1H, dd, J=8, 8Hz), 7.44 (1H, d, J=8Hz), 7.71 (1H, d, J=8Hz), 8.18 (1H, br), 8.38 (2H, br)
- 20
- 41) 4-[[2-[[2-(Dimethylamino)ethyl]amino]-1H-benzimidazol-4-yl]carbonylamino]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide trihydrochloride
- 25 NMR (DMSO-d₆, δ) : 1.38-1.65 (4H, m), 1.68-1.84 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=7Hz), 2.65-4.27 (28H, m), 4.42 (1H, m), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.88-6.99 (2H, m), 7.05 (1H, d, J=8Hz), 7.27 (1H, m), 7.55 (1H, d, J=8Hz), 7.83 (1H, d, J=8Hz)
- 30
- 42) 3-Methoxy-4-(2-mercaptomethyl-1H-benzimidazol-4-yl)-carbonylamino-N-methyl-N-[4-methyl-2-[5-(4-
- 35

methylnpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.66 (4H, m), 1.68-1.84 (2H,
m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.70-2.77
5 (3H, m), 2.79-3.12 (3H, m), 3.20 (3H, s), 3.30-4.18
(11H, m), 4.36-4.51 (1H, m), 6.66 (1H, d, J=8Hz),
6.82 (1H, s), 6.88-7.00 (2H, m), 7.05 (1H, d,
J=8Hz), 7.44 (1H, dd, J=8, 8Hz), 7.81 (1H, d,
J=8Hz), 8.00 (1H, d, J=8Hz), 8.17 (1H, m)

10

43) 4-[2-(3-Hydroxypropyl)-1H-benzimidazol-4-yl]-
carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylnpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide dihydrochloride

15

NMR (DMSO-d₆, δ) : 1.38-1.64 (4H, m), 1.68-1.84 (2H,
m), 1.94-2.10 (2H, m), 2.22 (3H, s), 2.39 (2H, t,
J=7Hz), 2.70-2.77 (3H, m), 2.78-3.16 (6H, m), 3.20
(3H, s), 3.26-3.57 (5H, m), 3.70 (3H, s), 3.79-4.14
(2H, m), 4.43 (1H, m), 6.66 (1H, d, J=8Hz), 6.83
20 (1H, s), 6.88-7.00 (2H, m), 7.05 (1H, d, J=8Hz),
7.46 (1H, m), 7.81 (1H, m), 7.93-8.10 (2H, m)

20

44) 4-[N-(1H-Benzimidazol-4-yl)carbonyl]-N-[2-[4,4-
dimethyl(2,5-oxazolinyl)]phenyl]-3-methoxy-N-
25 methylbenzamide dihydrochloride

25

NMR (DMSO-d₆, δ) : 1.30-1.55 (6H, m), 3.30 (3H, s),
3.67 (3H, s), 3.96-4.11 (1H, m), 4.22-4.38 (1H, m),
7.00-7.10 (1H, m), 7.17-8.04 (9H, m), 9.60 (1H, s)

30

45) 4-[N-[2-[(Dimethylamino)methyl]-1H-benzimidazol-4-yl]-
carbonyl]-N-[2-[4,4-dimethyl(2,5-oxazolinyl)]phenyl]-3-
methoxy-N-methylbenzamide trihydrochloride

35

NMR (DMSO-d₆, δ) : 1.30-1.56 (6H, m), 2.90 and 2.94
(Total 6H, s), 3.31 and 3.46 (Total 3H, s), 3.84
and 4.21 (Total 3H, s), 4.42-4.81 (4H, m), 7.06-

8.32 (10H, m)

- 46) N-[2-[4,4-Dimethyl(2,5-oxazolinyl)]phenyl]-3-methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-
5 carbonylaminobenzamide dihydrochloride
NMR (DMSO-d₆, δ) : 1.32-1.51 (6H, m), 2.78 (3H, s),
3.29 and 3.30 (Total 3H, s), 3.60 (3H, s), 3.92-
4.16 (1H, m), 4.30-4.49 (1H, m), 6.90-8.36 (10H, m)
- 10 47) 4-[N-(1H-Benzimidazol-4-yl)carbamoyl]-3-methoxy-N-methyl-N-[2-(1-pyrrolyl)phenyl]benzamide hydrochloride
NMR (DMSO-d₆, δ) : 3.40 (3H, s), 3.68 (3H, s), 6.23
(2H, s), 6.52 (2H, s), 6.60 (2H, s), 7.24 (1H, d,
J=8Hz), 7.30-7.58 (6H, m), 7.62 (1H, d, J=8Hz),
15 7.71 (1H, d, J=8Hz), 7.84 (1H, d, J=8Hz), 9.46 (1H,
s)
- 48) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)-
carbamoyl]-N-(2-piperidinophenyl)benzamide hydrochloride
20 NMR (DMSO-d₆, δ) : 1.38-1.70 (6H, m), 2.16-2.40 (2H,
m), 2.65-2.89 (5H, m), 3.40 (3H, s), 3.63 (3H, br
s), 6.83-7.03 (2H, m), 7.04-7.28 (3H, m), 7.40-7.72
(5H, m)
- 25 49) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-
carbonylamino-N-(2-piperidinophenyl)benzamide
dihydrochloride
NMR (DMSO-d₆, δ) : 1.37-1.72 (6H, m), 2.14-2.43 (2H,
m), 2.65-2.92 (5H, m), 3.39 (2H, s), 3.59 (3H, br
30 s), 6.80-7.32 (5H, m), 7.47 (1H, m), 7.58 (1H, dd,
J=8, 8Hz), 7.70 (1H, m), 7.94 (1H, d, J=8Hz), 8.14
(1H, d, J=8Hz)
- 50) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-
35 yl)carbamoyl]-N-[2-(4-methyl-1-piperazinyl)phenyl]-

benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 2.78 (3H, s), 2.80-3.26 (9H, m),
3.30-3.82 (8H, m), 6.90-7.10 (2H, m), 7.14-7.32
(3H, m), 7.40-7.63 (4H, m), 7.75 (1H, d, J=8Hz)

5

51) 3-Methoxy-N-methyl-4-(2-methyl-1H-benzimidazol-4-yl)-
carbonylamino-N-[2-(4-methyl-1-piperazinyl)phenyl]-
benzamide dihydrochloride

10 NMR (DMSO-d₆, δ) : 2.30 (1H, br s), 2.74 (3H, s), 2.81
and 2.82 (Total 3H, s), 2.86-3.26 (6H, m), 3.29-
3.49 (5H, m), 3.60 (3H, s), 6.90-7.02 (2H, m),
7.11-7.29 (3H, m), 7.42-7.58 (2H, m), 7.78-8.16
(3H, m)

15 52) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)-
carbamoyl]-N-[2-(2,5-oxazolyl)phenyl]benzamide
hydrochloride

20 NMR (DMSO-d₆, δ) : 2.77 (3H, s), 3.36 (3H, s), 3.61
(3H, s), 6.70 (1H, d, J=8Hz), 6.76 (1H, s), 7.37-
7.69 (8H, m), 7.84 (1H, d, J=8Hz), 8.32 (1H, s)

53) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-yl)-
carbamoyl]-N-[2-(2,5-oxazolinyl)phenyl]benzamide
dihydrochloride

25 NMR (DMSO-d₆, δ) : 2.78 (3H, s), 3.28 (3H, s), 3.38-
4.11 (7H, m), 7.08-8.80 (10H, m)

54) 3-Methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-
yl)carbamoyl]-N-[2-(3H,4H,5H-2,6-oxazinyl)phenyl]-
benzamide dihydrochloride

30 NMR (DMSO-d₆, δ) : 1.90-2.06 (2H, m), 2.72-2.85 (3H,
m), 3.12-4.08 (10H, m), 6.82-8.58 (10H, m)

55) N-[2-(1-Aza-3-oxaspiro[4.4]non-1-en-2-yl)phenyl]-3-
35 methoxy-N-methyl-4-[N-(2-methyl-1H-benzimidazol-4-

yl}carbamoyl}benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.52-2.07 (8H, m), 2.79 (3H, s),
3.30 (3H, s), 3.66 (3H, s), 4.38-4.58 (2H, m),
6.97-7.10 (2H, m), 7.20-8.08 (8H, m)

5

56) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-sulfamoylamino-1H-benzimidazol-4-yl)carbonylaminobenzamide dihydrochloride

10 NMR (DMSO-d₆, δ) : 1.38-1.52 (2H, m), 1.52-1.65 (2H, m), 1.65-1.83 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.75 (3H, d, J=4Hz), 2.80-3.08 (3H, m), 3.20 (3H, s), 3.33-3.42 (3H, m), 3.80-3.91 (1H, m), 3.91-4.02 (1H, m), 4.02-4.13 (1H, m), 4.39-4.50
15 (1H, m), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.89-6.98 (2H, m), 7.04 (1H, d, J=8Hz), 7.21 (1H, t, J=8Hz), 7.38-7.48 (1H, m), 7.54-7.67 (1H, m), 7.77 (1H, d, J=8Hz), 9.61 (1H, br peak), 10.55 (1H, br peak)

20

57) 4-[2-Carbamoyl-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

25 NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.65 (2H, m), 1.65-1.81 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.77 (3H, d, J=5Hz), 2.83-3.10 (3H, m), 3.20 (3H, s), 3.31-3.48 (3H, m), 3.77 (3H, s), 3.81-3.92 (1H, m), 3.92-4.01 (1H, m), 4.10 (1H, br d, J=15Hz), 4.45 (1H, br d, J=15Hz), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.91-7.03 (2H, m), 7.07 (1H, d, J=8Hz), 7.51 (1H, t, J=8Hz), 7.71 (1H, s), 7.80 (1H, d, J=8Hz), 8.04 (1H, d, J=8Hz), 8.30 (1H, s), 8.37 (1H, d, J=8Hz), 10.46 (1H, br peak), 12.00 (1H, s)

35

58) 4-[2-(N,N-Dimethylcarbamoyl)-1H-benzimidazol-4-yl]-
carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide dihydrochloride

5 NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.66 (2H,
m), 1.70-1.82 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=8Hz), 2.77 (3H, s), 2.83-3.10 (3H, m), 3.15 (3H,
s), 3.20 (3H, s), 3.33-3.60 (6H, m), 3.72 (3H, s),
3.80-3.93 (1H, m), 3.93-4.02 (1H, m), 4.02-4.15
10 (1H, m), 4.40-4.50 (1H, m), 6.65 (1H, d, J=8Hz),
6.82 (1H, s), 6.88-6.98 (2H, m), 7.05 (1H, d,
J=8Hz), 7.49 (1H, t, J=8Hz), 7.80 (1H, d, J=8Hz),
8.03 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz)

15 59) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-piperidyl)-1H-
benzimidazol-4-yl]carbonylaminobenzamide
trihydrochloride

20 NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.65 (2H,
m), 1.70-1.84 (2H, m), 2.03-2.20 (2H, m), 2.23 (3H,
s), 2.32-2.48 (4H, m), 2.75 (3H, d, J=5Hz), 2.83-
3.15 (4H, m), 3.19 (3H, s), 3.30-3.54 (7H, m), 3.77
(3H, s), 3.88 (1H, br peak), 3.97 (1H, br peak),
4.09 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz),
25 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.90-7.00 (2H,
m), 7.05 (1H, d, J=8Hz), 7.36 (1H, t, J=8Hz), 7.74
(1H, d, J=8Hz), 7.93 (1H, d, J=8Hz), 8.37 (1H, br
d, J=8Hz), 8.97 (1H, br peak), 9.27 (1H, br peak),
11.05 (1H, br peak)

30

60) 4-(2-Aminomethyl-1-methyl-1H-benzimidazol-4-yl)-
carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide trihydrochloride

35 NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.67 (2H,

- m), 1.70-1.84 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.80-3.10 (3H, m), 3.20 (3H, s), 3.32-3.54 (3H, m), 3.68-4.03 (8H, m), 4.10 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz), 4.51-4.61 (2H, m), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.91-7.00 (2H, m), 7.06 (1H, d, J=8Hz), 7.50 (1H, t, J=8Hz), 7.93 (1H, d, J=8Hz), 8.03 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz), 8.83-8.96 (3H, m)
- 10 61) 4-(2-Aminomethyl-3-methyl-3H-benzimidazol-4-yl)-carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide trihydrochloride
- 15 NMR (DMSO-d₆-D₂O, δ) : 1.41-1.53 (2H, m), 1.53-1.68 (2H, m), 1.68-1.85 (2H, m), 2.26 (3H, s), 2.41 (2H, t, J=7.5Hz), 2.80 (3H, s), 2.86-3.14 (3H, m), 3.21 (3H, s), 3.35-3.52 (2H, m), 3.68 (1H, s), 3.75 (3H, s), 3.84-4.05 (2H, m), 4.05-4.19 (1H, m), 4.42-4.53 (3H, m), 6.70 (1H, d, J=8Hz), 6.87 (1H, s), 6.91-7.01 (2H, m), 7.10 (1H, d, J=8Hz), 7.33 (1H, t, J=8Hz), 7.49 (1H, d, J=8Hz), 7.78 (1H, d, J=8Hz), 7.84 (1H, d, J=8Hz)
- 20
- 25 62) 4-(2-Methylthio-1H-benzimidazol-4-yl)carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- 30 NMR (CDCl₃, δ) : 1.40-1.52 (2H, m), 1.52-1.66 (2H, m), 1.66-1.82 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7Hz), 2.75 (3H, d, J=5Hz), 2.80-3.08 (5H, m), 3.20 (3H, s), 3.33-3.49 (3H, m), 3.72 (3H, s), 3.87 (1H, br peak), 3.95 (1H, br peak), 4.10 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.89-6.99 (2H, m), 7.04 (1H, d, J=8Hz), 7.30 (1H, t, J=8Hz), 7.63 (1H, d, J=8Hz), 7.88 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)
- 35

63) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-methylsulfonyl-1H-benzimidazol-4-yl)carbonylaminobenzamide dihydrochloride

5 NMR (DMSO-d₆, δ) : 1.36-1.51 (2H, m), 1.51-1.66 (2H, m), 1.66-1.83 (2H, m), 2.21 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.76 (3H, d, J=5Hz), 2.81-3.10 (3H, m), 3.19 (3H, s), 3.30-3.68 (6H, m), 3.80 (3H, s), 3.88 (1H, br peak), 3.96 (1H, br peak), 4.09 (1H, br d, J=15Hz),
10 4.45 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.90-7.00 (2H, m), 7.04 (1H, d, J=8Hz), 7.64 (1H, t, J=8Hz), 7.90 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

15 64) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-sulfamoyl-1H-benzimidazol-4-yl)carbonylaminobenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.67 (2H, m), 1.70-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.76 (3H, d, J=5Hz), 2.85-3.09 (3H, m),
20 3.20 (3H, s), 3.33-3.45 (3H, m), 3.75 (3H, s), 3.87 (1H, br peak), 3.97 (1H, br peak), 4.10 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.64 (1H, d, J=8Hz), 6.83 (1H, s), 6.89 (1H, s), 6.96 (1H, d, J=8Hz),
25 7.04 (1H, d, J=8Hz), 7.56 (1H, t, J=8Hz), 7.83 (1H, d, J=8Hz), 8.07 (1H, d, J=8Hz), 8.20 (2H, s), 8.36 (1H, d, J=8Hz)

65) 4-(2-Aminomethyl-1H-benzimidazol-4-yl)carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride

30 NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.68 (2H, m), 1.68-1.84 (2H, m), 2.22 (3H, s), 2.41 (2H, t, J=6Hz), 2.75 (3H, d, J=5Hz), 2.81-3.10 (3H, m),
35

- 3.22 (3H, s), 3.32-3.52 (4H, m), 3.82-4.02 (4H, m),
4.09 (1H, br d, J=12Hz), 4.30-4.50 (3H, m), 6.65
(1H, d, J=8Hz), 6.81 (1H, s), 7.02 (1H, d, J=8Hz),
7.07-7.16 (2H, m), 7.21 (1H, t, J=8Hz), 7.31 (1H,
5 d, J=8Hz), 7.88 (1H, d, J=8Hz), 8.05 (1H, br peak),
8.64-8.75 (3H, m)
- 66) 4-[2-(2-Aminoethyl)-1H-benzimidazol-4-yl]carbamoyl-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
10 yl)carbonylpent-1-yloxy]phenyl]benzamide
trihydrochloride
NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.67 (2H,
m), 1.71-1.84 (2H, m), 2.22 (3H, s), 2.41 (2H, t,
J=7.5Hz), 2.80-3.09 (3H, m), 3.21 (3H, d, J=5Hz),
15 3.27-3.74 (7H, m), 3.74-4.03 (5H, m), 4.09 (1H, br
d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.65 (1H, d,
J=8Hz), 6.82 (1H, s), 6.99 (1H, d, J=8Hz), 7.06
(1H, s), 7.11 (1H, d, J=8Hz), 7.26-7.49 (2H, m),
7.65-7.80 (1H, m), 7.90-7.93 (1H, m), 8.26 (3H, br
20 peak)
- 67) 4-(2-Hydroxymethyl-1H-benzimidazol-4-yl)carbamoyl-3-
methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-
yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
25 NMR (DMSO-d₆, δ) : 1.41-1.52 (2H, m), 1.52-1.66 (2H,
m), 1.71-1.85 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.76 (3H, s), 2.82-3.09 (3H, m), 3.21
(3H, s), 3.79 (3H, s), 3.85-4.03 (2H, m), 4.03-4.15
(1H, m), 4.38-4.51 (1H, m), 4.91 (2H, s), 6.66 (1H,
30 d, J=8Hz), 6.84 (1H, s), 7.00 (1H, d, J=8Hz), 7.04
(1H, s), 7.11 (1H, d, J=8Hz), 7.36-7.52 (2H, m),
7.63-7.72 (1H, m), 7.72-7.84 (1H, m)
- 68) 4-[2-Amino-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-
35 methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-

carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.68 (2H, m), 1.70-1.85 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.73 (3H, d, J=5Hz), 2.80-3.09 (3H, m), 3.20 (3H, s), 3.30-3.60 (3H, m), 3.71 (3H, s), 3.84-4.02 (2H, m), 4.08 (1H, br d, J=15Hz), 4.43 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.97 (1H, d, J=8Hz), 7.01 (1H, s), 7.10 (1H, d, J=8Hz), 7.16-7.23 (2H, m), 7.30-7.38 (1H, m), 7.54 (1H, d, J=8Hz), 8.37 (2H, s)

69) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-[(methylsulfonyl)amino]-1H-benzimidazol-4-yl]carbamoylbenzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.52 (2H, m), 1.52-1.67 (2H, m), 1.67-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.76 (3H, d, J=5Hz), 2.82-3.08 (3H, m), 3.23 (3H, s), 3.32-3.48 (3H, m), 3.51 (3H, s), 3.75-4.23 (6H, m), 4.45 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.80 (1H, s), 6.97-7.15 (4H, m), 7.15-7.30 (2H, m), 7.88 (1H, d, J=8Hz), 8.10 (1H, d, J=8Hz)

70) 3-Methoxy-4-[2-methoxymethyl-1H-benzimidazol-4-yl]-carbamoyl-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.67 (2H, m), 1.67-1.85 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.80-3.10 (3H, m), 3.21 (3H, s), 3.26-3.54 (6H, m), 3.77 (3H, s), 3.81-4.20 (3H, m), 4.43 (1H, br d, J=15Hz), 4.90 (2H, s), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 7.00 (1H, d, J=8Hz), 7.03 (1H, s), 7.11 (1H, d, J=8Hz),

7.40-7.58 (2H, m), 7.66 (1H, d, J=8Hz), 7.79 (1H, d, J=8Hz)

- 71) 3-Methoxy-N-methyl-4-[2-methyl-1H-benzimidazol-4-yl]carbamoyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
5 NMR (DMSO-d₆, δ) : 1.41-1.53 (2H, m), 1.53-1.67 (2H, m), 1.72-1.85 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.73 (3H, t, J=5Hz), 2.78 (3H, s), 2.82-3.13 (3H, m), 3.21 (3H, s), 3.30-3.56 (3H, m), 3.75 (3H, s), 3.80-4.17 (3H, m), 4.43 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.84 (1H, s), 6.98 (1H, d, J=8Hz), 7.02 (1H, s), 7.10 (1H, d, J=8Hz), 7.40-7.65 (4H, m)
10
- 72) 4-[1,2-dimethyl-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
15 NMR (DMSO-d₆, δ) : 1.39-1.53 (2H, m), 1.53-1.65 (2H, m), 1.71-1.85 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.67-2.80 (6H, m), 2.80-3.10 (3H, m), 3.21 (3H, s), 3.25-3.64 (3H, m), 3.73-4.20 (9H, m), 4.38-4.50 (1H, m), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.95-7.08 (2H, m), 7.11 (1H, d, J=8Hz), 7.45 (1H, br peak), 7.54-7.82 (3H, s)
20
- 73) 4-[2-Ethyl-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
25 NMR (DMSO-d₆, δ) : 1.37-1.52 (5H, m), 1.52-1.68 (2H, m), 1.71-1.85 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.75 (3H, s), 2.80-3.17 (5H, m), 3.21 (3H, s), 3.31-3.61 (3H, m), 3.77 (3H, s), 3.85-4.01 (2H, m), 4.09 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.99
30
- 35

(1H, d, J=8Hz), 7.03 (1H, s), 7.11 (1H, d, J=8Hz),
7.43 (1H, br peak), 7.53 (1H, br peak), 7.66 (2H,
br peak)

5 74) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(n-propyl)-1H-
benzimidazol-4-yl]carbamoylbenzamide dihydrochloride
NMR (DMSO-d₆, δ) : 0.97 (3H, t, J=7.5Hz), 1.40-1.53
 (2H, m), 1.53-1.69 (2H, m), 1.69-1.84 (4H, m), 2.22
10 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.75 (3H, s), 2.80-
 3.14 (5H, m), 3.20 (3H, s), 3.30-3.60 (3H, m), 3.78
 (3H, s), 3.83-4.16 (3H, m), 4.04 (1H, br peak),
 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 7.00 (1H, d,
 J=8Hz), 7.03 (1H, s), 7.10 (1H, d, J=8Hz), 7.43
15 (1H, br peak), 7.52 (1H, br peak), 7.66 (2H, br
 peak)

 75) 4-[2-Isopropyl-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-
N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-
20 carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
NMR (DMSO-d₆, δ) : 1.40-1.53 (8H, m), 1.53-1.69 (2H,
 m), 1.69-1.88 (2H, m), 2.23 (3H, s), 2.40 (2H, t,
 J=7.5Hz), 2.73 (3H, d, J=5Hz), 2.80-3.08 (3H, m),
 3.21 (3H, s), 3.31-3.56 (4H, m), 3.76 (3H, s),
25 3.85-4.03 (2H, m), 4.09 (1H, br d, J=15Hz), 4.43
 (1H, d, J=15Hz), 6.66 (1H, d, J=8Hz), 6.83 (1H, s),
 6.98 (1H, d, J=8Hz), 7.02 (1H, s), 7.10 (1H, d,
 J=8Hz), 7.39-7.69 (3H, m), 7.80 (1H, br peak)

30 76) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-trifluoromethyl-
1H-benzimidazol-4-yl]carbamoylbenzamide dihydrochloride
NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.68 (2H,
 m), 1.68-1.83 (2H, m), 2.21 (3H, s), 2.40 (2H, t,
 J=7.5Hz), 2.76 (3H, d, J=5Hz), 2.83-3.13 (3H, m),
35

3.22 (3H, s), 3.33-3.63 (3H, m), 3.82-4.02 (4H, m),
4.09 (1H, br d, J=15Hz), 4.45 (1H, br d, J=15Hz),
6.64 (1H, d, J=8Hz), 6.81 (1H, s), 7.00-7.16 (3H,
m), 7.32-7.46 (2H, m), 7.90 (1H, br peak), 8.23
5 (1H, br peak)

77) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(3-pyridyl)-1H-
benzimidazol-4-yl]carbamoylbenzamide trihydrochloride
10 NMR (DMSO-d₆, δ) : 1.41-1.53 (2H, m), 1.53-1.66 (2H,
m), 1.73-1.85 (2H, m), 2.21 (3H, s), 2.41 (2H, t,
J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.81-3.08 (3H, m),
3.23 (3H, s), 3.31-3.51 (3H, m), 3.71-4.16 (6H, m),
4.40-4.52 (1H, m), 6.66 (1H, d, J=8Hz), 6.81 (1H,
15 s), 7.04 (1H, d, J=8Hz), 7.08-7.16 (2H, m), 7.29
(1H, t, J=8Hz), 7.38 (1H, d, J=8Hz), 7.85 (1H, dd,
J=5, 8Hz), 7.90 (1H, d, J=8Hz), 8.19 (1H, d,
J=8Hz), 8.77 (1H, d, J=8Hz), 8.83 (1H, d, J=5Hz),
9.48 (1H, s)

20

78) 4-[2-(N,N-Dimethylcarbamoyl)-1H-benzimidazol-4-yl]-
carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methyl-
piperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
dihydrochloride
25 NMR (DMSO-d₆, δ) : 1.38-1.52 (2H, m), 1.52-1.65 (2H,
m), 1.65-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.79-3.07 (3H, m),
3.11 (3H, s), 3.21 (3H, s), 3.31-3.49 (3H, m), 3.71
(3H, s), 3.82-4.01 (4H, m), 4.01-4.38 (2H, m), 4.45
30 (1H, d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.80 (1H, s),
7.02 (1H, d, J=8Hz), 7.06-7.17 (2H, m), 7.24-7.31
(2H, m), 7.84-7.93 (1H, m), 8.19-8.29 (1H, m)

79) 4-[2-(N,N-Dimethylaminomethyl)-1H-benzimidazol-4-yl]-
carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-

35

methylypiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.53 (2H, m), 1.53-1.67 (2H,
m), 1.67-1.84 (2H, m), 2.20 (3H, s), 2.40 (2H, t,
5 J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.80-3.10 (9H, m),
3.21 (3H, s), 3.31-3.45 (3H, m), 3.81-4.01 (4H, m),
4.08 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz),
4.61 (2H, s), 6.65 (1H, d, J=8Hz), 6.81 (1H, s),
7.02 (1H, d, J=8Hz), 7.07-7.15 (2H, m), 7.24 (1H,
10 t, J=8Hz), 7.34 (1H, d, J=8Hz), 7.88 (1H, d,
J=8Hz), 8.10 (1H, br peak)

80) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(1-imidazolyl)-
15 methyl-1H-benzimidazol-4-yl]carbamoylbenzamide
trihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.68 (2H,
m), 1.68-1.85 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.74 (3H, s), 2.80-3.10 (3H, m), 3.23
20 (3H, s), 3.31-3.54 (3H, m), 3.70-4.20 (6H, m), 4.44
(1H, br peak), 5.84 (2H, s), 6.65 (1H, d, J=8Hz),
6.82 (1H, s), 7.01 (1H, d, J=8Hz), 7.04-7.15 (2H,
m), 7.21 (1H, t, J=8Hz), 7.30 (1H, d, J=8Hz), 7.78
(1H, s-like), 7.87 (1H, d, J=8Hz), 7.92 (1H,
25 s-like), 8.08 (1H, br peak), 9.40 (1H, s)

81) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-[(4-methyl-
piperazin-1-yl)methyl]-1H-benzimidazol-4-yl]-
30 carbamoylbenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.64 (2H,
m), 1.70-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.70-3.18 (9H, m), 3.21 (3H, s), 3.32-
4.20 (19H, m), 4.43 (1H, br d, J=15Hz), 6.66 (1H,
35 d, J=8Hz), 6.84 (1H, s), 7.00 (1H, d, J=8Hz), 7.05

(1H, s), 7.10 (1H, d, J=8Hz), 7.37 (1H, br peak),
7.45 (1H, br peak), 7.70 (1H, br peak), 7.89 (1H,
br peak)

- 5 82) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(morpholin-4-
ylmethyl)-1H-benzimidazol-4-yl]carbamoylbenzamide
trihydrochloride

10 NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.66 (2H,
m), 1.66-1.85 (2H, m), 2.21 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.78-3.11 (3H, m),
3.21 (3H, s), 3.30-4.00 (16H, m), 4.09 (1H, d,
J=15Hz), 4.44 (1H, d, J=15Hz), 4.64 (2H, s), 6.65
(1H, d, J=8Hz), 6.82 (1H, s), 7.03 (1H, d, J=8Hz),
15 7.06-7.15 (2H, m), 7.26 (1H, t, J=8Hz), 7.36 (1H,
d, J=8Hz), 7.87 (1H, d, J=8Hz), 8.08 (1H, br peak)

- 20 83) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(pyrrolidin-1-
ylmethyl)-1H-benzimidazol-4-yl]carbamoylbenzamide
trihydrochloride

25 NMR (DMSO-d₆, δ) : 1.39-1.53 (2H, m), 1.53-1.68 (2H,
m), 1.68-1.87 (2H, m), 1.87-2.11 (4H, m), 2.21 (3H,
s), 2.40 (2H, t, J=7.5Hz), 2.74 (3H, d, J=5Hz),
2.80-3.11 (3H, m), 3.21 (3H, s), 3.25-3.79 (7H, m),
3.79-4.02 (5H, m), 4.09 (1H, br d, J=15Hz), 4.44
(1H, br d, J=15Hz), 4.70 (2H, s), 6.65 (1H, d,
J=8Hz), 6.81 (1H, s), 7.01 (1H, d, J=8Hz), 7.04-
7.15 (2H, m), 7.24 (1H, t, J=8Hz), 7.32 (1H, d,
30 J=8Hz), 7.87 (1H, d, J=8Hz), 8.03-8.13 (1H, m)

- 35 84) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(piperidino-
methyl)-1H-benzimidazol-4-yl]carbamoylbenzamide
trihydrochloride

NMR (DMSO-d₆, δ) : 1.38-1.90 (12H, m), 2.21 (3H, s),
2.40 (2H, t, J=7.5Hz), 2.75 (3H, d, J=5Hz), 2.80-
3.17 (3H, m), 3.23 (3H, s), 3.28-4.02 (15H, m),
4.09 (1H, br d, J=15Hz), 4.44 (1H, br d, J=15Hz),
5 4.59 (2H, s), 6.64 (1H, d, J=8Hz), 6.81 (1H, s),
7.02 (1H, d, J=8Hz), 7.06-7.15 (2H, m), 7.24 (1H,
t, J=8Hz), 7.34 (1H, d, J=8Hz), 7.87 (1H, d,
J=8Hz), 8.09 (1H, br peak)

10 85) 4-[2-[2-(Dimethylamino)ethyl]-1H-benzimidazol-4-yl]-
carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.67 (2H,
15 m), 1.70-1.86 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.75 (3H, d, J=5Hz), 2.81-3.08 (9H, m),
3.22 (3H, s), 3.28-3.72 (7H, m), 3.82-4.02 (5H, m),
4.09 (1H, br d, J=15Hz), 4.43 (1H, br d, J=15Hz),
6.65 (1H, d, J=8Hz), 6.81 (1H, s), 7.01 (1H, d,
20 J=8Hz), 7.05-7.17 (2H, m), 7.20-7.40 (2H, m), 7.80
(1H, d, J=8Hz), 7.93 (1H, br peak)

86) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-[2-(4-
25 methylpiperazin-1-yl)ethyl]-1H-benzimidazol-4-yl]-
carbamoylbenzamide trihydrochloride

NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.66 (2H,
m), 1.71-1.86 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.70-2.79 (6H, m), 2.79-3.65 (21H, m),
30 3.78-4.01 (5H, m), 4.09 (1H, br d, J=15Hz), 4.44
(1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.82 (1H,
s), 7.00 (1H, d, J=8Hz), 7.05 (1H, s), 7.10 (1H, d,
J=8Hz), 7.35 (1H, t, J=8Hz), 7.46 (1H, d, J=8Hz),
7.66-7.80 (2H, m)

- 87) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-methylpiperazin-1-yl)-1H-benzimidazol-4-yl]carbamoylbenzamide trihydrochloride

5 NMR (DMSO-d₆, δ) : 1.48-1.53 (2H, m), 1.53-1.65 (2H, m), 1.70-1.85 (2H, m), 2.21 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.80 (3H, s), 2.83-3.10 (3H, m), 3.16-3.83 (13H, m), 3.83-4.15 (2H, m), 4.33 (1H, br d, J=15Hz), 4.43 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.81 (1H, s), 6.96 (1H, d, J=8Hz), 7.01 (1H, s), 7.09 (1H, d, J=8Hz), 7.13-7.24 (2H, m), 7.61 (1H, br peak), 7.79 (1H, br peak)

- 15 88) 4-[2-Dimethylamino-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride
- 20 NMR (DMSO-d₆, δ) : 1.41-1.52 (2H, m), 1.52-1.67 (2H, m), 1.71-1.83 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.75 (3H, s), 2.80-3.10 (3H, m), 3.15 (6H, s), 3.21 (3H, s), 3.70-4.50 (6H, m), 4.50 (1H, br peak), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.95-7.06 (2H, m), 7.11 (1H, d, J=8Hz), 7.50-8.10 (4H, m)

- 25 89) 3-Methoxy-N-methyl-4-[2-[[2-(methylamino)ethyl]amino]-1H-benzimidazol-4-yl]carbamoyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide trihydrochloride
- 30 NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.65 (2H, m), 1.65-1.84 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.61 (2H, t-like, J=5Hz), 2.74 (3H, d, J=5Hz), 2.78-3.11 (3H, m), 3.11-3.29 (5H, m), 3.29-3.55 (3H, m), 3.72 (3H, s), 3.73-4.01 (2H, m), 4.09 (1H, br d, J=15Hz), 4.43 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.83 (1H, s), 6.95 (1H, d, J=8Hz),
- 35

7.00 (1H, s), 7.10 (1H, d, J=8Hz), 7.18-7.27 (2H, m), 7.47-7.61 (2H, m), 9.10 (3H, br peak)

- 90) 4-[2-[(2-Aminoethyl)methylamino]-1H-benzimidazol-4-yl]-
5 carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide trihydrochloride
- NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.67 (2H, m), 1.70-1.83 (2H, m), 2.23 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.79-3.11 (3H, m), 3.11-3.23 (5H, m), 3.26 (3H, s), 3.33-3.56 (5H, m), 3.71 (3H, s), 3.84-4.02 (5H, m), 4.09 (1H, br d, J=15Hz), 4.43 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.93 (1H, d, J=8Hz), 6.99
10 (1H, s), 7.09 (1H, d, J=8Hz), 7.18-7.28 (2H, m), 7.50 (1H, br peak), 7.80 (1H, br peak), 8.29 (3H, br peak)

- 91) 4-[2-(1-Imidazolyl)-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride
- NMR (DMSO-d₆, δ) : 1.41-1.52 (2H, m), 1.52-1.67 (2H, m), 1.67-1.84 (2H, m), 2.20 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.74 (3H, s-like), 2.81-3.10 (3H, m), 3.22 (3H, s), 3.30-3.53 (3H, m), 3.78-4.22 (6H, m), 4.39-4.51 (1H, m), 6.66 (1H, d, J=8Hz), 6.81 (3H, s), 7.04 (1H, d, J=8Hz), 7.09-7.19 (2H, m), 7.23-7.40 (2H, m), 7.65 (1H, s), 7.91 (1H, d, J=8Hz),
25 8.15-8.29 (2H, m), 9.28 (1H, s)

- 92) 4-[2-[[2-(Dimethylamino)ethyl]amino]-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide trihydrochloride
- 35

NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.66 (2H, m), 1.70-1.85 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7.5Hz), 2.73 (3H, d, J=5Hz), 2.84 (6H, s), 2.88-3.10 (3H, m), 3.20 (3H, s), 3.28-3.53 (7H, m), 3.84-4.02 (5H, m), 4.08 (1H, br d, J=15Hz), 4.43 (1H, br d, J=15Hz), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 6.92-7.02 (2H, m), 7.10 (1H, d, J=8Hz), 7.19-7.27 (2H, m), 7.50 (1H, d, J=8Hz), 7.58 (1H, br peak), 9.14 (1H, br peak)

10

93) 4-[2-[[2-(Dimethylamino)ethyl]methylamino]-1H-benzimidazol-4-yl]carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride

15

NMR (DMSO-d₆, δ) : 1.41-1.53 (2H, m), 1.53-1.66 (2H, m), 1.70-1.85 (2H, m), 2.22 (3H, s), 2.40 (2H, d, J=7.5Hz), 2.74 (3H, d, J=5Hz), 2.81-3.09 (9H, m), 3.21 (3H, s), 3.26 (3H, s), 3.32-3.55 (7H, m), 3.71 (3H, br s), 3.81-4.14 (3H, m), 4.44 (1H, br d, J=15Hz), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.92 (1H, d, J=8Hz), 6.98 (1H, s), 7.09 (1H, d, J=8Hz), 7.15-7.25 (2H, m), 7.49 (1H, br peak), 7.85 (1H, br peak)

20

94) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(1,2,4-triazol-1-yl)-1H-benzimidazol-4-yl]carbamoylbenzamide trihydrochloride

25

NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.67 (2H, m), 1.72-1.85 (2H, m), 2.22 (3H, s), 2.41 (2H, t, J=7.5Hz), 2.76 (3H, d, J=5Hz), 2.80-3.10 (3H, m), 3.21 (3H, s), 3.28-3.64 (3H, m), 3.83-4.03 (5H, m), 4.10 (1H, br d, J=15Hz), 4.45 (1H, br d, J=15Hz), 6.66 (1H, d, J=8Hz), 6.83 (1H, s), 7.05 (1H, d, J=8Hz), 7.08-7.17 (2H, m), 7.21-7.32 (2H, m), 7.90.

30

35

(1H, br peak), 8.19 (1H, br peak), 8.50 (1H, s),
9.41 (1H, s)

- 5 95) 4-[2-[(2-Methoxyethyl)amino]-1H-benzimidazol-4-yl]-
carbamoyl-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide dihydrochloride

10 NMR (DMSO-d₆, δ) : 1.39-1.52 (2H, m), 1.52-1.67 (2H,
m), 1.70-1.85 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=7.5Hz), 2.73 (3H, s), 2.80-3.10 (3H, m), 3.21
(3H, s), 3.27-3.49 (6H, m), 3.53-3.64 (4H, m), 3.73
(3H, s), 3.84-4.16 (3H, m), 4.43 (1H, br peak),
6.65 (1H, d, J=8Hz), 6.84 (1H, s), 6.97 (1H, d,
J=8Hz), 7.00 (1H, s), 7.10 (1H, d, J=8Hz), 7.16-
15 7.26 (2H, m), 7.40 (1H, br peak), 7.55 (1H, br
peak), 8.80 (1H, br peak)

- 20 96) 4-(2-Dimethylaminomethyl-1H-benzimidazol-4-yl)-
carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-
methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]-
benzamide trihydrochloride

25 NMR (DMSO-d₆, δ) : 1.40-1.52 (2H, m), 1.52-1.63 (2H,
m), 1.72-1.82 (2H, m), 2.22 (3H, s), 2.40 (2H, t,
J=7Hz), 2.73 (3H, s), 2.90-3.05 (10H, m), 3.20 (3H,
s), 3.35-3.50 (3H, m), 3.78 (3H, s), 3.95-4.13 (2H,
m), 4.40-4.45 (1H, m), 4.68 (2H, s), 6.64 (1H, d,
J=8Hz), 6.82 (1H, s), 6.90-6.95 (2H, m), 7.04 (1H,
d, J=8Hz), 7.46 (1H, t, J=8Hz), 7.89 (1H, d,
J=8Hz), 8.01 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz)

30

- 97) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methyloperazin-
1-yl)carbonylpent-1-yloxy]phenyl]-4-[2-(4-methyl-
piperazin-1-yl)methyl-1H-benzimidazol-4-yl]-
carbonylamino benzamide trihydrochloride

35 NMR (DMSO-d₆, δ) : 1.42-1.52 (2H, m), 1.52-1.64 (2H,

5 m), 1.73-1.82 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7Hz), 2.72 (3H, s), 2.77 (3H, s), 2.85-3.10 (4H, m), 3.20 (3H, s), 3.20-3.40 (7H, m), 3.45-3.56 (4H, m), 3.78 (3H, s), 3.83-4.10 (2H, m), 4.37-4.43 (3H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.90-6.93 (2H, m), 7.03 (1H, d, J=8Hz), 7.47 (1H, t, J=8Hz), 7.87 (1H, d, J=8Hz), 8.03 (1H, d, J=8Hz), 8.20-8.23 (1H, m)

10 98) 4-[2-(4-Dimethylaminopiperidino)methyl-1H-benzimidazol-4-yl]carbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide trihydrochloride


15 NMR (DMSO-d₆, δ) : 1.40-1.53 (2H, m), 1.53-1.66 (2H, m), 1.70-1.82 (2H, m), 2.10-2.35 (7H, m), 2.42 (2H, t, J=7Hz), 2.63-2.73 (7H, m), 2.85-3.08 (4H, m), 3.18 (3H, s), 3.33-3.53 (3H, m), 3.73-4.30 (9H, m), 4.40-4.47 (1H, m), 4.69 (2H, s), 6.63 (1H, d, J=8Hz), 6.83 (1H, s), 6.92-6.97 (2H, m), 7.04 (1H, d, J=8Hz), 7.47 (1H, t, J=8Hz), 7.91 (1H, d, J=8Hz), 8.02 (1H, d, J=8Hz), 8.30-8.35 (1H, m)

25 99) 3-Methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-4-(2-morpholinomethyl-1H-benzimidazole-4-yl)carbonylamino benzamide trihydrochloride

30 NMR (DMSO-d₆, δ) : 1.41-1.52 (2H, m), 1.52-1.64 (2H, m), 1.72-1.80 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=8Hz), 2.73 (3H, s), 2.80-3.10 (4H, m), 3.18 (3H, s), 3.34-3.55 (7H, m), 3.80 (3H, s), 3.82-4.10 (6H, m), 4.37-4.45 (1H, m), 4.72 (2H, s), 6.64 (1H, d, J=8Hz), 6.82 (1H, s), 6.92-6.94 (2H, m), 7.03 (1H, d, J=8Hz), 7.46 (1H, t, J=8Hz), 7.90 (1H, d, J=8Hz), 8.03 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz)

CLAIMS

1. A compound of the formula :



wherein

R^1 is aryl, cyclo(lower)alkyl or a heterocyclic group,

each of which may be substituted with substituent(s)

selected from the group consisting of halogen;

hydroxy; nitro; protected amino; amino; acyl;

substituted acyl; acyl(lower)alkylsulfinyl;

acyl(lower)alkylsulfonyl; acyloxy;

lower alkylamino(lower)alkylcarbamoyloxy;

aryl; cyano; a heterocyclic group;

lower alkenyl optionally substituted with acyl,

substituted acyl, aryl or acyl-substituted aryl;

lower alkynyl optionally substituted with amino,

acylamino or substituted acylamino;

lower alkyl optionally substituted with halogen, amino,

lower alkylamino, acylamino, substituted acylamino,

hydroxy, acyloxy, acyl(lower)alkanoyloxy, acyl,

substituted acyl, acyl(lower)alkoxyimino, aryl or

acyl-substituted aryl;

lower alkylthio optionally substituted with acyl or

substituted acyl;

alkoxy optionally substituted with aryl, substituted

aryl, hydroxy, acyloxy, amino, lower alkylamino,

protected amino, a heterocyclic group, acyl-substituted

pyridyl, substituted acyl-substituted pyridyl, halogen,

acyl(lower)alkylamino, N-protected-acyl(lower)-

- alkylamino, N-acyl(lower)alkyl-N-lower alkylamino, acyl, substituted acyl, acylamino, substituted acylamino, lower alkylhydrazinocarbonylamino, hydroxyimino, acyl(lower)alkoxyimino, substituted
- 5 acyl(lower)alkoxyimino, acyl(lower)alkoxy, guanidino or N-protected guanidino; and lower alkenyloxy optionally substituted with acyl or substituted acyl;
- R^2 is hydrogen; lower alkyl optionally substituted with hydroxy, aryl or acyl; or cyclo(lower)alkyl;
- 10 R^3 is hydrogen; halogen; hydroxy; acyloxy; substituted acyloxy; lower alkyl optionally substituted with hydroxy or lower alkoxy; lower alkoxy optionally substituted with aryl, amino, protected amino, acyl, hydroxy, cyano or lower alkylthio; nitro; amino; acyl; substituted
- 15 acyl; or cyclo(lower)alkyloxy;
- A is a single bond, O or NH;
- E is lower alkylene, lower alkenylene, $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}- \end{array}$, $\begin{array}{c} \text{O} \\ \parallel \\ -\text{S}- \\ \parallel \\ \text{O} \end{array}$, or a group of the formula :
- 20 -G-J-
- in which G is lower alkylene or $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}- \end{array}$ and
- $\begin{array}{c} \text{R}^4 \\ | \\ \text{-N-} \end{array}$
- 25 J is O or $\begin{array}{c} \text{R}^4 \\ | \\ \text{-N-} \end{array}$ (wherein R^4 is hydrogen or N-protective group);
- X is -CH=CH- , -CH=N- or S; and
- Y is aryl which may be substituted with acyl, protected amino(lower)alkanoyl, protected amino and nitro, amino and nitro or diamino;
- 30 or a condensed heterocyclic group which may be substituted with substituent(s) selected from the group consisting of halogen, acyl, lower alkoxy, hydroxy, guanidino, mercapto, acylamino, amino, a heterocyclic group, cyanoamino, amino(lower)alkyl(lower)alkylamino,
- 35 lower alkylamino, lower alkylamino(lower)alkylamino,

substituted-heterocyclic group, lower alkylhydrazino,
aryloxy, lower alkylthio, aryl, protected amino,
N-protected lower alkylamino(lower)alkylamino,
N-protected amino(lower)alkyl(N'-lower alkyl)amino,
5 amino(lower)alkyl(N-lower alkyl)amino, lower
alkylamino(lower)alkyl(N-lower alkyl)amino, lower
alkoxy(lower)alkylamino and lower alkyl optionally
substituted with aryl, ar(lower)alkoxy, cyano,
hydroxyimino, mercapto, lower alkylamino, acyloxy,
10 halogen, lower alkoxy, protected hydroxy, hydroxy, lower
alkoxyaryl, protected amino, amino, a heterocyclic group
or substituted heterocyclic group;
provided that when Y is phenyl which may be substituted with
lower alkyl or acyl,
15 then A is a single bond and

$$\begin{array}{c} \text{OR}^4 \\ ||| \\ \text{E is } -\text{CN}- \end{array}$$
 (wherein R⁴ is as defined
above);

and pharmaceutically acceptable salt thereof.

20

2. A compound according to claim 1, wherein
R¹ is aryl, cyclo(lower)alkyl or a heterocyclic group,
each of which may be substituted with substituent(s)
selected from the group consisting of halogen;
hydroxy; nitro; amino; acyl; substituted acyl;
25 acyl(lower)alkylsulfinyl; acyl(lower)alkylsulfonyl;
acyloxy; lower alkylamino(lower)alkylcarbamoyloxy;
aryl; cyano; a heterocyclic group;
lower alkenyl optionally substituted with acyl,
substituted acyl, aryl or acyl-substituted aryl;
30 lower alkynyl optionally substituted with amino,
acylamino or substituted acylamino;
lower alkyl optionally substituted with halogen, amino,
lower alkylamino, acylamino, substituted acylamino,
35 hydroxy, acyloxy, acyl(lower)alkanoyloxy, acyl,

substituted acyl, acyl(lower)alkoxyimino, aryl or
acyl-substituted aryl;

lower alkylthio optionally substituted with acyl or
substituted acyl;

5 alkoxy optionally substituted with aryl, substituted
aryl, hydroxy, acyloxy, amino, lower alkylamino,
protected amino, a heterocyclic group, acyl-substituted
pyridyl, substituted acyl-substituted pyridyl, halogen,
acyl(lower)alkylamino, N-protected-acyl(lower)-
10 alkylamino, N-acyl(lower)alkyl-N-lower alkylamino, acyl,
substituted acyl, acylamino, substituted acylamino,
lower alkylhydrazinocarbonylamino, hydroxyimino,
acyl(lower)alkoxyimino, substituted
acyl(lower)alkoxyimino, acyl(lower)alkoxy, guanidino or
15 N-protected guanidino; and lower alkenyloxy optionally
substituted with acyl or substituted acyl;

R^2 is hydrogen; lower alkyl optionally substituted with
hydroxy, aryl or acyl; or cyclo(lower)alkyl;

R^3 is hydrogen; halogen; hydroxy; acyloxy; substituted
20 acyloxy; lower alkyl optionally substituted with hydroxy
or lower alkoxy; lower alkoxy optionally substituted
with aryl, amino, protected amino, acyl, hydroxy, cyano
or lower alkylthio; nitro; amino; acyl; substituted
acyl; or cyclo(lower)alkyloxy;

25 A is a single bond, O or NH;

E is lower alkylene, lower alkenylene, $\begin{smallmatrix} \text{O} \\ \parallel \\ -\text{C}- \end{smallmatrix}$, $\begin{smallmatrix} \text{O} \\ \parallel \\ -\text{S}- \\ \parallel \\ \text{O} \end{smallmatrix}$, or
a group of the formula :

30 $-G-J-$

in which G is lower alkylene or $\begin{smallmatrix} \text{O} \\ \parallel \\ -\text{C}- \end{smallmatrix}$ and

J is O or $\begin{smallmatrix} \text{R}^4 \\ | \\ -\text{N}- \end{smallmatrix}$

(wherein R^4 is hydrogen or N-protective group);

X is $-\text{CH}=\text{CH}-$, $-\text{CH}=\text{N}-$ or S; and

Y is aryl which is substituted with protected amino and
nitro, amino and nitro or diamino; or a condensed
heterocyclic group which may be substituted with
substituent(s) selected from the group consisting of
halogen, acyl, lower alkoxy, hydroxy, guanidino,
mercapto, acylamino, amino and lower alkyl optionally
substituted with lower alkylamino, acyloxy, halogen,
lower alkoxy, protected hydroxy, hydroxy, lower
alkoxyaryl, protected amino, amino or a heterocyclic
group.

3. A compound according to claim 2, wherein

R^1 is aryl which may be substituted with lower alkoxy
substituted with acyl or acylamino,

R^2 is lower alkyl,

R^3 is hydrogen, lower alkyl or lower alkoxy,

A is a single bond or NH,

E is $-\overset{\text{O}}{\underset{\parallel}{\text{C}}}-$ or $-\overset{\text{O}}{\underset{\parallel}{\text{CNH}}}-$,

X is $-\text{CH}=\text{CH}-$, and

Y is a condensed heterocyclic group which is substituted with
lower alkyl optionally substituted with lower
alkylamino, acyloxy, halogen, lower alkoxy, protected
hydroxy, hydroxy, lower alkoxyaryl, protected amino,
amino or a heterocyclic group.

4. A compound according to claim 3, wherein

R^1 is phenyl or tolyl, each of which is substituted with
lower alkoxy substituted with N-lower
alkylpiperazinylcarbonyl,

R^3 is lower alkoxy,

A is a single bond,

E is $\text{-}\overset{\text{O}}{\parallel}\text{CNH-}$, and

Y is benzimidazol which is substituted with lower alkyl
optionally substituted with amino, hydroxy or N-lower
alkylpiperazinyl.

5

5. A compound according to claim 3, wherein

R^1 is phenyl or tolyl, each or which is substituted with
lower alkoxy substituted with N-lower
alkylpiperazinylcarbonyl,

10 R^3 is lower alkoxy,


A is NH,

E is $\text{-}\overset{\text{O}}{\parallel}\text{C-}$,

15 Y is benzimidazolyl which is substituted with lower alkyl
optionally substituted with amino, hydroxy or N-lower
alkylpiperazinyl.

6. A process for preparing the formula :

20



25

wherein

R^1 is aryl, cyclo(lower)alkyl or a heterocyclic group,
each of which may be substituted with substituent(s)
selected from the group consisting of halogen;
hydroxy; nitro; protected amino; amino; acyl;
30 substituted acyl; acyl(lower)alkylsulfinyl;
acyl(lower)alkylsulfonyl; acyloxy;
lower alkylamino(lower)alkylcarbamoxyloxy;
aryl; cyano; a heterocyclic group;
35 lower alkenyl optionally substituted with acyl,

- substituted acyl, aryl or acyl-substituted aryl;
 lower alkynyl optionally substituted with amino,
 acylamino or substituted acylamino;
 lower alkyl optionally substituted with halogen, amino,
 5 lower alkylamino, acylamino, substituted acylamino,
 hydroxy, acyloxy, acyl(lower)alkanoyloxy, acyl,
 substituted acyl, acyl(lower)alkoxyimino, aryl or
 acyl-substituted aryl;
 lower alkylthio optionally substituted with acyl or
 10 substituted acyl;
 alkoxy optionally substituted with aryl, substituted
 aryl, hydroxy, acyloxy, amino, lower alkylamino,
 protected amino, a heterocyclic group, acyl-substituted
 pyridyl, substituted acyl-substituted pyridyl, halogen,
 15 acyl(lower)alkylamino, N-protected-acyl(lower)-
 alkylamino, N-acyl(lower)alkyl-N-lower alkylamino, acyl,
 substituted acyl, acylamino, substituted acylamino,
 lower alkylhydrazinocarbonylamino, hydroxyimino,
 acyl(lower)alkoxyimino, substituted
 20 acyl(lower)alkoxyimino, acyl(lower)alkoxy, guanidino or
 N-protected guanidino; and lower alkenyloxy optionally
 substituted with acyl or substituted acyl;
 R^2 is hydrogen; lower alkyl optionally substituted with
 hydroxy, aryl or acyl; or cyclo(lower)alkyl;
 25 R^3 is hydrogen; halogen; hydroxy; acyloxy; substituted
 acyloxy; lower alkyl optionally substituted with hydroxy
 or lower alkoxy; lower alkoxy optionally substituted
 with aryl, amino, protected amino, acyl, hydroxy, cyano
 or lower alkylthio; nitro; amino; acyl; substituted
 30 acyl; or cyclo(lower)alkyloxy;
 A is a single bond, O or NH;
 E is lower alkylene, lower alkenylene, $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}- \end{array}$, $\begin{array}{c} \text{O} \\ \parallel \\ -\text{S}- \\ \parallel \\ \text{O} \end{array}$, or
 a group of the formula :

in which G is lower alkylene or $\text{-}\overset{\text{O}}{\parallel}\text{C-}$ and

J is O or $\text{-}\overset{\text{R}^4}{\underset{|}{\text{N}}}\text{-}$

5 (wherein R^4 is hydrogen or N-protective group);

X is -CH=CH- , -CH=N- or S; and

Y is aryl which may be substituted with acyl, protected amino(lower)alkanoyl, protected amino and nitro, amino and nitro or diamino;

10 or a condensed heterocyclic group which may be substituted with substituent(s) selected from the group consisting of halogen, acyl, lower alkoxy, hydroxy, guanidino, mercapto, acylamino, amino, a heterocyclic group, cyanoamino, amino(lower)alkyl(lower)alkylamino, 15 lower alkylamino, lower alkylamino(lower)alkylamino, substituted-heterocyclic group, lower alkylhydrazino, aryloxy, lower alkylthio, aryl, protected amino, N-protected lower alkylamino(lower)alkylamino, N-protected amino(lower)alkyl(N'-lower alkyl)amino, 20 amino(lower)alkyl(N-lower alkyl)amino, lower alkylamino(lower)alkyl(N-lower alkyl)amino, lower alkoxy(lower)alkylamino and lower alkyl optionally substituted with aryl, ar(lower)alkoxy, cyano, hydroxyimino, mercapto, lower alkylamino, acyloxy, 25 halogen, lower alkoxy, protected hydroxy, hydroxy, lower alkoxyaryl, protected amino, amino, a heterocyclic group or substituted heterocyclic group;

provided that when Y is phenyl which may be substituted with lower alkyl or acyl,


30 then A is a single bond and

E is $\text{-}\overset{\text{OR}^4}{\parallel}\text{CN-}$ (wherein R^4 is as defined above);

or pharmaceutically acceptable salt thereof,

35 which comprises,

1) reacting a compound of the formula :




or its salt with a compound of the formula :

10




or its reactive derivative at the carboxy group or the
sulfo group, or a salt thereof to provide a compound of
the formula :

15



25

or its salt, in the above formulas,
R¹, R², R³, X and Y are each as defined above, and




30

2) reacting a compound of the formula :

35

5



or its salt with a compound of the formula :


10



in the presence of a base to provide a compound of the formula :

15

20



25


or its salt, in the above formulas,
 R^1 , R^2 , R^3 , A, E and X are each as defined above,
 Ya is indolyl,
 R^5 is lower alkyl,
 Z^1 is an acid residue, and
 Yb is N-(lower alkyl)indolyl, or

30

3) subjecting a compound of the formula :

35


5



10

or its salt to reduction to provide a compound of the formula :

15




20

or its salt, in the above formulas,
 R^1 , R^2 , R^3 , A, E and X are each as defined above,
 Yc is phenyl substituted with amino and nitro, and
 Yd is phenyl substituted with diamino,

25


4) reacting a compound of the formula :

30



35

or its salt with aroyl halide,
 cyano(lower)alkylcarboxylic acid,
 mercapto(lower)alkylcarboxylic acid, lower alkylactone,
 1,1-dihalo-1,1-diphenoxymethane, diphenyl
 5 N-sulfamoylcarbonimide, diphenyl N-cyanocarbonimide,
 dicyandiamide, 1,1'-thiocarbonylimidazole, cyanogen
 bromide, lower alkoxycarbonyl isothiocyanate,
 tri(lower)alkyl orthoformate, tetra(lower)alkyl
 orthoformate, lower alkylcarboxylic acid,
 10 halo(lower)alkylcarboxylic acid, protected
 amino(lower)alkylcarbonyl halide or
 a heterocyclic(lower)alkylcarbonyl halide to provide a
 compound of the formula :




or its salt, in the above formulas,
 R¹, R², R³, A, E, X and Yd are each as defined above,
 and

Ye is benzimidazolyl optionally 2-position substituted
 25 with aryl, phenoxy, sulfamoylamino, cyanoamino,
 guanidino, mercapto, amino, lower
 alkoxycarbonylamino, lower alkoxy or lower alkyl
 optionally substituted with cyano, mercapto,
 hydroxy, halogen, protected amino or a heterocyclic
 30 group, or

5) reacting a compound of the formula :


5



10

or its salt with glyoxal and sodium hydrogen sulfite, or
sodium nitrite to provide a compound of the formula :

15




20

or its salt, in the above formulas,
R¹, R², R³, A, E, X and Yd are each as defined above,
and
Yf is quinoxalinylyl or benzotriazolyl, or

25

6) reacting a compound of the formula :


30



or its salt with an acylating agent to provide
a compound of the formula :

35

5



10

or its salt, in the above formulas,

R^1 , R^2 , R^3 , A, E, X and Ya are each as defined above,


and

Yg is N-acylindolyl, or

15

7) subjecting a compound of the formula :


20



25

or its salt to elimination reaction of the N-substituent group to provide a compound of the formula :

30



35

or its salt, in the above formulas,


R^1 , R^2 , R^3 , A, E and X are each as defined above,

Yh is (N-acyl)acylindolinyl, N-acylindolinyl,

(N-acyl)hydroxy(lower)alkylindolinyl, lower
 alkylamino(lower)alkylamino(N-acyl)indolinyl,
 (N-lower alkoxyarylmethyl)acylbenzimidazolyl,
 (N-lower alkoxycarbonyl)phthalimido(lower)-
 5 alkylindolyl, N-protected lower alkylamino(lower)-
 alkylamino(N-acyl)benzimidazolyl,
 (N-acyl)benzimidazolyl, (N-acyl)(lower)-
 alkylbenzimidazolyl, N-protected
 amino(lower)alkyl(N-lower alkyl)amino(N-
 10 acyl)benzimidazolyl, N-acylindolyl,
 (N-acyloxymethyl)indolyl,
 (N-acyl)acylindolyl, (N-arylmethyl)lower
 alkoxy(lower)alkylbenzimidazolyl or (N-lower
 alkoxyarylmethyl)acylbenzimidazolyl; and
 15 Yi is acylindolinyl, indolinyl,
 hydroxy(lower)alkylindolinyl,
 lower alkylamino(lower)alkylaminoidolinyl,
 acylbenzimidazolyl, phthalimido(lower)alkylindolyl,
 amino(lower)alkylindolyl,
 20 lower alkylamino(lower)alkylaminobenzimidazolyl,
 benzimidazolyl, lower alkylbenzimidazolyl,
 amino(lower)alkyl(N-lower
 alkyl)aminobenzimidazolyl, indolyl, acylindolyl,
 lower alkoxy(lower)alkylbenzimidazolyl or
 25 acylbenzimidazolyl;

8) subjecting a compound of the formula :


30



35

or its salt to elimination reaction of the N-protective group to provide a compound of the formula :

5



10

or its salt, in the above formulas,

R^1 , R^2 , R^3 , A, E and X are each as defined above,

Y_j is aryl which is substituted with protected amino and

15

nitro; or a condensed heterocyclic group which is

substituted with protected amino or lower alkyl

substituted with protected amino; and

Y_k is aryl which is substituted with amino and nitro; or

20


a condensed heterocyclic group which is substituted

with amino or lower alkyl substituted with amino;

or

9) subjecting a compound of the formula :

25




30

or its salt to deesterification reaction to provide a compound of the formula :

35

5



or its salt, in the above formulas,

R^1 , R^2 , R^3 , A, E and X are each as defined above,

10

Yl is aryl substituted with esterified carboxy, or a condensed heterocyclic group substituted with esterified carboxy, and


Ym is aryl substituted with carboxy, or a condensed heterocyclic group substituted with

15

carboxy, or

10) reacting a compound of the formula :

20




25

or its reactive derivative at the carboxy group

or a salt thereof with an amine or its salt to provide a compound of the formula :

30




35

or its salt, in the above formulas,


R^1 , R^2 , R^3 , A, E, X and Y_m are each as defined above,
and

Y_n is aryl or a condensed heterocyclic group, each of
which is substituted with substituted or
unsubstituted N-containing heterocycliccarbonyl,
carbamoyl, heterocycliccarbamoyl, or substituted or
unsubstituted lower alkylcarbamoyl; or

11) subjecting a compound of the formula :



or its salt to elimination reaction of methyl or the
hydroxy-protective group to provide a compound of the
formula :



or its salt, in the above formulas,

R^1 , R^2 , R^3 , A, E and X are each as defined above,


Y_o is a condensed (N-acyl)N-containing heterocyclic
group or a condensed heterocyclic group, each of
which is substituted with methoxy or lower alkyl
substituted with protected hydroxy; and

Y_p is a condensed (N-acyl)N-containing heterocyclic
group or a condensed heterocyclic group, each of
which is substituted with hydroxy or lower alkyl

substituted with hydroxy; or

12) reacting a compound of the formula :


5



10

or its salt with an acylating agent to provide a compound of the formula :

15




20

or its salt, in the above formulas,
 R^1 , R^2 , R^3 , A, E and X are each as defined above,
 Yq is a condensed heterocyclic group which is
substituted with amino or amino(lower)alkyl, and
 Yr is a condensed heterocyclic group which is
substituted with acylamino or
acylamino(lower)alkyl, or

25

13) reacting a compound of the formula :


30



35

or its salt with N-lower alkylmethyleammonium halide
to provide a compound of the formula :

5



10


or its salt, in the above formulas,
 R^1 , R^2 , R^3 , A , E , X and Y_a are each as defined above,
and

15

Y_s is indolyl which is substituted with methyl
substituted with lower alkylamino, or

14) subjecting a compound of the formula :


20



25

or its salt to oxidation reaction to provide a compound
of the formula :

30



35

or its salt, in the above formulas,

R^1 , R^2 , R^3 , A, E and X are each as defined above,

Yt is a condensed heterocyclic group which is

substituted with lower alkyl substituted with


5 hydroxy, and

Yu is a condensed heterocyclic group which is


substituted with lower alkyl substituted with

formyl, or

10 15) subjecting a compound of the formula :



or its salt to deesterification reaction to provide a
20 compound of the formula :



or its salt, in the above formulas,

A, E and X are each as defined above,

30 R_a^1 is aryl substituted with esterified carboxy or lower
alkoxy substituted with esterified carboxy,

R_b^1 is aryl substituted with carboxy or lower alkoxy
substituted with carboxy,


R_a^2 is lower alkyl,

35 R_a^3 is hydrogen or lower alkoxy, and

Yv is benzimidazolyl optionally substituted with lower alkyl or protected amino(lower)alkyl, or


16) subjecting a compound of the formula :

5



15 or its salt to elimination reaction of methyl substituted with aryl or substituted aryl to provide a compound of the formula :

20



25

or its salt, in the above formulas,

R_a^2 , R_a^3 , A, E, X and Y are each as defined above,


30 R_C^1 is aryl substituted with methoxy which is substituted with substituted or unsubstituted aryl, and R_d^1 is aryl substituted with hydroxy, or

17) reacting a compound of the formula :

35

310


5



10

or its reactive derivative at the carboxy group or a salt thereof with an amine or its salt to provide a compound of the formula :

15



20

or its salt, in the above formulas,
 R_b^1 , R_a^2 , R_a^3 , A, E, X and Yv are each as defined above,
 and


25

R_e^1 is aryl substituted with N-protected
 piperazinylcarbonyl, oxopiperidinylcarbonyl,
 carbamoyl, lower alkylcarbamoyl, lower
 alkylaminocarbamoyl or lower alkylamino(lower)-
 alkyl(N-lower)alkylcarbamoyl, or aryl which is
 substituted with lower alkoxy substituted with
 N-protected piperazinylcarbonyl,
 oxopiperidinylcarbonyl, carbamoyl, lower
 alkylcarbamoyl, lower alkylaminocarbamoyl or lower
 alkylamino(lower)alkyl(N-lower)alkylcarbamoyl,

30

35 18) reacting a compound of the formula :


5



10

or its salt with a reducing agent to provide a compound
of the formula :

15




20

or its salt, in the above formulas,
 R_a^2 , R_a^3 , A, E, X and Yv are each as defined above,
 R_f^1 is aryl which is substituted with lower alkoxy
substituted with oxopiperidinylcarbonyl, and
 R_g^1 is aryl which is substituted with lower alkoxy
substituted with hydroxypiperidinylcarbonyl, or

25

19) reacting a compound of the formula :


30



35


312

or its salt with an acylating agent to provide a compound of the formula :



10 or its salt, in the above formulas,
 R_d^1 , R_a^2 , R_a^3 , A, E, X and Y are each as defined above,
 and
 R_h^1 is aryl substituted with acyloxy, or


15 20) reacting a compound of the formula :



or its salt with a compound of the formula :

25 $Z^2 - R^6$ (V)

to provide a compound of the formula :



35

313

or its salt, in the above formulas,


R_d^1 , R_a^2 , R_a^3 , A, E, X and Y are each as defined above,

R_1^1 is aryl which is substituted with lower alkoxy
substituted with protected amino,

5 R^6 is lower alkyl substituted with protected amino, and
 Z^2 is an acid residue, or

21) subjecting a compound of the formula :


10



15

or its salt to elimination reaction of N-protective
group to provide a compound of the formula :

20



25

or its salt, in the above formulas,


R_1^1 , R_a^2 , R_a^3 , A, E, X and Y are each as defined above,
and

30 R_j^1 is aryl which is substituted with lower alkoxy
substituted with amino, or

22) reacting a compound of the formula :

35


5



10

or its salt with an acylating agent to provide a compound of the formula :

15




20

or its salt, in the above formulas,
 R_J^1 , R_a^2 , R_a^3 , A, E, X and Y are each as defined above,
 and
 R_K^1 is aryl which is substituted with acylamino, or

25

23) reacting a compound of the formula :

30




35

or its salt with lower alkanal in the presence of a reducing agent to provide a compound of the formula :

315

5



or its salt, in the above formulas,

10


R_j^1 , R_a^2 , R_a^3 , A, E, X and Y are each as defined above,
and

R_j^1 is aryl which is substituted with lower alkylamino,
or

15

24) subjecting a compound of the formula :


20



or its salt to reduction to provide a compound of the
formula :

25

30



or its salt, in the above formulas,

35


R_a^2 , R_a^3 , A, E, X and Y are each as defined above,

316

R_m^1 is aryl substituted with nitro, and
 R_n^1 is aryl substituted with amino, or


25) reacting a compound of the formula :

5



or its salt with an azide compound to provide a compound
 of the formula :

15



25

or its salt, in the above formulas,

R_n^1 , R_a^2 , R_a^3 , A, E, X and Yv are each as defined above,
 and


R_{ba}^1 is aryl substituted with carboxy, or

30

26) reacting a compound of the formula :

35


5



10

or its reactive derivative at the carboxy group or a salt thereof with a reducing agent to provide a compound of the formula :

15



20

or its salt, in the above formulas,

R_a^2 , R_a^3 , A, E, X and Yv are each as defined above,


R_{bb}^1 is aryl which is substituted with lower alkoxy substituted with carboxy, and

25

R_O^1 is aryl which is substituted with lower alkoxy substituted with hydroxymethyl, or


27) reacting a compound of the formula :

30



35

or its salt with an acylating agent to provide a compound of the formula :




10 or its salt, in the above formulas,


R_a^2 , R_a^3 , A, E, X and Y_v are each as defined above,
 R_p^1 is aryl which is substituted with lower alkoxy
 substituted with hydroxy, and

15 R_q^1 is aryl which is substituted with lower alkoxy
 substituted with acyloxy, or

28) reacting a compound of the formula :



or its salt with an alkali metal salt of phthalimide to provide a compound of the formula :




or its salt, in the above formulas,


R^1_Q , R^2_a , R^3_a , A, E, X and Yv are each as defined above,
and

R^1_r is aryl which is substituted with lower alkoxy
substituted with phthalimido, or

29) reacting a compound of the formula :



or its salt with an amine to provide a compound of the
formula :




or its salt, in the above formulas,


R^1 , R^2 , R^3 , A, E and X are each as defined above,
Yw is benzimidazolyl substituted with halogen, and
Yx is benzimidazolyl substituted with N-lower
alkylpiperidyl, morpholino, lower alkylamino,
di(lower)alkylaminopiperidino,
di(lower)alkylhydrazino,
amino(lower)alkyl(N-lower alkyl)amino or
di(lower)alkylamino(lower)alkylamino, or

326

30) subjecting a compound of the formula :




10 or its salt to elimination reaction of N-protective group to provide a compound of the formula :




20 or its salt, in the above formulas,
 R^1 , R^2 , R^3 , A, E and X are each as defined above,
 Yy is benzimidazolyl substituted with N-protected
 piperidyl, and
 25 Yz is benzimidazolyl substituted with piperidyl, or

31) reacting a compound of the formula :



or its salt with hydroxylamine or its salt to provide a compound of the formula :

5



10

or its salt, in the above formulas,

R^1 , R^2 , R^3 , A, E and X are each as defined above,

Y_1 is benzimidazolyl or indolyl, each of which is

15

substituted with formyl or cyano(lower)alkyl, and

Y_2 is benzimidazolyl or indolyl, each of which is


substituted with hydroxyiminomethyl or

amino(hydroxyimino)(lower)alkyl, or

20

32) reacting a compound of the formula :

25



30

or its reactive derivative at the carboxy group


or a salt thereof with a compound of the formula :



35

or its salt to provide a compound of the formula :

5



10

or its salt, in the above, formulas,
R¹, R², R³, X and Y are each as defined above.

15

7. A pharmaceutical composition comprising a compound of claim 1, as an active ingredient, in association with a pharmaceutically acceptable, substantially non-toxic carrier or excipient.

8. A compound of claim 1 for use as a medicament.

20

9. A method of therapeutic treatment and/or prevention of hypertension, heart failure, renal insufficiency, edema, ascites, vasopressin parasecretion syndrome, hepatocirrhosis, hyponatremia, hypokalemia, diabetic, circulation disorder, cerebrovascular disease, Meniere's syndrome or motion sickness which comprises administering an effective amount of a compound of claim 1 to human beings or animals.

25

30

10. Use of a compound of claim 1 for the manufacture of a medicament for treating and/or preventing hypertension, heart failure, renal insufficiency, edema, ascites, vasopressin parasecretion syndrome, hepatocirrhosis, hyponatremia, hypokalemia, diabetic, circulation disorder, cerebrovascular disease, Meniere's syndrome or motion sickness in human beings or animals.

35

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP 97/04192

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D235/08 A61K31/415 A61K31/40 C07D235/14 C07D235/30
C07D209/08 C07D209/42 C07D235/12 C07D235/24 C07D235/06
C07D235/10 C07D235/26 C07D209/12 C07D401/04 C07D401/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 91 05549 A (OTSUKA PHARMA CO LTD) 2 May 1991 cited in the application see pages 216-221; examples 120-127, 129, and 130 see page 343; example 374 see pages 479-480; examples 613 and 614 ---	1, 2, 7-10
X	EP 0 640 592 A (AMERICAN CYANAMID CO) 1 March 1995 see page 55; table I, the compounds of the examples 123 and 124 ---	1, 2, 7-10
X	EP 0 636 625 A (AMERICAN CYANAMID CO) 1 February 1995 see pages 188-189; table XIII, the compounds of the examples 484 and 485 ---	1, 2, 7-10
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

25 February 1998

Date of mailing of the international search report

03.04.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2

4032 Sassenheim, The Netherlands

Authorized officer

INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 97/04192

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 29152 A (FUJISAWA PHARMACEUTICAL CO ;SETOI HIROYUKI (JP); OHKAWA TAKEHIKO () 2 November 1995 cited in the application see page 233 - page 234; claim 1	1,7-10
Y	see the whole document	1-10
Y	WO 95 34540 A (OTSUKA PHARMA CO LTD ;OGAWA HIDENORI (JP); KONDO KAZUMI (JP); YAMA) 21 December 1995 see pages 634-639, claim 1; and in particular, page 636, lines 12-15 therein	1-10
X	DE 28 02 023 A (SANDOZ AG) 3 August 1978 see page 14 - page 15; examples 10,17	1-3,7,8
X	Y.G. PERRON ET AL.: "Derivatives of 6-Aminopenicillanic Acid. III. Reactions with N-Substituted Phthalamic Acids" JOURNAL OF MEDICINAL AND PHARMACEUTICAL CHEMISTRY., vol. 5, September 1962, EASTON US, pages 1016-1025, XP002056852 see page 1019; table II, compound no. 10	1,2
X	CHEMICAL ABSTRACTS, vol. 90, no. 21, 21 May 1979 Columbus, Ohio, US; abstract no. 168396r, M.T. NGUYEN: "Synthesis of indole ketoamides." page 583; column 1; XP002056856 see abstract; and Chemical Abstracts, CHEMICAL SUBSTANCES, 10th Collective Index, vol. 86-95, 1977-1981, page 4185CS, the compound with the RN: '69971-86-4! & TAP CHI HOA HOC, vol. 16, no. 1, 1978, pages 26-29,	1,2
X	T. YAMAZAKI ET AL.: "Dipolar Addition and Acyl Migration in Reactions of Benzyne and Acetylenes with 5-Membered Ring α -Diazo Ketones" TETRAHEDRON LETTERS., no. 44, October 1972, OXFORD GB, pages 4533-4536, XP002056853 see page 4535, the compound no. 31	1,2

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 97/04192

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CHEMICAL ABSTRACTS, vol. 108, no. 13, 28 March 1988 Columbus, Ohio, US; abstract no. 112144e, M.A.I. SALEM ET AL.: "Reaction of 3-carbethoxy-5,6-benzocoumarin with anthranilic acid and synthesis and some reactions of 3-(3',1'-benzoxazin-4'-one)-5,6-benzocouma rin." page 599; column 1; XP002056857 see abstract; and Chemical Abstracts, CHEMICAL SUBSTANCES, 12th Collective Index, vol. 106-115, 1987-1991, page 57433CS: the compound with the RN: '113120-43-7! & J. CHEM. SOC. PAK., vol. 9, no. 1, 1987, pages 177-189,</p>	1,2
X	<p>--- CHEMICAL ABSTRACTS, vol. 119, no. 21, 22 November 1993 Columbus, Ohio, US; abstract no. 225895w, M.R. SELIM ET AL.: "Synthesis of some benzoxazin-4-one derivatives and study of their reaction with nucleophilic reagents." page 1009; column 1; XP002056858 see abstract; and Chemical Abstracts, CHEMICAL SUBSTANCES, 13th Collective Index, vol. 116-125, 1992-1996, book 11, page 410: the compound with the RN: '150711-88-9! & J. INDIAN CHEM. SOC., vol. 69, no. 10, 1992, pages 688-690,</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1,2

INTERNATIONAL SEARCH REPORT

Inter. Patent Application No
PCT/JP 97/04192

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CHEMICAL ABSTRACTS, vol. 93, no. 1, 7 July 1990 Columbus, Ohio, US; abstract no. 8105d, M.M. ADDALLA ET AL.: "Synthesis of some 3-(3',1'-benzoxazin-4'-one)-6- substituted coumarins and their chemical reactions." page 745; column 1; XP002056859 see abstract; in particular the compounds of formula I, wherein R = H, and R1 = PhNH, 4-MeOC6H4NH, and 4-MeC6H4NH & EGYPT. J. CHEM., vol. 20, no. 3, 1979, pages 245-257,</p> <p style="text-align: center;">---</p>	1,2
X	<p>A. VARNAVAS ET AL.: "Anthranoyl-anthranilic acid: a template for the development of a new class of cholecystokinin receptor ligands" PHARMAZIE., vol. 51, no. 10, October 1996, BERLIN DD, pages 697-700, XP002056854 see page 698; table 1, the compounds no. 7 and 14</p> <p style="text-align: center;">---</p>	1,2
X	<p>M.-F. GRENIER-LOUSTALOT ET AL.: "Mechanics and Kinetics of Polymerization of Thermoplastic Polyimides. II. Study of "Bridged" Dianhydride/Aromatic Amine Systems" JOURNAL OF POLYMER SCIENCE, POLYMER CHEMISTRY EDITION., vol. 31, no. 12, November 1993, NEW YORK US, pages 3049-3063, XP002056855 see page 3050; the compound no. 5, wherein X = C=O</p> <p style="text-align: center;">-----</p>	1

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/JP 97/04192

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

Claims Nos. 1, 2, 5-16

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Claims 1 and 2 are so broad that for determining the scope of a meaningful international Search due account has been taken of Rule 33.3 PCT; special emphasis was put on the following subject-matter:
The compounds of claims 3-5, the methods of their preparation, pharmaceutical compositions containing them, and their use as medicaments

Remark : Although claim 9 is directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/JP 97/04192

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9105549 A	02-05-91	AU 630281 B	22-10-92
		AU 7291791 A	19-12-91
		CN 1051038 A,B	01-05-91
		CN 1107146 A	23-08-95
		DE 69026708 D	30-05-96
		EP 0450097 A	09-10-91
		ES 2089033 T	01-10-96
		JP 2052333 C	10-05-96
		JP 4154765 A	27-05-92
		JP 7076214 D	16-08-95
		US 5258516 A	02-11-93
		MX 9203571 A	01-07-92
		US 5559232 A	24-09-96
EP 0640592 A	01-03-95	AU 676737 B	20-03-97
		AU 6877691 A	09-02-95
		CA 2128911 A	30-01-95
		CN 1106832 D	15-08-95
		CZ 9401791 A	15-02-95
		FI 943542 A	30-01-95
		HU 71543 A	23-12-95
		JP 7179432 D	18-07-95
		NO 942311 A	30-01-95
		NZ 264111 A	26-11-96
		PL 304491 A	06-02-95
		SK 80391 A	10-05-95
		US 5512582 A	30-04-96
		US 5633441 A	11-11-97
		US 5693611 A	02-12-97
		ZA 940561 A	19-03-95
EP 0636625 A	01-02-95	AU 683611 A	10-11-97
		AU 6877691 A	09-02-95
		CA 2128911 A	30-01-95
		CN 1106831 D	15-08-95
		CZ 9401791 A	15-02-95
		FI 943542 A	30-01-95
		HU 71543 A	23-11-95
		JP 715711 A	02-06-95
		NO 942311 A	30-01-95

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 97/04192

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0636623		PL 304498 A	06-02-95
		SK 88194 A	12-04-95
		US 5516774 A	14-05-96
		US 5624923 A	29-04-97
		ZA 9405603 A	09-03-95
WO 9529152 A	02-11-95	AU 2267495 A	16-11-95
		EP 0757670 A	12-02-97
		JP 9512528 T	16-12-97
WO 953454 A	21-12-95	AU 2629395 A	05-01-96
		CA 2192928 A	21-12-95
		EP 0765314 A	02-04-97
		JP 8301848 A	19-11-96
DE 2802023	03-08-78	AU 3281078 A	02-08-79
		BE 863366 A	26-07-78
		CA 1105008 A	14-07-81
		DK 28278 A	29-07-78
		FI 780168 A	29-07-78
		FR 2378781 A	25-08-78
		GB 1595606 A	12-08-81
		JP 53095999 A	22-08-78
		NL 7800817 A	01-08-78
		SE 7800633 A	29-07-78
		US 4248871 A	03-02-81
		US 4180581 A	25-12-79
		ZA 7800521 A	26-09-79